


Forward-backward asymmetry in top-antitop production in proton-antiproton collisions


Regina Demina
University of Rochester
for
D0 collaboration
7/23/2011

Asymmetry in top-antitop production

- In early 80s asymmetry observed in $e^+e^- \rightarrow \mu^+\mu^-$ at $\sqrt{s}=34.6 \text{ GeV} \ll M_Z$ was used to verify the validity of EW theory (Phys. Rev. Lett. 48, 1701–1704 (1982))



- Similarly, asymmetry in $p\bar{p} \rightarrow t\bar{t}$ production could give information about new physics
 - Mediator with axial coupling in s-channel
 - Abnormally enhanced t-channel production
- Complications:
 - Top is not observed directly, but reconstructed through its decay products
 - Proton and antiproton are not point-like objects, lab frame is different from rest frame




Definitions

- Asymmetry defined for $ee \rightarrow \mu\mu$

$$A = \frac{N(\cos\theta > 0) - N(\cos\theta < 0)}{N(\cos\theta > 0) + N(\cos\theta < 0)}$$

- In proton-antiproton collisions $\theta \rightarrow y$
- Δy is invariant to boosts along z -axis
- Asymmetry based on Δy is the same in lab and tt rest frame
- Asymmetry based on rapidity of lepton from top decay
 - Lepton angles are measured with a good precision



$$\Delta y = y_t - y_{\bar{t}} = q_l (y_{leptonic} - y_{hadronic})$$

$$A = \frac{N(\Delta y > 0) - N(\Delta y < 0)}{N(\Delta y > 0) + N(\Delta y < 0)}$$

$$A_l = \frac{N(q_l y_l > 0) - N(q_l y_l < 0)}{N(q_l y_l > 0) + N(q_l y_l < 0)}$$

History of measurements and predictions

D0, reconstruction level

- PRL 100, 142002(2008) $A(0.9 fb^{-1}) = (12 \pm 8)\%$
- ICHEP2010 $A(4.3 fb^{-1}) = (8 \pm 4)\%$
 $A(MC @ NLO) = (0.8 \pm 1)\%$

CDF, generator level

- PRL 101, 202001(2008) $A(1.9 fb^{-1}) = (24 \pm 14)\%$
- Phys. Rev. D 83, 112003 (2011) $A(5.3 fb^{-1}) = (15.7 \pm 7.4)\%$
 $A(MC @ NLO) = (5.0 \pm 0.1)\%$

Reconstruction of top-antitop signal

Require :

→ 1 lepton with $p_T > 20 GeV$

→ $E_T > 20 GeV$

→ ≥ 4 jets with $p_T > 20 GeV$

→ leading jet with $p_T > 40 GeV$


→ ≥ 1 b - tag

→ In kinematic fit constrain

- $M_W = 80.4 GeV$

- $M_t = 172.5 GeV$


→ Charge of lepton determines
which reconstructed quark is top



1581 events pass the selection requirements in $5.4 fb^{-1}$


Asymmetry at reconstruction level

- Using kinematic variables of $l+jets$ events construct a discriminant and fit events with $\Delta y > 0$ and $\Delta y < 0$ for top fraction



$$A = (9.2 \pm 3.6^{+0.8}_{-0.9})\% \Leftrightarrow A(MC @ NLO) = (2.4 \pm 0.3^{+0.7}_{-0.5})\%$$


Asymmetry dependence on $M_{t\bar{t}}$ and $|\Delta y|$



$$A = (9.2 \pm 3.6_{-0.9}^{+0.8})\%$$

$$A(\text{MC@NLO}) = (2.4 \pm 0.3_{-0.5}^{+0.7})\%$$

Forward-Backward Top Asymmetry, %



Generated asymmetry

- “Unfolding” = correcting for acceptance (A) and detector resolution (S)
- **Method 1**: 4 bin Likelihood unfolding :

$$\vec{n}_{reco} = SA\vec{n}_{gen} \Rightarrow \vec{n}_{gen} = A^{-1}S^{-1}\vec{n}_{reco}$$

$$\Rightarrow A = (16.9 \pm 7.7^{+1.8}_{-2.6})\%$$

Problem with Method 1: migration of events near inner bin edge ($\Delta y \rightarrow 0$) is underestimated, while for the outer edge it is overestimated
Solution: *fine* bins closer to $\Delta y=0$


Problem: statistical fluctuations in data make the fine bin unfolding unstable

Solution: employ *regularization*

Bonus: reduced statistical uncertainties

Method 2: fine bin unfolding with regularization

$$A = (19.6 \pm 6.0^{+1.8}_{-2.6})\%$$



Results for asymmetry, in %

- Reconstruction level (experiments **cannot** be directly compared, only to Monte Carlo after reconstruction and selection)

- D0 (5.4 fb⁻¹) $9.2 \pm 3.6^{+0.8}_{-0.9}$

- MC@NLO (D0) $2.4 \pm 0.3^{+0.7}_{-0.5}$

- CDF (5.3 fb⁻¹) 7.5 ± 3.7

- MC@NLO (CDF) 2.4 ± 0.5

- Generator level


(experiments can be directly compared)

- D0 $19.6 \pm 6.0^{+1.8}_{-2.6}$

- CDF $15.8 \pm 7.2 \pm 1.7$


- MC@NLO 5.0 ± 0.1

Forward-Backward Top Asymmetry, %



Lepton-based asymmetry, in %

- Since lepton direction is defined with a very good precision, lepton based asymmetry is simpler to extract
- Lepton from top decay carries information about underlying asymmetry at production
- Can be directly compared to theoretical predictions



Reconstruction level

$$A_l = 14.2 \pm 3.7 \pm 0.7$$

$$A_l(MC@NLO) = 0.8 \pm 0.3 \pm 0.5$$


Generated level

$$A_l = 15.2 \pm 3.8^{+1.0}_{-1.3}$$

$$A_l(MC@NLO) = 2.1 \pm 0.1$$


Interpretation of the Asymmetry

- Coulomb repulsion
- QED: $e^+ \mu^+$
- QCD: quark-top



- Coulomb attraction
- QED: $e^- \mu^+$
- QCD: antiquarkquark-top


Predicted asymmetry in SM



$l+4 \text{ jets}$

data : $12.2 \pm 4.2\%$

MC@NLO : $3.9 \pm 0.3\%$



$l+ \geq 5 \text{ jets}$

data : $-3.0 \pm 7.8\%$

MC@NLO : $-2.9 \pm 0.7\%$



Born(α_s^2) and box(α_s^4)

- Coulomb-like repulsion of top and quark and attraction of antitop and quark in QCD
- Interference – α_s^3
- Positive asymmetry
- Final state with no extra partons \rightarrow small transverse momentum of the $t\bar{t}$ system

ISR (α_s^3) and FSR(α_s^3)


- Interference – α_s^3
- Negative asymmetry
- Final state with extra gluons \rightarrow large transverse momentum of the $t\bar{t}$ system
- Possible extra jets

Modeling of gluon radiation



- $p_T^{t\bar{t}}$ spectrum suggests that gluon radiation might be mismodeled by MC@NLO+HERWIG
- lower radiation is preferred
- best agreement with PYTHIA ISR off
- This suggests a higher contribution from $2 \rightarrow 2$ processes, e.g. Born+box

Asymmetry and gluon radiation



- MC@NLO+HERWIG suggests strong dependence of asymmetry on $p_T^{t\bar{t}}$
- Some PYTHIA tunes suggest even more dramatic dependence while other do not – the main parameter that affects this behavior is angular coherence of ISR
- Asymmetry dependence on $p_T^{t\bar{t}}$ is a source of systematic uncertainty on the measured value of asymmetry
- Higher weight of $2 \rightarrow 2$ processes (Born+box) would shift the predicted asymmetry toward more positive and higher values

Conclusions

- Using 5.4 fb^{-1} of data D0 measured asymmetry in top-antitop production

$$A = (19.6 \pm 6.0_{-2.6}^{+1.8})\%$$


$$A(MC@NLO) = (5.0 \pm 0.1)\%$$

- Asymmetry in leptons from top decay is

$$A_l = (15.2 \pm 3.8_{-1.3}^{+1.0})\%$$

$$A_l(MC@NLO) = (2.1 \pm 0.1)\%$$

Forward-Backward Top Asymmetry, %



Systematics on A

TABLE VII. Systematic uncertainties on A_{FB} .

Source	Absolute uncertainty ^a (%)		
	Reconstruction level		Prod. level
	Prediction	Measurement	Measurement
Jet reco	± 0.3	± 0.5	± 1.0
JES/JER	+0.5	-0.5	-1.3
Signal modeling	± 0.3	± 0.5	+0.3/-1.6
<i>b</i> -tagging	-	± 0.1	± 0.1
Charge ID	-	+0.1	+0.2/-0.1
Bg subtraction	-	± 0.1	+0.8/-0.7
Unfolding Bias	-	-	+1.1/-1.0
Total	+0.7/-0.5	+0.8/-0.9	+1.8/-2.6

Systematics on A_1

TABLE VIII. Systematic uncertainties on A_{FB}^l .

Source	Absolute uncertainty ^a (%)		
	Reconstruction level		Prod. level
	Prediction	Measurement	Measurement
Jet reco	± 0.3	± 0.1	± 0.8
JES/JER	+0.1	-0.4	+0.1/-0.6
Signal modeling	± 0.3	± 0.5	+0.2/-0.6
b -tagging	-	± 0.1	± 0.1
Charge ID	-	+0.1	+0.2/-0.0
Bg subtraction	-	± 0.3	± 0.6
Total	± 0.5	± 0.7	+1.0/-1.3

Migration matrix

