



Theoretical update of B -mixing



Alexander Lenz

CERN, Theory Division

TeVatron gave us many presents, and then ...





Mixing I

Time evolution of a decaying particle: $B(t) = \exp [-im_B t - \Gamma_B / 2t]$
can be written as

$$i \frac{d}{dt} \begin{pmatrix} |B(t)\rangle \\ |\bar{B}(t)\rangle \end{pmatrix} = \left(\hat{M} - \frac{i}{2} \hat{\Gamma} \right) \begin{pmatrix} |B(t)\rangle \\ |\bar{B}(t)\rangle \end{pmatrix}$$

BUT: In the neutral B -system transitions like $B_{d,s} \rightarrow \bar{B}_{d,s}$ are possible due to weak interaction: **Box diagrams**





Mixing II

Mixing is a macroscopic quantum effect!

It was observed in

- K^0 -system: 1950s (see text books, regeneration...)
- B_d -system: 1986
- B_s -system: 2006
- D^0 -system: 2007

Strongly suppressed in the SM (due to virtual top-quarks)
New physics effects might be of comparable size


?Is QCD under control?

Mixing III

Time evolution of a decaying particle: $B(t) = \exp[-im_B t - \Gamma_B/2t]$
can be written as

$$i \frac{d}{dt} \begin{pmatrix} |B(t)\rangle \\ |\bar{B}(t)\rangle \end{pmatrix} = \left(\hat{M} - \frac{i}{2} \hat{\Gamma} \right) \begin{pmatrix} |B(t)\rangle \\ |\bar{B}(t)\rangle \end{pmatrix}$$

BUT: In the neutral B -system transitions like $B_{d,s} \rightarrow \bar{B}_{d,s}$ are possible due to weak interaction: Box diagrams



\Rightarrow off-diagonal elements in \hat{M} , $\hat{\Gamma}$: M_{12} , Γ_{12} (complex)

Diagonalization of \hat{M} , $\hat{\Gamma}$ gives the physical eigenstates B_H and B_L with the masses M_H , M_L and the decay rates Γ_H , Γ_L

CP-odd: $B_H := p B + q \bar{B}$, CP-even: $B_L := p B - q \bar{B}$ with $|p|^2 + |q|^2 = 1$

Mixing IV

$|M_{12}|$, $|\Gamma_{12}|$ and $\phi = \arg(-M_{12}/\Gamma_{12})$ can be related to three observables:

- Mass difference: $\Delta M := M_H - M_L = 2|M_{12}| \left(1 - \frac{1}{8} \frac{|\Gamma_{12}|^2}{|M_{12}|^2} \sin^2 \phi + \dots \right)$

$|M_{12}|$: heavy internal particles: t, SUSY, ...

- Decay rate difference: $\Delta\Gamma := \Gamma_L - \Gamma_H = 2|\Gamma_{12}| \cos \phi \left(1 + \frac{1}{8} \frac{|\Gamma_{12}|^2}{|M_{12}|^2} \sin^2 \phi + \dots \right)$

$|\Gamma_{12}|$: light internal particles: u, c, ... (almost) no NP!!!

- Flavor specific/semileptonic CP asymmetries:

$\bar{B}_q \rightarrow f$ and $B_q \rightarrow \bar{f}$ forbidden

No direct CP violation: $|\langle f | B_q \rangle| = |\langle \bar{f} | \bar{B}_q \rangle|$

e.g. $B_s \rightarrow D_s^- \pi^+$ or $B_q \rightarrow X l \nu$ (semileptonic)

$$a_{sl} \equiv a_{fs} = \frac{\Gamma(\bar{B}_q(t) \rightarrow f) - \Gamma(B_q(t) \rightarrow \bar{f})}{\Gamma(\bar{B}_q(t) \rightarrow f) + \Gamma(B_q(t) \rightarrow \bar{f})} = -2 \left(\left| \frac{q}{p} \right| - 1 \right) = \text{Im} \frac{\Gamma_{12}}{M_{12}} = \frac{\Delta\Gamma}{\Delta M} \tan \phi$$



The Mass Difference ΔM

Calculating the box diagram with an internal top-quark yields

$$M_{12,q} = \frac{G_F^2}{12\pi^2} (V_{tq}^* V_{tb})^2 M_W^2 S_o(x_t) B_{B_q} f_{B_q}^2 M_{B_q} \hat{\eta}_B$$

(Inami, Lim '81)

- Hadronic matrix element: $\frac{8}{3} B_{B_q} f_{B_q}^2 M_{B_q} = \langle \bar{B}_q | (\bar{b}q)_{V-A} (\bar{b}q)_{V-A} | B_q \rangle$
- Perturbative QCD corrections $\hat{\eta}_B$ (Buras, Jamin, Weisz, '90)

Theory 1102.4274 vs. Experiment : HFAG 11

$$\Delta M_d = 0.543 \pm 0.091 \text{ } ps^{-1}$$

$$\Delta M_d = 0.507 \pm 0.004 \text{ } ps^{-1}$$

ALEPH, CDF, D0, DELPHI, L3,

OPAL, BABAR, BELLE, ARGUS, CLEO

$$\Delta M_s = 17.30 \pm 2.6 \text{ } ps^{-1}$$

$$\Delta M_s = 17.70 \pm 0.12 \text{ } ps^{-1}$$

CDF, D0, LHCb

Important bounds on the unitarity triangle and new physics

Determination of Γ_{12}

Sensitive to real intermediate states \Rightarrow much more complicated than M_{12}

1. OPE I: Integrate out W: like $M_{12} \propto f_B^2 B$
2. OPE II: Heavy quark expansion $\Rightarrow \Gamma_i^{(j)} \propto f_B^2 \sum C_k B_K$

$$\Gamma_{12} = \left(\frac{\Lambda}{m_b}\right)^3 \left(\Gamma_3^{(0)} + \frac{\alpha_s}{4\pi} \Gamma_3^{(1)} + \dots \right) + \left(\frac{\Lambda}{m_b}\right)^4 \left(\Gamma_4^{(0)} + \dots \right) + \left(\frac{\Lambda}{m_b}\right)^5 \left(\Gamma_5^{(0)} + \dots \right) + \dots$$

1996: Beneke, Buchalla, Dunietz

1998: Beneke, Buchalla, Greub, A.L., Nierste

2003: Ciuchini, Franco, Lubicz, Mescia, Tarantino; Beneke, Buchalla, A.L., Nierste

2006: A.L., Nierste

2007: Badin, Gabbiani, Petrov

$$\begin{aligned}\Delta\Gamma_s &= \Delta\Gamma_s^0 (1 + \delta^{\text{Lattice}} + \delta^{\text{QCD}} + \delta^{\text{HQE}}) \\ &= 0.142 \text{ ps}^{-1} (1 - 0.14 - 0.06 - 0.19)\end{aligned}$$



HQE under attack!

OPE II might be questionable - relies on quark hadron duality

- Mid 90's: Missing Charm puzzle $n_c^{\text{Exp.}} < n_c^{\text{SM}}$, semi leptonic branching ratio
- Mid 90's: Λ_b lifetime is too short
- before 2003: $\tau_{B_s}/\tau_{B_d} \approx 0.94 \neq 1$
- 2010/2011: Di-muon asymmetry too large

Theory arguments for HQE

- ⇒ calculate corrections in all possible “directions”, to test convergence
⇒ Γ_{12} seems to be ok!
- ⇒ test reliability of OPE II via lifetimes (no NP effects expected) “directions”, to test convergence
⇒ $\tau(B^+)/\tau(B_d)$ Experiment and theory agree within hadronic uncertainties



HQE under attack!

OPE II might be questionable - relies on quark hadron duality

- 2012: $n_c^{2011\text{PDG}} = 1.20 \pm 0.06$ vs. $n_c^{\text{SM}} = 1.20 \pm 0.04$
Eberhardt, Krinner, A.L., Rauh in prep.
- Mid 90's: Λ_b lifetime is too short
- before 2003: $\tau_{B_s}/\tau_{B_d} \approx 0.94 \neq 1$
- 2010/2011: Di-muon asymmetry too large

Theory arguments for HQE

- ⇒ calculate corrections in all possible “directions”, to test convergence
⇒ Γ_{12} seems to be ok!
- ⇒ test reliability of OPE II via lifetimes (no NP effects expected) “directions”, to test convergence
⇒ $\tau(B^+)/\tau(B_d)$ Experiment and theory agree within hadronic uncertainties



HQE under attack!

OPE II might be questionable - relies on quark hadron duality

- 2012: $n_c^{2011\text{PDG}} = 1.20 \pm 0.06$ vs. $n_c^{\text{SM}} = 1.20 \pm 0.04$
Eberhardt, Krinner, A.L., Rauh in prep.
- HFAG '03 $\tau_{\Lambda_b} = 1.212 \pm 0.052 \text{ ps}^{-1}$ → HFAG '11 $\tau_{\Lambda_b} = 1.425 \pm 0.032 \text{ ps}^{-1}$
Shift by 4σ ⇒ Eagerly waiting for new LHCb results!!!
- before 2003: $\tau_{B_s}/\tau_{B_d} \approx 0.94 \neq 1$
- 2010/2011: Di-muon asymmetry too large

Theory arguments for HQE

- ⇒ calculate corrections in all possible “directions”, to test convergence
⇒ Γ_{12} seems to be ok!
- ⇒ test reliability of OPE II via lifetimes (no NP effects expected) “directions”, to test convergence
⇒ $\tau(B^+)/\tau(B_d)$ Experiment and theory agree within hadronic uncertainties



HQE under attack!

OPE II might be questionable - relies on quark hadron duality

- 2012: $n_c^{2011\text{PDG}} = 1.20 \pm 0.06$ vs. $n_c^{\text{SM}} = 1.20 \pm 0.04$
Eberhardt, Krinner, A.L., Rauh in prep.
- HFAG '03 $\tau_{\Lambda_b} = 1.212 \pm 0.052 \text{ ps}^{-1}$ → HFAG '11 $\tau_{\Lambda_b} = 1.425 \pm 0.032 \text{ ps}^{-1}$
Shift by 4σ ⇒ Eagerly waiting for new LHCb results!!!
- Moriond 2012 LHCb: $\tau_{B_s}/\tau_{B_d} = 1.001 \pm 0.014$ Talk by Peter Clarke
- 2010/2011: Di-muon asymmetry too large

Theory arguments for HQE

- ⇒ calculate corrections in all possible “directions”, to test convergence
⇒ Γ_{12} seems to be ok!
- ⇒ test reliability of OPE II via lifetimes (no NP effects expected) “directions”, to test convergence
⇒ $\tau(B^+)/\tau(B_d)$ Experiment and theory agree within hadronic uncertainties



The B_s lifetime

Moriond 2012 LHCb vs SM A.L., Nierste 2011

$$\frac{\tau_{B_s}^{\text{Exp}}}{\tau_{B_d}} = 1.001 \pm 0.014 \quad \frac{\tau_{B_s}^{\text{SM}}}{\tau_{B_d}} = 0.996 \dots 1.000$$

- 0.940 ± 0.014 would have been a desaster for SM = may be NP :-)
- Update of effective lifetimes
Fleischer et al used 1011.1096, 1109.1112, 1109.5115: $\tau_{B_s} = 1.477$ ps

	Exp.	SM-old	SM-new
$\tau^{\text{Eff}}(K^+ K^-)$	1.44 ± 0.10	1.390 ± 0.032	1.43 ± 0.03
$\tau^{\text{Eff}}(\psi f_0)$	1.70 ± 0.12	1.582 ± 0.036	1.63 ± 0.03
τ^{FS}	1.417 ± 0.042	— — —	1.54 ± 0.03



HQE under attack!

OPE II might be questionable - relies on quark hadron duality

- 2012: $n_c^{2011\text{PDG}} = 1.20 \pm 0.06$ vs. $n_c^{\text{SM}} = 1.20 \pm 0.04$
Eberhardt, Krinner, A.L., Rauh in prep.
- HFAG '03 $\tau_{\Lambda_b} = 1.212 \pm 0.052 \text{ ps}^{-1}$ → HFAG '11 $\tau_{\Lambda_b} = 1.425 \pm 0.032 \text{ ps}^{-1}$
Shift by 4σ ⇒ Eagerly waiting for new LHCb results!!!
- Moriond 2012 LHCb: $\tau_{B_s}/\tau_{B_d} = 1.001 \pm 0.014$ Talk by Peter Clarke
- 2010/2011: Di-muon asymmetry too large — Test Γ_{12} with $\Delta\Gamma_s$!

Theory arguments for HQE

- ⇒ calculate corrections in all possible “directions”, to test convergence
⇒ Γ_{12} seems to be ok!
- ⇒ test reliability of OPE II via lifetimes (no NP effects expected) “directions”, to test convergence
⇒ $\tau(B^+)/\tau(B_d)$ Experiment and theory agree within hadronic uncertainties



$\Delta\Gamma_s$ in NLO-QCD I

A brief history of theory predictions

'81... Hagelin; Buras et al.;...	$\Delta\Gamma \propto \mathcal{O}(0.15 \text{ ps}^{-1})$
'93 Aleksan et al.; ...	$\Delta\Gamma \propto \mathcal{O}(0.10 \text{ ps}^{-1})$
'96 Beneke, Buchalla, Dunietz	$\Delta\Gamma_s = (0.11^{+0.07}_{-0.06}) \text{ ps}^{-1}$
'00 Beneke, A.L.	$\Delta\Gamma_s = (0.06 \pm 0.03) \text{ ps}^{-1}$
'03 Ciuchini, et al	$\Delta\Gamma_s = (0.050 \pm 0.016) \text{ ps}^{-1}$
'06 A.L., Nierste	$\Delta\Gamma_s = (0.096 \pm 0.036) \text{ ps}^{-1}$
'11 A.L., Nierste	$\Delta\Gamma_s = (0.087 \pm 0.021) \text{ ps}^{-1}$

Crucial dependence on non-perturbative parameters!

2011 $f_{B_s} = 231 \pm 15 \text{ MeV}$ used.

Newer Results:


- 1110.4510 - HPQCD: $f_{B_s} = 225 \pm 4 \text{ MeV} \Rightarrow \Delta\Gamma_s = (0.083 \pm 0.017) \text{ ps}^{-1}$
- 1112.3051 - Fermilab: $f_{B_s} = 242 \pm 9.5 \text{ MeV} \Rightarrow \Delta\Gamma_s = (0.095 \pm 0.021) \text{ ps}^{-1}$
- 1201.3956 - chiral QM: $f_{B_s} = 262 \pm ? \text{ MeV} \Rightarrow \Delta\Gamma_s = (0.112 \pm ?) \text{ ps}^{-1}$



$\Delta\Gamma_s$ in NLO-QCD II

Improvement in theoretical accuracy

$\Delta\Gamma_s^{\text{SM}}$	2011	2006
Central Value	0.087 ps^{-1}	0.096 ps^{-1}
$\delta(\mathcal{B}_{\tilde{R}_2})$	17.2%	15.7%
$\delta(f_{B_s})$	13.2%	33.4%
$\delta(\mu)$	7.8%	13.7%
$\delta(\tilde{\mathcal{B}}_{S, B_s})$	4.8%	3.1%
$\delta(\mathcal{B}_{R_0})$	3.4%	3.0%
$\delta(V_{cb})$	3.4%	4.9%
$\delta(\mathcal{B}_{B_s})$	2.7%	6.6%
...
$\sum \delta$	24.5%	40.5%



Finally $\Delta\Gamma_s$ is measured! (naive: 6.1σ)

$$\Delta\Gamma_s^{\text{SM}} = (0.087 \pm 0.021) \text{ ps}^{-1}$$

LHCb from $B_s \rightarrow J/\psi\phi$

$$\text{LP 2011 } \Delta\Gamma_s = (0.123 \pm 0.031) \text{ ps}^{-1} \Rightarrow \frac{\Delta\Gamma_s^{\text{Exp}}}{\Delta\Gamma_s^{\text{SM}}} = 1.41 \pm 0.50$$

$$\text{Moriond 2012 } \Delta\Gamma_s = (0.116 \pm 0.019) \text{ ps}^{-1} \Rightarrow \frac{\Delta\Gamma_s^{\text{Exp}}}{\Delta\Gamma_s^{\text{SM}}} = 1.33 \pm 0.39$$

- D0 8fb^{-1} 1109.3166: $\Delta\Gamma_s = (0.163 \pm 0.065) \text{ ps}^{-1}$
- CDF 9.6fb^{-1} : Talk by G. Borissov $\Delta\Gamma_s < \Delta\Gamma_s^{\text{SM}}$



Finally $\Delta\Gamma_s$ is measured! (naive: 6.1σ)

Get rid off the dependence on f_{B_s} (No NP in ΔM)

$$\begin{aligned}\frac{\Delta\Gamma_s}{\Delta M_s} &= 10^{-4} \cdot \left[46.2 + 10.6 \frac{\tilde{B}'_S}{B} - \left(13.2 \frac{B_{\tilde{R}_2}}{B} - 2.5 \frac{B_{R_0}}{B} + 1.2 \frac{B_R}{B} \right) \right] \\ &= 0.0050 \pm 0.0010\end{aligned}$$

HQE vs. Experiment

$$\left(\frac{\Delta\Gamma_s}{\Delta M_s} \right)^{\text{Exp}} / \left(\frac{\Delta\Gamma_s}{\Delta M_s} \right)^{\text{SM}} = 1.30 \pm 0.34$$

HQE works also for Γ_{12} !

How precise does it work? 30%? 10%?

Still more accurate data needed! **TeVatron, LHCb, Super-B(elle)**


$$\Delta\Gamma_s^{\text{CP}}/\Gamma_s = 2Br(B_s \rightarrow D_s^{(*)+} + D_s^{(*)-})?$$

- 1993 Aleksan; Le Yaouanc, Olivre, Pene, Raynal:
The above equation holds in the limit: $m_c \rightarrow \infty$; $m_b - 2m_c \rightarrow 0$; $N_c \rightarrow \infty$
Corresponds to negligible 3-body final state contributions to Γ_{12}^s

$$\frac{\Delta\Gamma_s}{\Gamma_s} \propto \mathcal{O}(0.15)$$

- 1107.4325 Chua, Hou, Shen Reanalysis of the exclusive approach
 - ◆ 2-body final states contribute 0.100 ± 0.030 to $\Delta\Gamma/\Gamma$
Aleksan et al were lucky...
 - ◆ 3-body final states contribute about $0.06...0.08$
This is comparable to 2-body final states! \Rightarrow bad approximation \Rightarrow test exp.

We strongly discourage from the inclusion of $Br(B_s \rightarrow D_s^{(*)+} + D_s^{(*)-})$ in averages with $\Delta\Gamma_s$ determined from clean methods.

A.L., Nierste; hep-ph/0612167

Semi leptonic CP-asymmetries a_{fs} and $\Delta\Gamma_d$

SM predictions: A.L., U. Nierste, 1102.4274; A.L. 1108.1218

$$\begin{aligned} a_{fs}^s &= (1.9 \pm 0.3) \cdot 10^{-5} & \phi_s &= 0.22^\circ \pm 0.06^\circ \\ a_{fs}^d &= -(4.1 \pm 0.6) \cdot 10^{-4} & \phi_d &= -4.3^\circ \pm 1.4^\circ \\ A_{sl}^b &= 0.406a_{sl}^s + 0.594a_{sl}^d & = & (-2.3 \pm 0.4) \cdot 10^{-4} \end{aligned}$$

CP

Experimental bounds

$$\begin{aligned} a_{fs}^s &= (-1150 \pm 610) \cdot 10^{-5} \text{ (HFAG 11)} \\ \phi_s &= -51.6^\circ \pm 12^\circ \quad (\text{A.L., Nierste, CKMfitter, 1008.1593}) \\ &= -0.1^\circ \pm 5.0^\circ \quad \text{LHCb Moriond 2012} \\ a_{fs}^d &= -(49 \pm 38) \cdot 10^{-4} \quad \text{ (HFAG 11)} \\ \frac{\Delta\Gamma_d}{\Gamma_d} &= (-17 \pm 21) \cdot 10^{-3} \quad \text{ (Belle EPS 2011)} \\ A_{sl}^b &= -(7.87 \pm 1.72 \pm 0.93) \cdot 10^{-3} \text{ (D0, 1106.6308)} \end{aligned}$$



$$A_{sl}^b(\text{Exp.})/A_{sl}^b(\text{Theory}) = 34 \quad 3.9 - \sigma\text{-effect}$$

New Physics in B-Mixing I

$$\Gamma_{12,s} = \Gamma_{12,s}^{\text{SM}}, \quad M_{12,s} = M_{12,s}^{\text{SM}} \cdot \Delta_s; \quad \Delta_s = |\Delta_s| e^{i\phi_s^\Delta}$$

$$\Delta_s = r_s^2 e^{2i\theta_s} = C_{B_s} e^{2i\phi_{B_s}} = 1 + h_s e^{2i\sigma_s}$$

$$\Delta M_s = 2|M_{12,s}^{\text{SM}}| \cdot |\Delta_s|$$


$$\Delta \Gamma_s = 2|\Gamma_{12,s}| \cdot \cos(\phi_s^{\text{SM}} + \phi_s^\Delta)$$

$$\frac{\Delta \Gamma_s}{\Delta M_s} = \frac{|\Gamma_{12,s}|}{|M_{12,s}^{\text{SM}}|} \cdot \frac{\cos(\phi_s^{\text{SM}} + \phi_s^\Delta)}{|\Delta_s|}$$

$$a_{fs}^s = \frac{|\Gamma_{12,s}|}{|M_{12,s}^{\text{SM}}|} \cdot \frac{\sin(\phi_s^{\text{SM}} + \phi_s^\Delta)}{|\Delta_s|}$$



$$\sin(\phi_s^{\text{SM}}) \approx 1/240$$

For $|\Delta_s| = 0.9$ and $\phi_s^\Delta = -\pi/4$ one gets the following bounds in the **complex Δ -plane**:



New Physics in B-Mixing II

Combine all data till end of 2011 and neglect penguins
 fit of Δ_d and Δ_s 1203.0238 (update of 1008.1593) soon v2!



- Fits not so good anymore (LHCb vs. Dzero)
- $B \rightarrow \tau\nu$ vs. $\sin 2\beta$ solved with ϕ_d^Δ — No tension for ϵ_K



The dimuon asymmetry

The central value of the di μ asymmetry is larger than *theoretically possible!*

$$\begin{aligned} A_{sl}^{Max.} &\approx (0.594 \pm 0.022)(5.4 \pm 1.0) \cdot 10^{-3} \frac{\sin(\phi_d^{SM} + \phi_d^\Delta)}{|\Delta_d|} \\ &\quad + (0.406 \pm 0.022)(5.0 \pm 1.1) \cdot 10^{-3} \frac{\sin(\phi_s^{SM} + \phi_s^\Delta)}{|\Delta_s|} \\ &\approx (-3.1; -4.8[1\sigma]; -9.0[3\sigma]) \cdot 10^{-3} \\ A_{sl}^{D0} &= (-7.8 \pm 2.0) \cdot 10^{-3} && \text{A.L. 1108.1218} \end{aligned}$$

Possible solutions:


- HQE violated by $\mathcal{O}(200\% - 3300\%)$ now excluded!
- Huge new physics in Γ_{12} ? - see talk by Uli Haisch
- Contradiction to $B_s \rightarrow J/\psi \phi$ from LHCb? - Penguins
- Stat. fluctuation (1.5σ) of the D0 result? (Actual value is below -4.8 per mille?)
Independent measurements of semi leptonic asymmetries needed!

?New physics in Γ_{12} ?


- Large ($\mathcal{O}(200 - 3400\%)$) NP effects in Γ_{12} ?

Why not seen somewhere else?

A new operator $bs \rightarrow X$ with $M_x < M_B$ contributes not only to a_{sl}^s but also to many more observables, e.g.:



$$\Gamma_3 \Rightarrow \begin{cases} \tau(B_s)/\tau(B_d) \\ \Delta\Gamma_s \end{cases}$$



$$\Gamma_0 \Rightarrow \begin{cases} \tau(B_x) \\ B_{sl} \\ Br(b \rightarrow s \text{ no charm}) \end{cases}$$

◆ ...

◆ A promising candidate for X seems to be $\tau^+ + \tau^- \rightarrow$ Uli Haisch.



?New physics in Γ_{12} ?

■ Missing charm puzzle, e.g.

Bigi et al '94; Bagan et al. '94; Falk, Wise, Dunietz '95, Neubert '97... A.L.
,hep-ph/0011258

Look at inclusive b -decay into 0, 1, 2 c -quarks

Define $r(x \text{ charm}) := \frac{\Gamma(b \rightarrow x \text{ charm})}{\Gamma_{sl}}$: $m_b^5 V_{cb}^2$ cancels; Γ_{sl} seems safe

The average number of charm quarks per b -decay reads

$$\begin{aligned} n_c &= 0 + [r(1c) + 2r(2c)] B_{sl}^{\text{Exp.}} \\ &= 1 + [r(2c) - r(0c)] B_{sl}^{\text{Exp.}} \\ &= 2 - [r(1c) + 2r(0c)] B_{sl}^{\text{Exp.}} \end{aligned}$$


Buchalla, Dunietz, Yamamoto '95

- ◆ $n_c^{\text{Exp.}} < n_c^{\text{Theory}}$ = missing charm puzzle
May be enhanced $b \rightarrow s g \dots$ Kagan ...
- ◆ latest Data from BaBar and CLEO agree within large uncertainties
Recent and future experiments can do better!
- ◆ Any unknown, even invisible decay mode has an effect on $r(0, 1, 2 \text{ charm})$

!!! \Rightarrow Need new experimental values for $r(0c, 1c, 2c) = \Gamma_{0c,1c,2c}/\Gamma_{sl}$ and B_{sl} !!!

?New physics in Γ_{12} ?

Step I: Forget about all the bounds and fit $\Delta\Gamma$, a_{sl} and ΔM :



Step II: Take your favourite model which gives new contributions to Γ_{12}

- Determine contributions to δ_d , δ_s
- Determine contributions to τ_{B_s} , n_c ,
- *Exclude the model :-)*



How large are Penguins?

Angular analysis of $B_s \rightarrow J/\psi\phi$ at CDF, D0 and LHCb:

$$S_{\psi\phi}^{\text{SM}} = 0.0036 \pm 0.002 \rightarrow \sin(2\beta_s - \phi_s^\Delta - \delta_s^{\text{Peng,SM}} - \delta_s^{\text{Peng,NP}}) = 0.002 \pm 0.087$$

LHCb Moriond 2012

Is this a contraction to the dimuon asymmetry?

Depends on the possible size of penguin contributions

- SM penguins are expected to be very small
but see also Faller, Fleischer; Mannel 2008
- NP penguins might be larger

But: even small penguin contributions have a sizeable effect! A.L. 1106.3200



Wish-list for Experiments

a) Congratulations to LHCb for the first measurement of $\Delta\Gamma_s$!

- ◆ Still more precision needed: LHCb, TeVatron, Super-B $B_s \rightarrow J/\psi\eta^{(')}$
- ◆ Do not use $Br(B_s \rightarrow D_s^{(*)+} D_s^{(*)-}) = \frac{\Delta\Gamma_s^{\text{CP}}}{2\Gamma}$ - check size of 3-body FS!

b) $\tau_{B_s} = (1.001 \pm 0.014)\tau_{B_d}$: strong constraint on NP and duality violation

- ◆ Combine with other determinations of τ_{B_s} : LHCb, ATLAS?, CMS?
- ◆ B_s : Effective lifetimes, flavor specific lifetimes (2.x sigma deviation)
- ◆ τ_{Λ_b}, \dots

c) Di muon asymmetry A_{sl}^b

- ◆ HQE fails? **No! At most** 30 – 40% — more precise test via $\tau(B_s), \Delta\Gamma_s, \dots$
- ◆ NP acts in Γ_{12} ? **No! At most** 40%! — More precise tests via
 $\tau(B_s), \Delta\Gamma_s, \Delta\Gamma_d, n_c, B_{sl}, r(0, 1, 2 \text{ charm}), B_s \rightarrow \tau\tau, B \rightarrow K\tau\tau, \dots$
- ◆ ???
- ◆ **Experimental cross-check via** a_{sl}^d **and** a_{sl}^s !

d) $\phi_s^{LHCb} \ll \phi_s^{A_{sl}^b}$ How large is the penguin pollution?

- ◆ Even small penguins can be important!
- ◆ **Values** for many penguin modes e.g. $B_s \rightarrow J/\psi K_s, K^0 \bar{K}^0, \phi\phi, \eta^{(')}\eta^{(')} \dots$



What to do list - Theory

Test of HQE with lifetimes

- τ_{B^+}/τ_{B_d} and τ_{B_s}/τ_{B_d} fits well \Rightarrow currently no hints for deviations from HQE
- Precise non-perturbative matrix elements for 4-quark operators urgently needed
Beautiful Mesons and Baryons on the Lattice ECT* Trento, 2-6 April 2012
- Perturbative improvements of lifetime predictions

Theoretical predictions for mixing observables

- Precise decay constants and Bag parameter for ΔM
- Additional Bag parameters at dimension 6 and 7 for Γ_{12}
- α_s/m_b corrections for Γ_{12}
- α_s^2 corrections for Γ_{12}

Theoretical predictions for charm mixing observables

- Push HQE to its limits
- Try to improve the exclusive approach

Update of theoretical predictions for inclusive rates

Moriond 2012: Conclusion from *B*-Mixing

It is actually not bad, what the Grinch left for us



Expansion in $1/m_b$ works so well,
What does this tell about charm? $1/m_c \approx 3 \cdot 1/m_b$



CKM⁻: How large are Penguins? II

Many observables in the B_s mixing system:

Elimination of $\Gamma_{12}^{\text{Theo}}$ via (No hint for incorrectness of $\Gamma_{12}^{\text{Theo}}$ except: A_{sl}^b is 1.5σ above bound)


$$a_{sl}^s = -\frac{\Delta\Gamma}{\Delta M} \frac{S_{\psi\phi}}{\sqrt{1 - S_{\psi\phi}^2}} \cdot \delta$$

not possible at that simple level, because $\delta \neq 1$

$$\delta = \frac{\tan(\phi_s^{\text{SM}} + \phi_s^\Delta)}{\tan(-2\beta_s^{\text{SM}} + \phi_s^\Delta + \delta_s^{\text{peng,SM}} + \delta_s^{\text{peng,NP}})}$$

A.L. 1106.3200

CKM⁻: How large are Penguins? III



- Above relation can be used to determine $\delta_s^{\text{peng,SM}} + \delta_s^{\text{peng,NP}}$

- To extract ϕ_s^Δ one needs $\Gamma_{12}^{s,\text{SM}}$


A.L. 1106.3200

Lifetimes: τ_{B^+}/τ_{B_d} in NLO-QCD I

$$\frac{\tau_1}{\tau_2} = 1 + \left(\frac{\Lambda}{m_b} \right)^3 \left(\Gamma_3^{(0)} + \frac{\alpha_s}{4\pi} \Gamma_3^{(1)} + \dots \right) + \left(\frac{\Lambda}{m_b} \right)^4 \left(\Gamma_4^{(0)} + \dots \right) + \dots$$

2002: Beneke, Buchalla, Greub, A.L., Nierste; Franco, Lubicz, Mescia, Tarantino

2004: Greub, A.L., Nierste; 2008 A.L.




$$\left[\frac{\tau(B^+)}{\tau(B_d^0)} \right]_{\text{LO,NLO,HFAG10}} = 1.047 \pm 0.049 \leftrightarrow 1.063 \pm 0.027 \leftrightarrow 1.071 \pm 0.009$$

Lifetimes: τ_{B^+}/τ_{B_d} in NLO-QCD II

$$\frac{\tau_1}{\tau_2} = 1 + \left(\frac{\Lambda}{m_b} \right)^3 \left(\Gamma_3^{(0)} + \frac{\alpha_s}{4\pi} \Gamma_3^{(1)} + \dots \right) + \left(\frac{\Lambda}{m_b} \right)^4 \left(\Gamma_4^{(0)} + \dots \right) + \dots$$

2002: Beneke, Buchalla, Greub, A.L., Nierste; Franco, Lubicz, Mescia, Tarantino

2004: Greub, A.L., Nierste; 2008 A.L.



$$\left[\frac{\tau(B^+)}{\tau(B_d^0)} \right]_{\text{LO,NLO,HFAG11}} = 1.047 \pm 0.049 \leftrightarrow 1.044 \pm 0.024 \leftrightarrow 1.079 \pm 0.007$$



Lifetimes: τ_{B^+}/τ_{B_d} in NLO-QCD III

$$\frac{\tau_{B^+}}{\tau_{B_d}} - 1 = 0.0324 \left(\frac{f_B}{200 \text{MeV}} \right)^2 [(1.0 \pm 0.2)B_1 + (0.1 \pm 0.1)B_2 - (17.8 \pm 0.9)\epsilon_1 + (3.9 \pm 0.2)\epsilon_2 - 0.26]$$

with non-perturbative input from [Becirevic hep-ph/0110124](#)

$$\begin{aligned} B_1 &= 1.10 \pm 0.20 \\ B_2 &= 0.79 \pm 0.10 \\ \epsilon_1 &= -0.02 \pm 0.02 \\ \epsilon_2 &= 0.03 \pm 0.01 \end{aligned}$$

Update urgently needed!



Lifetimes: Lifetimes of heavy hadrons

- $\tau(B^+)/\tau(B_d)$: HQE seems to fit, but we need urgently more precise hadronic matrix elements

$$\frac{\tau(B_s)}{\tau(B_d)} = 0.996 \dots 1.000 \leftrightarrow 0.969 \pm 0.017 \text{ HFAG 2011}$$

$$\text{A.L. 1102.4274} \leftrightarrow 1.004 \pm 0.018 \text{ LHCb-Conf2011-049}$$

More data as well as non-perturbative matrix elements needed

- $\tau(\Lambda_b)$, $\tau(\Xi_b)$ and $\tau(B_c)$: more data and further theory work (perturbative and non-perturbative) necessary

- $\tau(D)$, D-mixing: work in progress

Bigi, Uraltsev 2001; Bobrowski, A.L., Riedl, Rohrwild 1002.4794; 1011.5608;
Bobrowski, A.L. Nierste, Prill, to appear

It is not unplausible that HQE might give reasonable estimates

Theory statements about CPV in D before LHCb

[1002.4794] How large can the SM contribution to CP violation in $D^0-\bar{D}^0$ mixing be? - Mozilla Firefox

Alexander Lenz - Outlook W... [1002.4794] How large can t... arxiv.org/abs/arXiv:1002.4794

Most Visited Interactions.org - Par... SPIEGEL ONLINE - Na... Latest Headlines CERN Hot News

Cornell University Library We gratefully support

arXiv.org > hep-ph > arXiv:1002.4794 Search or Article-id (Help | All papers)

High Energy Physics - Phenomenology

How large can the SM contribution to CP violation in $D^0-\bar{D}^0$ mixing be?

M. Bobrowski, A. Lenz, J. Riedl, J. Rohrwild

(Submitted on 25 Feb 2010)

We investigate the maximum size of CP violating effects in $D^0-\bar{D}^0$ -mixing within the Standard Model (SM), using Heavy Quark Expansion (HQE) as theoretical working tool. For this purpose we determine the leading HQE contributions and also α_s corrections as well as subleading $1/m_c$ corrections to the absorptive part of the mixing amplitude of neutral D^0 mesons. It turns out that these contributions to Γ_{12} do not vanish in the exact $SU(3)\text{F}$ limit. Moreover, while the leading HQE terms give a result for Γ_{12} orders of magnitude lower than the current experimental value, we do find a sizeable phase. In the literature it was suggested that higher order terms in the HQE might be much less affected by the severe GIM cancellations of the leading terms; it is even not excluded that these higher order terms can reproduce the experimental value of Γ_{12} . If such an enhancement is realized in nature, the phase discovered in the leading HQE terms can have a sizeable effect. Therefore, we think that statements like: "CP violating effects in $D^0-\bar{D}^0$ -mixing of the order of 10^{-3} to 10^{-2} are an unambiguous sign of new physics"--given our limited knowledge of the SM prediction--are premature. Finally, we give an example of a new physics model that can enhance the leading HQE terms to Γ_{12} by one to two orders of magnitude.

Comments: 14 pages, considerably extended version of 0904.3971 with completely new main aspect; text (except title and abstract) identical to the version accepted by JHEP
Subjects: High Energy Physics - Phenomenology (hep-ph); High Energy Physics - Experiment (hep-ex)
Journal reference: JHEP 03(2010)009
Report number: DO-TH 10/04, TTK-10-2
Cite as: arXiv:1002.4794v1 [hep-ph]

Download:

- PDF
- PostScript
- Other formats

Current browse coauthors hep-ph < prev | next > new | recent | 1002 Change to browse mode hep-ex References & Citations

- INSPIRE HEP (refers to | cited by)
- NASA ADS

Bookmark (what is this)

ScienceWISE

Submission history

From: Alexander Lenz [view email]

[v1] Thu, 25 Feb 2010 14:27:00 GMT (97kb)

Which authors of this paper are endorsers?