



Summer School in Particle and
Astroparticle physics
of Annecy-le-Vieux

16-22 July 2015






IN2P3
Les deux infinis





Centre de
Physique Théorique
Grenoble-Alpes



A computing exercise using ROOT

Aim: give a taste of data analysis ..

- What is ROOT ?
 - ROOT is an object-oriented C++ analysis package
 - User-compiled code can be called to produce 1-d, 2-d, and 3-d graphics and histograms...



Outline

- Variables used in the analysis of $p - p$ collisions
- Useful relations
- Concept of **invariant mass**: example: 'inclusive' Z boson production
- Kinematics of $p - p$ collisions
- Analysis in $p - p$ collisions :
 - * **Signal**: Production of a W and a Z
 - * **Background**: Production of a pair of top-antitop
- Example: Macro.C

In all the following slides we assume the speed of the light

$$c=1$$


Variables used in the analysis of $p - p$ collisions

A particle ($Z, W, e^+, e^-, \text{etc ...}$) is described by its **four-momentum**:

$$\tilde{p} = (E, p_x, p_y, p_z)$$

The particle mass is $m = \sqrt{E^2 - p_x^2 - p_y^2 - p_z^2}$

When dealing with pp collision the following variables are used:




For **each** particle ($Z, W, e^+, e^-, \text{etc ...}$):

- 1. Transverse momentum/energy : $p_T = p \sin \theta$ $E_T = E \sin \theta$
- 2. Rapidity $Y = \frac{1}{2} \ln \frac{E+p_z}{E-p_z}$
or Pseudorapidity $\eta = - \ln \left(\tan \frac{\theta}{2} \right)$
- 3. Azimuthal angle Φ

Why?

Variables used in the analysis of $p - p$ collisions

Why? Many reasons.




1. The **longitudinal momentum of initial partons is unknown** while we know that $\vec{p}_T^{\text{initial partons}} \sim 0$
 → To exploit momentum conservation use transverse quantities (**in the plane \perp to the beam**) → p_T
2. p_T and ΔY are invariants for Lorentz transformations along the z axis 😊
3. $\sum^{\text{initial partons}} \vec{p}_T = \sum_{\text{vis}} \vec{p}_T + \sum_{\text{invis}} \vec{p}_T \approx 0$ → Allows to evaluate the p_T of particles not detected (ν)

$$\sum_{\text{invis}} \vec{p}_T = - \sum_{\text{vis}} \vec{p}_T$$

$$|\sum_{\text{invis}} \vec{p}_T| \text{ is the "missing } E_T \text{"}$$
4. The “interesting” physics is due to hard scattering processes → high p_T particles (selection of high p_T particles assures “interesting” physics)

Useful relations



$$\begin{aligned} p_x &= p_T \cos(\Phi); \\ p_y &= p_T \sin(\Phi); \\ p_z &= E \tanh(\eta); \end{aligned}$$

$$\begin{aligned} p_T &= p \sin \theta \\ \eta &= - \ln \left(\tan \frac{\theta}{2} \right) \end{aligned}$$

- $m \ll E \rightarrow Y \approx \eta$ (η doesn't require particle identification)
- $m \ll E \rightarrow p_T \approx E_T \quad E_T = E \sin \theta$

Concept of invariant mass: inclusive Z boson production

$$p - p \rightarrow Z X$$

With $Z \rightarrow e+e-$

$X = p_1, p_2, p_3, \dots$

Very 'clean' processes (low bkg)!!

Invariant mass M_{ee} of ee system from the 4-momentum conservation
(it allows to measure the Z mass, M_Z):


$$\tilde{p}_Z^2 = (\tilde{p}_{e1} + \tilde{p}_{e2})^2$$

$$M_{ee}^2 = (\tilde{p}_{e1} + \tilde{p}_{e2})^2 \approx 2 (E_{e1} E_{e2} - |\vec{p}_{e1}| |\vec{p}_{e2}| \cos \vartheta)$$

$$M_{ee} \approx \sqrt{2} E_{e1} E_{e2} (1 - \cos \vartheta_{e1 e2})$$

(the electron mass is neglected)

Why M_{ee} gives a distribution and not a single value?



$$1. \Delta E * \Delta t > \hbar/2 \quad \Delta m * \Delta \tau > \hbar/2$$


$$\Gamma * \tau > \hbar/2$$

width lifetime

2. Experimental resolution

Our signal : Production of a W and a Z

p - p 'hard' collisions in the $q_1 \bar{q}_2$ center of mass:




Kinematics of p - p collisions

★ 4-mom of the initial partons : $[(x_1+x_2)E_{\text{beam}}, 0, 0, (x_1-x_2) p_{\text{beam}}]$

Our background: Production of a pair of top-antitop

p - p 'hard' collisions. In the $q_1 \bar{q}_2$ center of mass:



Aim of the exercise:

- 1) look at some important variables,
- 2) build the Z invariant mass,
- 3) how one can discriminate between the 'signal' and the 'background'

On the GRASPA 2015 Web page you will be given :

<https://indico.in2p3.fr/event/11292/other-view?view=standard>

[GRASPA2015explanation.pptx.pdf](#)

[Selected_All_EEM.root](#)

[Exercise2015.pdf](#)

[macro.C](#)

[Tutorial_ROOT_Bose.pdf](#)

[macro_final.C](#)

GRASPA 2015 - L. Di Ciaccio - LAPP, 16-22 July 2015

9

1) an input file containing the physics: [Selected_All_EEM.root](#)

```
==== MOST ENERGETIC LEPTON FROM THE Z
Br 4 :pt1 : pt1
Br 5 :eta1 : eta1
Br 6 :phi1 : phi1
Br 7 :E1 : E1

==== SECOND ENERGETIC LEPTON FROM THE Z
Br 8 :pt2 : pt2
Br 9 :eta2 : eta2
Br 10 :phi2 : phi2
Br 11 :E2 : E2

==== LEPTON FROM W
Br 12 :pt3 : pt3
Br 13 :eta3 : eta3
Br 14 :phi3 : phi3
Br 15 :E3 : E3
```

List of variables given
per each collision event
(kinematics of the final
state leptons)

2) Instructions to make the computing exercise : [Exercise2015.pdf](#)

GRASPA 2015

16-22 July 2015



COMPUTING EXERCISE

Study of the production of a pair of gauge bosons (W and Z) at the LHC

The data to analyse are organised into a 'Root n-tuple' which we will provide to you. The Root n-tuple is a file containing information about the kinematics of

GRASPA 2015 - L. Di Ciaccio - LAPP, 16-22 July 2015

10

3) A skeleton of an analysis program using ROOT: **macro.C**

```
#include "TCanvas.h"
#include "TR00T.h"
#include "TFile.h"
#include "TTree.h"
#include "TBrowser.h"
#include "TH2.h"
#include "TRandom.h"

void tree1r()
{
    // Read Selected_All_EEM.root file
    //Root file
    TFile *f = new TFile("Selected_All_EEM.root");

    // Signal events
    TTree *sig = (TTree*)f->Get("WZSignal");
    Double_t pt1, eta1, phi1, E1;
    Double_t pt2, eta2, phi2, E2;
    Double_t pt3, eta3, phi3, E3;
    Double_t MZ, MET, trackd0cutWMu, TrackIsoWmu;
    Double_t Weight;

    //get some variables for SIGNAL EVENTS
    sig->SetBranchAddresses("pt1",&pt1);
```

Look also to:

ROOT Tutorial

4) A 'manual' with ROOT instructions: **Tutorial_ROOT_Bose.pdf**

Tulika Bose
Brown University
NEPPSR 2007

Useful links about root:

<http://root.cern.ch/drupal/content/root-primer-534>

<http://root.cern.ch/drupal/content/root-users-guide-534>

Example of analysis program

macro.C

23/07/2013 00:21

```
#include "TCanvas.h"
#include "TROOT.h"
#include "TFile.h"
#include "TTree.h"
#include "TBrowser.h"
#include "TH2.h"
#include "TRandom.h"

void tree1r()
{
  // Read Selected_All_EEM.root file
  //Root file
  TFile *f = new TFile("Selected_All_EEM.root");

  // Signal events
  TTree *sig = (TTree*)f->Get("WZSignal");
  Double_t pt1, eta1, phi1, E1;
  Double_t pt2, eta2, phi2, E2;
  Double_t pt3, eta3, phi3, E3;
  Double_t MZ, MET, trackd0cutWMu, TrackIsoWmu;
  Double_t Weight;

  //get some variables for SIGNAL EVENTS
  sig->SetBranchAddresses("pt1",&pt1);
  sig->SetBranchAddresses("eta1",&eta1);
  sig->SetBranchAddresses("phi1",&phi1);
  sig->SetBranchAddresses("E1",&E1);
  sig->SetBranchAddresses("MZ",&MZ);
  sig->SetBranchAddresses("Weight",&Weight);
  // add other variables ...
}
```

Open the input file

Access the Signal info

Variables per each lepton

GRASPA 2015 - L. Di Ciaccio - LAPP, 16-22 July 2015

13

```
////get some variables for BACKGROUND EVENTS
TTree *ttbar = (TTree*)f->Get("ttbar");
Double_t pt1_bkg, eta1_bkg, phi1_bkg, E1_bkg;
Double_t MZ_bkg;
Double_t Weight_bkg;

//get some variables for ttbar
ttbar->SetBranchAddresses("pt1",&pt1_bkg);
ttbar->SetBranchAddresses("eta1",&eta1_bkg);
ttbar->SetBranchAddresses("phi1",&phi1_bkg);
ttbar->SetBranchAddresses("E1",&E1_bkg);
ttbar->SetBranchAddresses("MZ",&MZ_bkg);
ttbar->SetBranchAddresses("Weight",&Weight_bkg);
// add other variables ...

//create two histograms (for sig and ttbar)
TH1F *h_MZ = new TH1F("h_MZ","MZ distribution All events",40,65,115);
TH1F *h_MZ_bkg = new TH1F("h_MZ_bkg","MZ distribution BKG",40,65,115);
TH1F *h_MZ_sig = new TH1F("h_MZ_sig","MZ distribution SIG",40,65,115);

//read all SIGNAL entries and fill the histograms
Int_t nentries = (Int_t)sig->GetEntries();

for (Int_t i=0;i<nentries;i++) {
  sig->GetEntry(i);
  h_MZ->Fill(MZ,Weight);
  h_MZ_sig->Fill(MZ,Weight);
}
```

Access the background info

Page 1 of 2

GRASPA 2015 - L. Di Ciaccio - LAPP, 16-22 July 2015

14

Have fun !!



$W(jj) Z(jj)$

$jj = J$ (1 fat jet)

Another example: search for di-boson resonances

- Is there something hiding in the data, waiting to be discovered?

