

Dark Matter from μ Anomalies

Cédric Delaunay
CNRS/LAPTh
Annecy-le-Vieux
France

content based on:

G. Bélanger, CD, S. Westhoff
Phys.Rev. D92 (2015) 5, 055021
arXiv:1507.06660 [hep-ph]

CoDyCE workshop
25-11-2015 | Lyon

Outline

1. $g_\mu - 2$ and $b \rightarrow s\mu\mu$ anomalies \Rightarrow DM
2. a toy model for muon-philic DM
3. DM relic density
4. collider signatures

First signs of DM
in B physics?*

**non-gravitational*

$B \rightarrow (K, K^*, \phi) + \mu^+ \mu^-$ anomalies

Decay	obs.	q^2 bin	SM pred.	measurement	pull
$\bar{B}^0 \rightarrow \bar{K}^{*0} \mu^+ \mu^-$	F_L	[2, 4.3]	0.81 ± 0.02	0.26 ± 0.19	ATLAS +2.9
$\bar{B}^0 \rightarrow \bar{K}^{*0} \mu^+ \mu^-$	F_L	[4, 6]	0.74 ± 0.04	0.61 ± 0.06	LHCb +1.9
$\bar{B}^0 \rightarrow \bar{K}^{*0} \mu^+ \mu^-$	S_5	[4, 6]	-0.33 ± 0.03	-0.15 ± 0.08	LHCb -2.2
$\bar{B}^0 \rightarrow \bar{K}^{*0} \mu^+ \mu^-$	P'_5	[1.1, 6]	-0.44 ± 0.08	-0.05 ± 0.11	LHCb -2.9
$\bar{B}^0 \rightarrow \bar{K}^{*0} \mu^+ \mu^-$	P'_5	[4, 6]	-0.77 ± 0.06	-0.30 ± 0.16	LHCb -2.8
$B^- \rightarrow K^{*-} \mu^+ \mu^-$	$10^7 \frac{d\text{BR}}{dq^2}$	[4, 6]	0.54 ± 0.08	0.26 ± 0.10	LHCb +2.1
$\bar{B}^0 \rightarrow \bar{K}^0 \mu^+ \mu^-$	$10^8 \frac{d\text{BR}}{dq^2}$	[0.1, 2]	2.71 ± 0.50	1.26 ± 0.56	LHCb +1.9
$\bar{B}^0 \rightarrow \bar{K}^0 \mu^+ \mu^-$	$10^8 \frac{d\text{BR}}{dq^2}$	[16, 23]	0.93 ± 0.12	0.37 ± 0.22	CDF +2.2
$B_s \rightarrow \phi \mu^+ \mu^-$	$10^7 \frac{d\text{BR}}{dq^2}$	[1, 6]	0.48 ± 0.06	0.23 ± 0.05	LHCb +3.1


$$\mathcal{H}_{\text{eff}}^{\text{NP}} = -\frac{\alpha G_F}{2\sqrt{2}\pi} V_{tb} V_{ts}^* \sum_i C_i \mathcal{O}_i + \text{h.c.}$$

$$\begin{aligned} \mathcal{O}_9^\ell &\equiv \bar{b} \gamma_\rho (1 - \gamma_5) s \bar{\ell} \gamma^\rho \ell \\ \mathcal{O}_{10}^\ell &\equiv \bar{b} \gamma_\rho (1 - \gamma_5) s \bar{\ell} \gamma^\rho \gamma_5 \ell \end{aligned}$$

2 favored simple
NP structure:

$C_9 \simeq -1, C_{10} = C'_{9,10} = 0$
or TH favored

$C_{10} = -C_9 \simeq 0.5, C'_{9,10} = 0$




R_K anomaly

Adding electron data, in particular

$$R_K \equiv \frac{\text{BR}(\text{B} \rightarrow \text{K}\mu^+\mu^-)_{[1,6]}}{\text{BR}(\text{B} \rightarrow \text{K}\text{e}^+\text{e}^-)_{[1,6]}} = 0.745^{+0.090}_{-0.074} \pm 0.036 \lesssim 1$$

global fits indicate violation of
Lepton Flavor Universality

$$C_{9,10}^e \ll C_{9,10}^\mu$$



$g_\mu - 2$ anomaly

Long story with many bounces between TH and EXP,
But current status still anomalous:

$$(a_\mu \equiv \frac{g_\mu - 2}{2})$$


$$\Delta a_\mu \equiv a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (287 \pm 80) \times 10^{-11}$$

If discrepancy originates from one-loop new physics,
weak scale/coupling are favored

$$a_\mu^{\text{BSM}} \sim \frac{g^2 m_\mu^2}{16\pi^2 M^2} \sim 3 \times 10^{-9} \left(\frac{g}{0.6}\right)^2 \left(\frac{100 \text{ GeV}}{M}\right)^2$$

A hint for DM

- Tree-level explanations of $b \rightarrow s$ typically yields a too small contribution to $g_\mu - 2$:



$$\rightarrow a_\mu^{Z'} \sim \mathcal{O}(10^{-11}) \left(\frac{1 \text{ TeV}}{m_{Z'}} \right)^2$$

- If $b \rightarrow s$ arises at one-loop, $g_\mu - 2$ in the right ballpark.
- A simple way with a Z' : seclude it from SM, break $U(1)_X \rightarrow Z_2$ \rightarrow DM candidate

*A toy model
for loop-induced Z'
couplings to muons*

The simplest toy model

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \boxed{\mathcal{L}_{\text{dark}}} + \mathcal{L}_{\text{portal}}$$

Dark $\text{U}(1)_X$ field, SM carry no X charge,
Dark sector fields:

mediators to
SM fermions
 $\text{U}(1)$ breaking
Dark Matter

	spin	$\text{SU}(3)_c$	$\text{SU}(2)_L$	$\text{U}(1)_Y$	$\text{U}(1)_X$
L, L^c	$1/2$	1	2	$-1/2$	1
Q, Q^c	$1/2$	3	2	$1/6$	-2
ϕ	0	1	1	0	2
χ	0	1	1	0	-1

The simplest toy model

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \mathcal{L}_{\text{dark}} + \boxed{\mathcal{L}_{\text{portal}}}$$

$$\mathcal{L}_{\text{portal}} = \boxed{\epsilon B \cancel{\chi_{\text{closed}}^{\mu\nu}}} - \lambda_{\chi H} \cancel{|\chi|^2} - \lambda_{\phi H} |\phi|^2 |H|^2$$

hypercharge portal *Higgs portal*


$$- y (\bar{l} L) \chi - w (\bar{q} Q) \phi + \text{h.c.}$$

‘Yukawa’ portal *mass mixing = “partial darkness”*

The X charge assignment forbids tree-level $Z' \bar{l} l$ coupling, makes DM leptophilic, explaining direct detection null results

Z' coupling to SM fermions

- Quarks: partial darkness $w(\bar{q}Q)\langle\phi\rangle + \text{h.c.}$



switch on only $\bar{b}_L s_L$ coupling
motivated by LHCb anomalies
(not much changes with also $\bar{s}_L s_L$)


$B - \bar{B}$ mixing constraint*:

$$g_{bs}^L \lesssim 2.4 \times 10^{-3} \left(\frac{m_{Z'}/g_X}{300 \text{ GeV}} \right)$$

*NP<0.1SM

Z' coupling to SM fermions

- Leptons: $(\bar{l}L)\phi$ not allowed \longrightarrow one-loop



$$g_{\mu\mu}^L = \frac{|y|^2}{32\pi^2} F(\tau, \delta)$$


$$\tau \equiv \frac{m_L^2}{m_\chi^2}, \delta \equiv \frac{m_{\chi'}^2}{m_\chi^2} - 1$$

For $\delta \rightarrow 0$, $g_{\mu\mu}^L(q^2 = 0) = 0$, $\langle \phi \rangle$ -insertion needed

$$(\phi^* D_\rho \phi \bar{\mu}_L \gamma^\rho \mu_L)$$

\longrightarrow significant DM-DM' splitting

Explaining $b \rightarrow s\mu\mu$ anomalies



$$C_{10,-9}^\mu = \frac{g_X^2 \Lambda_{\text{SM}}^2}{m_{Z'}^2} \frac{|V_{ts}^* V_{tb}|}{V_{ts}^* V_{tb}} g_{bs}^L g_{\mu\mu}^L$$

$$\Lambda_{\text{SM}} \equiv [2\sqrt{2}\pi/(\alpha G_F |V_{ts}^* V_{tb}|)]^{1/2} \simeq 50 \text{ TeV}$$


Under $B - \bar{B}$ constraint,
explaining LHCb requires:

$$g_{\mu\mu}^L \gtrsim 0.03 \left(\frac{m_{Z'}/g_X}{300 \text{ GeV}} \right)$$

In turns, one needs: $y \gg 1$, $\delta \gg 1$, a light Z'
and $\tau \simeq [1, \delta]$

Accomodating Δa_μ sets DM mass

$g_\mu - 2$ further constrains the parameter space:



$$\Delta a_\mu = \frac{|y|^2 m_\mu^2}{32\pi^2 m_\chi^2} G(\tau, \delta)$$


$$\tau \equiv \frac{m_L^2}{m_\chi^2}, \delta \equiv \frac{m_{\chi'}^2}{m_\chi^2} - 1$$

2 limiting scenarios:

$\tau \simeq 1 \longrightarrow m_\chi \simeq 30|y|(1 + 2/\delta) \text{ GeV}$

$\tau \simeq \delta \longrightarrow m_\chi \simeq 55 \frac{|y|}{\sqrt{\delta}} \text{ GeV}$

So, muon anomalies favors:



Large δ needs tuning in
the dark scalar potential:


$$\delta \sim \mathcal{O}(10)$$

which guarantees DM is
scalar (see relic later)

*DM relic prediction
from muon anomalies*

Scalar DM annihilation

- At freeze-out dominant process is d-wave:



$$\langle\sigma v\rangle = \frac{|y|^4 x^{-2}}{2\pi(1+\tau)^4 m_\chi^2}$$

$$\langle v^2 \rangle^{-1} \sim x \equiv \frac{m_\chi}{T} \simeq 25$$

- Thus DM relic also favors large y ,
for weak scale m_χ

Relic density prediction

- Note that: $\Delta a_\mu \propto \frac{|y|^2}{m_\chi^2}$, $C_{9,-10} \propto |y|^2$

Hence: $\Omega_\chi \sim \int_T \langle \sigma v \rangle^{-1} \propto \Delta a_\mu^{-1} \times C_{9,-10}^{-1}$
and

$$\Omega_\chi h^2 \simeq 0.01 R(\tau, \delta) \times \left(\frac{x}{25} \right)^3 \left(\frac{100 \text{ GeV}}{m_{Z'} / g_X} \right)$$

with


$$R(\tau, \delta) \equiv F(\tau, \delta) G(\tau, \delta) (1 + \tau)^4 \sim \mathcal{O}(10)$$

Muon anomalies yields DM relic
in the right ballpark!

$$\Omega_{\text{DM}}^{\text{CMB}} h^2 \simeq 0.11$$

Is Relic prediction robust?

Sniffing around the parameter space with micrOMEGAs,
under the anomalies constraints




Relatively robust, except in peculiar corners with:

- 1) strong coannihilation $m_L \gtrsim m_\chi$
and/or
- 2) open Z' channel $m_{Z'} < m_\chi$

*Tests at colliders
and DM detection experiments*

EW Precision test

- The $Z\bar{\mu}\mu$ coupling is shifted at one-loop:




corrects R_μ and A_{FB}^μ

$$\frac{\delta g_{\mu_L}^Z}{g_{\text{SM}}} \simeq (1 - 3) \times 10^{-3}$$

- Mild tension with LEP data: $\delta g_{\mu_L}^Z / g_{\text{SM}} \lesssim 10^{-3}$
- May-be solved in complete model?

Heavy lepton searches at LHC

- $pp \rightarrow Z^* \rightarrow LL \rightarrow \mu\mu + \cancel{E}_T$



- Recasting SUSY searches for smuons:

$m_L \gtrsim 450 \text{ GeV}$


unless $\tau \simeq 1$
where missing E_T is missing

Dimuons from resonant Z'

- LHC bound on Z' with Z -like fermions couplings are rather strong: $m_{Z'_{\text{SSM}}} \gtrsim 3 \text{ TeV}$
- Here Z' produced through $\bar{b}s + \bar{s}b$ with $\text{BR}(Z' \rightarrow \mu\mu) \sim \mathcal{O}(50\%)$, so one finds $\sigma(pp \rightarrow Z' \rightarrow \mu\mu) \sim \mathcal{O}(\text{fb})$ for $m_{Z'} = 300 \text{ GeV}$
- LHC8 sensitivity $\sim 10 \text{ fb}$, it's not clear the low mass region will improve at 13-14 TeV...

Indirect DM detection

- Today $v_{\text{halo}} \sim 10^{-3}$: 2 body ann. suppressed
- Dominant process = $\chi\chi \rightarrow \mu\mu\gamma$ (s-wave)
- Could give gamma-ray signals at FERMI:



Wrapping up

Conclusions

- Loop-induced Z' couplings to SM allow to simultaneously address $g_\mu - 2$ and $b \rightarrow s\mu\mu$
- The symmetry forbidding tree-level couplings typically breaks to a discrete subgroup yielding (leptophilic) DM!
- The same interaction controlling the muon anomalies drives DM annihilation in the early Universe, giving $\Omega_\chi \simeq 0.1$!



Beauty in the sky with muons

Direct detection below ν -bkg

