

# Interpreting a 2 TeV resonance in WW scattering

**Domènec Espriu**

Institut de Ciències del Cosmos (ICCUB),  
Universitat de Barcelona

Lyon, November 2015



Institut de Ciències del Cosmos






EXCELENCIA  
MARÍA  
DE MAEZTU

- D.E. and B. Yencho, PRD 87, 055017 (2013) [1212.4158]
- D.E., F. Mescia and B. Yencho, PRD 88, 055002 (2013) [1307.2400]
- D.E. and F. Mescia, PRD 90, 015035 (2014) [1403.7386]
- P. Arnan, D.E. and F. Mescia, 1508.00174
- R. Delgado, A. Dobado, D.E., F. Llanes, 1510.03761


Closely related work:

- R. Delgado, A. Dobado and F. Llanes, JHEP 1402 (2014) 121 [1311.5993] & 1502.04841, 1509.04725
- R. Delgado, A. Dobado, M.J. Hererro and J.J. Sanz-Cillero, JHEP 1407 (2014) 149 [1404.2866]
- A. Dobado, F-K. Guo, F. Llanes, 1508.03544


# A resonance at 2 TeV?



# Excess events at 2 TeV




ATLAS WZ events



CMS WW events

# $W_L W_L$ resonances?

We know that in the SM the Higgs boson **unitarizes**  $W_L W_L$  scattering. Consider e.g.  $W_L^+ W_L^- \rightarrow Z_L Z_L$



If any of these couplings are different from SM values, the careful balance necessary for perturbative unitarity is lost.

The first 3 diagrams are fixed by gauge invariance, but we can contemplate other Higgs-gauge boson couplings. For  $s \gg M_W^2$  the amplitude in the SM goes for  $s \rightarrow \infty$  as

$$\frac{s}{v^2} \frac{M_H^2}{s - M_H^2} \sim \frac{M_H^2}{v^2}$$

# $W_L W_L$ resonances?

... but on dimensional grounds it *should* go as (cf. pion physics)

$$\frac{s}{v^2} \frac{s}{s - M_H^2} \sim \frac{s}{v^2}$$

This is what happens after *any modification* of the Higgs couplings and produces *non-unitary* amplitudes.

Adding *new effective operators* typically spoils unitarity too.

$$\mathcal{L}_{SM} \rightarrow \mathcal{L}_{SM} + \sum_i a_i \mathcal{O}_i \quad \mathcal{O}_i \sim s^2$$

*New physics may produce either type of modifications*

*What can the unitarity of longitudinal WW scattering tell us about anomalous couplings in EW sector?*

... but on dimensional grounds it *should* go as (cf. pion physics)

$$\frac{s}{v^2} \frac{s}{s - M_H^2} \sim \frac{s}{v^2}$$

This is what happens after *any modification* of the Higgs couplings and produces *non-unitary* amplitudes.

Adding *new effective operators* typically spoils unitarity too.

$$\mathcal{L}_{SM} \rightarrow \mathcal{L}_{SM} + \sum_i a_i \mathcal{O}_i \quad \mathcal{O}_i \sim s^2$$

*New physics may produce either type of modifications*

*What can the unitarity of longitudinal WW scattering tell us about anomalous couplings in EW sector?*

... but on dimensional grounds it *should* go as (cf. pion physics)

$$\frac{s}{v^2} \frac{s}{s - M_H^2} \sim \frac{s}{v^2}$$

This is what happens after *any modification* of the Higgs couplings and produces *non-unitary* amplitudes.

Adding *new effective operators* typically spoils unitarity too.

$$\mathcal{L}_{SM} \rightarrow \mathcal{L}_{SM} + \sum_i a_i \mathcal{O}_i \quad \mathcal{O}_i \sim s^2$$

*New physics may produce either type of modifications*

*What can the unitarity of longitudinal WW scattering tell us about anomalous couplings in EW sector?*



# Parametrizing composite Higgs physics

A light “Higgs boson” with mass  $M_H \sim 125$  GeV is coupled to the EW bosons according to (non-linear realization)

$$\mathcal{L}_{\text{eff}} \supset -\frac{1}{2}\text{Tr}W_{\mu\nu}W^{\mu\nu} - \frac{1}{4}\text{Tr}B_{\mu\nu}B^{\mu\nu} + \mathcal{L}_{\text{GF}} + \mathcal{L}_{\text{FP}} + \sum_i \mathcal{L}_i$$
$$+ \left[ 1 + 2a \left( \frac{h}{v} \right) + b \left( \frac{h}{v} \right)^2 \right] \frac{v^2}{4} \text{Tr} D_\mu U^\dagger D^\mu U - V(h)$$

$$U = \exp(i \omega \cdot \tau / v)$$

$$D_\mu U = \partial_\mu U + \frac{1}{2} ig W_\mu^i \tau^i U - \frac{1}{2} ig' B_\mu U \tau^3$$

and additional gauge-invariant operators are encoded in  $\mathcal{L}_i$ .

Setting  $a = b = 1$  (and  $\mathcal{L}_i = 0$ ) reproduces the **SM interactions**

# $\mathcal{O}(p^4)$ operators

The  $\mathcal{L}_i$  are a full set of  $C$ ,  $P$ , and  $SU(2)_L \times U(1)_Y$  gauge invariant,  $d = 4$  operators that parameterize the *low-energy effects* of the *model-dependent high-energy EWSB sector* along with  $a, b$ . The two relevant *custodial-symmetry preserving* operators are


$$\mathcal{L}_4 = a_4 (\text{Tr} [V_\mu V_\nu])^2 \quad \mathcal{L}_5 = a_5 (\text{Tr} [V_\mu V^\mu])^2 \quad V_\mu = (D_\mu U) U^\dagger$$

The  $a_i$  could be *functions of  $\frac{\hbar}{v}$*

- For example: *Heavy Higgs* *QCD-like technicolor*

$$\begin{array}{lcl} a_4 = & 0 & -2a_5 \\ a_5 = & \frac{v^2}{8M_H^2} & \frac{N_{TC}}{96\pi^2} \end{array}$$

(up to logarithmic corrections)



$$ig^4 [a_4 (g^{\mu\sigma} g^{\nu\rho} + g^{\mu\rho} g^{\nu\sigma}) + 2a_5 g^{\mu\nu} g^{\rho\sigma}]$$

# What if the $hWW$ couplings are not exactly the SM ones?

$$\begin{aligned}\mathcal{L}_{\text{eff}} = & -\frac{1}{2}\text{Tr}W_{\mu\nu}W^{\mu\nu} - \frac{1}{4}\text{Tr}B_{\mu\nu}B^{\mu\nu} + \sum_i \mathcal{L}_i + \mathcal{L}_{\text{GF}} + \mathcal{L}_{\text{FP}} \\ & + \left[1 + 2a\left(\frac{h}{v}\right) + b\left(\frac{h}{v}\right)^2\right] \frac{v^2}{4}\text{Tr}D_\mu U^\dagger D^\mu U + \frac{1}{2}(\partial_\mu h)^2 - \frac{1}{2}M_H^2 h^2 \\ & - d_3(\lambda v)h^3 - d_4\frac{1}{4}h^4\end{aligned}$$

This *effective theory* is *non-renormalizable* and the  $a_i$  will be required to absorb the divergences. They will be running parameters (unlike for  $a = b = 1$ )

$$\delta a_4 = \Delta_\epsilon \frac{1}{(4\pi)^2} \frac{-1}{12} (1 - a^2)^2$$

$$\delta a_5 = \Delta_\epsilon \frac{1}{(4\pi)^2} \frac{-1}{24} \left[ (1 - a^2)^2 + \frac{3}{2} ((1 - a^2) - (1 - b))^2 \right]$$

We have set  $d_3 = d_4 = 1$  for simplicity.


# Early Higgs measurements

There are solid indications that the “Higgs” couples to the  $W, Z$  *very similarly to the SM rules*

$$\mathcal{L}_{\text{eff}} \simeq \mathcal{L}_{\text{SM}} + a_4 (\text{Tr} [V_\mu V_\nu])^2 + a_5 (\text{Tr} [V_\mu V^\mu])^2$$

Then  $a_4$  and  $a_5$  represent anomalous 4-point couplings of the  $W$  bosons due to an extended EWSBS that however does not manifest with  $O(p^2)$  couplings *noticeably different to the ones in the SM*. **Assume now that  $a = b = 1$  exactly.**

These operators will lead to **violations** of perturbative unitarity at loop level ( $\sim g^4$ )



$$\sim \left( \frac{s}{v^2} \right)^2$$

*Violations of unitarity are cured by the appearance of new particles or resonances*

*We can now use well-understood unitarization techniques to constrain these resonances and the effective couplings  $\{a_i\}$*


# Early Higgs measurements

There are solid indications that the “Higgs” couples to the  $W, Z$   
*very similarly to the SM rules*

$$\mathcal{L}_{\text{eff}} \simeq \mathcal{L}_{\text{SM}} + a_4 (\text{Tr} [V_\mu V_\nu])^2 + a_5 (\text{Tr} [V_\mu V^\mu])^2$$

Then  $a_4$  and  $a_5$  represent anomalous 4-point couplings of the  $W$  bosons due to an extended EWSBS that however does not manifest with  $O(p^2)$  couplings *noticeably different to the ones in the SM*. **Assume now that  $a = b = 1$  exactly.**

These operators will lead to **violations** of perturbative unitarity at loop level ( $\sim g^4$ )



$$\sim \left( \frac{s}{v^2} \right)^2$$

*Violations of unitarity are cured by the appearance of new particles or resonances*

*We can now use well-understood unitarization techniques to constrain these resonances and the effective couplings  $\{a_i\}$*


# Early Higgs measurements

There are solid indications that the “Higgs” couples to the  $W, Z$  *very similarly to the SM rules*

$$\mathcal{L}_{\text{eff}} \simeq \mathcal{L}_{\text{SM}} + a_4 (\text{Tr} [V_\mu V_\nu])^2 + a_5 (\text{Tr} [V_\mu V^\mu])^2$$

Then  $a_4$  and  $a_5$  represent anomalous 4-point couplings of the  $W$  bosons due to an extended EWSBS that however does not manifest with  $O(p^2)$  couplings *noticeably different to the ones in the SM*. **Assume now that  $a = b = 1$  exactly.**

These operators will lead to **violations** of perturbative unitarity at loop level ( $\sim g^4$ )



$$\sim \left( \frac{s}{v^2} \right)^2$$

*Violations of unitarity are cured by the appearance of new particles or resonances*

*We can now use well-understood unitarization techniques to constrain these resonances and the effective couplings  $\{a_i\}$*

We would like to

- Determine how much room is left for the  $a_i$
- Find possible additional resonances imposed by unitarity
- Should we have already seen any?
- To what extent an extended EWSBS is excluded?
- What about this putative 2 TeV resonance?

*Yes, there are new resonances even with relatively light masses  
Their signal is very weak. Can a 2 TeV EWSBS resonance be visible?*

*Searching for resonances is an efficient (albeit indirect) way of setting constraints on  $a_{TGC}$  and  $a_{QGC} \leftarrow \{a_i\}$*

IMPORTANT: even very small values of  $a_4$  and  $a_5$  can lead to resonances. Their effect is more important than having  $a \neq 1$  (within reason).

We would like to

- Determine how much room is left for the  $a_i$
- Find possible additional resonances imposed by unitarity
- Should we have already seen any?
- To what extent an extended EWSBS is excluded?
- What about this putative 2 TeV resonance?

*Yes, there are new resonances even with relatively light masses  
Their signal is very weak. Can a 2 TeV EWSBS resonance be visible?*

*Searching for resonances is an efficient (albeit indirect) way of setting constraints on  $a_{TGC}$  and  $a_{QGC} \leftarrow \{a_i\}$*

IMPORTANT: even very small values of  $a_4$  and  $a_5$  can lead to resonances. Their effect is more important than having  $a \neq 1$  (within reason).



We would like to

- Determine how much room is left for the  $a_i$
- Find possible additional resonances imposed by unitarity
- Should we have already seen any?
- To what extent an extended EWSBS is excluded?
- What about this putative 2 TeV resonance?

*Yes, there are new resonances even with relatively light masses  
Their signal is very weak. Can a 2 TeV EWSBS resonance be visible?*

*Searching for resonances is an efficient (albeit indirect) way of setting constraints on  $a_{TGC}$  and  $a_{QGC} \leftarrow \{a_i\}$*

IMPORTANT: even very small values of  $a_4$  and  $a_5$  can lead to resonances. Their effect is more important than having  $a \neq 1$  (within reason).

We would like to

- Determine how much room is left for the  $a_i$
- Find possible additional resonances imposed by unitarity
- Should we have already seen any?
- To what extent an extended EWSBS is excluded?
- What about this putative 2 TeV resonance?

*Yes, there are new resonances even with relatively light masses  
Their signal is very weak. Can a 2 TeV EWSBS resonance be visible?*

*Searching for resonances is an efficient (albeit indirect) way of setting constraints on  $a_{TGC}$  and  $a_{QGC} \leftarrow \{a_i\}$*


IMPORTANT: even very small values of  $a_4$  and  $a_5$  can lead to resonances. Their effect is more important than having  $a \neq 1$  (within reason).

# Equivalence Theorem

Most studies concerning unitarity at high energies are (understandably) carried out using the ET

$$A(W_L^+ W_L^- \rightarrow Z_L Z_L) \rightarrow A(\omega^+ \omega^- \rightarrow \omega^0 \omega^0) + O(M_W/\sqrt{s})$$

For a light Higgs the region one needs to include tree-level Higgs exchange as well



Then one could make use of the well known chiral lagrangian techniques to derive the amplitudes and compare with experiment, including the Higgs as an explicit resonance.

*However for  $s$  not too large (which obviously is now an interesting region) the ET may be too crude an approximation.*


*We shall use as much as possible exact amplitudes*

# Equivalence Theorem

Most studies concerning unitarity at high energies are (understandably) carried out using the ET

$$A(W_L^+ W_L^- \rightarrow Z_L Z_L) \rightarrow A(\omega^+ \omega^- \rightarrow \omega^0 \omega^0) + O(M_W/\sqrt{s})$$

For a light Higgs the region one needs to include tree-level Higgs exchange as well



Then one could make use of the well known chiral lagrangian techniques to derive the amplitudes and compare with experiment, including the Higgs as an explicit resonance.

*However for  $s$  not too large (which obviously is now an interesting region) the ET may be too crude an approximation.*


*We shall use as much as possible exact amplitudes*

# Equivalence Theorem

Most studies concerning unitarity at high energies are (understandably) carried out using the ET

$$A(W_L^+ W_L^- \rightarrow Z_L Z_L) \rightarrow A(\omega^+ \omega^- \rightarrow \omega^0 \omega^0) + O(M_W/\sqrt{s})$$

For a light Higgs the region one needs to include tree-level Higgs exchange as well



Then one could make use of the well known chiral lagrangian techniques to derive the amplitudes and compare with experiment, including the Higgs as an explicit resonance.

*However for  $s$  not too large (which obviously is now an interesting region) the ET may be too crude an approximation.*

*We shall use as much as possible exact amplitudes.*

# Partial wave amplitudes

We will assume  $e = 0$  (*no e.m.*) i.e. the custodial limit  $c_W = 1$ .  
The  $WW$  scattering amplitudes can then be deconstructed into amplitudes of *fixed isospin*  $T_I$

$$\begin{aligned}T_0 &= 3A^{+-00} + A^{++++} \\T_1 &= 2A^{+--+} - 2A^{+-00} - A^{++++} \\T_2 &= A^{++++}\end{aligned}$$

where  $A^{+-00} = A(W_L^+ W_L^- \rightarrow Z_L Z_L)$  and all others may be expressed in terms of this amplitude through isospin and crossing symmetries

These can then be written in terms of *partial waves*

$$t_{IJ}(s) = \frac{1}{64\pi} \int_{-1}^1 d(\cos \theta) P_J(\cos \theta) T_I$$

which are constrained by unitarity at high energies to be  $|t_{IJ}| < 1$ .  
Most discussions based on unitarity are based on this simple constraint (*tree-level unitarity*)

# Inverse Amplitude method

Partial wave unitarity **requires**

$$\text{Im } t_{IJ}(s) = \underbrace{\sigma(s)|t_{IJ}(s)|^2}_{\substack{\text{Elastic} \\ WW \rightarrow WW}} + \underbrace{\sigma_H(s)|t_{H,IJ}(s)|^2}_{\substack{\text{Inelastic} \\ WW \rightarrow hh}}$$

where  $\sigma$  and  $\sigma_H$  are phase space factors.

Given a perturbative expansion

$$t_{IJ} \approx \underbrace{t_{IJ}^{(2)}}_{\text{tree}} + \underbrace{t_{IJ}^{(4)}}_{\substack{\text{one-loop} \\ + a_i \text{ terms}}} + \dots$$

we can require unitarity to hold *exactly* by defining (*Note: non-coupled channels*)

$$t_{IJ} \approx \frac{t_{IJ}^{(2)}}{1 - t_{IJ}^{(4)}/t_{IJ}^{(2)}}$$

Several mild analyticity assumptions are implied.

The **unitarization** of the amplitudes may result in the appearance of **new heavy resonances** associated with the high-energy theory

$t_{00}$  → Scalar isoscalar

$t_{11}$  → Vector isovector

$t_{20}$  → Scalar isotensor

Will search for poles in  $t_{IJ}(s)$  up to  $(4\pi v) \sim 3 \text{ TeV}$  (domain of applicability)










**True** resonances will have the phase shift pass through  $+\pi/2$

$$\delta_{IJ} = \tan^{-1} \left( \frac{\text{Im } t_{IJ}}{\text{Re } t_{IJ}} \right)$$

This method is known to work remarkably well in strong interactions:  $\pi\pi$  scattering  $\Rightarrow$   $\sigma$  and  $\rho$  meson masses and widths



# In hadronic physics



Truong '89, Truong, Dobado, Herrero, '90, Dobado, Pelaez, '93, '96

Integrating out a heavy higgs and recovering the resonance:

	Tree level		1-loop ( $\mu = M_W$ )		1-loop ( $\mu = M_H$ )	
MH	Mass	Width	Mass	Width	Mass	Width
500	446	28	442	27	455	30
600	521	46	511	43	538	50
700	599	70	580	63	627	81
800	680	105	647	90	726	133
900	767	157	712	121	840	201
1000	862	226	776	164	980	351
1100	968	344	838	210	1166	617
1200	1091	485	899	260	1460	1868

$$a_4 = \frac{-1}{16\pi^2} \frac{1}{12} \left( \Delta_\epsilon - \log \left( \frac{M_H^2}{\mu^2} \right) + \frac{17}{6} \right)$$

$$a_5 = \frac{M_W^2}{2g^2 M_H^2} - \frac{1}{16\pi^2} \frac{1}{24} \left( \Delta_\epsilon - \log \left( \frac{M_H^2}{\mu^2} \right) + \frac{79}{3} - \frac{27\pi}{2\sqrt{3}} \right)$$

*(On-shell scheme)*

See also recent work by Corbett, Eboli, Gonzalez-Garcia

Is this unitarization method unique?

*No, it is not. Many methods exist: IAM, K-matrix approach, N/D expansions, Roy equations,...*

*While the quantitative results differ slightly, the gross picture does not change*

For a very detailed discussion of different methods see 1502.0484 (Delgado et al.)

Is this unitarization method unique?

*No, it is not. Many methods exist: IAM, K-matrix approach, N/D expansions, Roy equations,...*

*While the quantitative results differ slightly, the gross picture does not change*

For a very detailed discussion of different methods see 1502.0484 (Delgado et al.)


Is this unitarization method unique?

*No, it is not. Many methods exist: IAM, K-matrix approach, N/D expansions, Roy equations,...*

*While the quantitative results differ slightly, the gross picture does not change*

For a very detailed discussion of different methods see 1502.0484 (Delgado et al.)

Real problem: one-loop calculation *extremely difficult*



+

*more than  
1000 other  
diagrams!*

Denner & Hahn (1998) [hep-ph/9711302]

$$t_{IJ}^{(4)} = \text{Re } t_{IJ}^{(4)} + i\text{Im } t_{IJ}^{(4)}$$

The **Optical Theorem** implies the *perturbative* relation

$$\text{Im } t_{IJ}^{(4)}(s) = \underbrace{\sigma(s)}_{\text{one-loop}} |t_{IJ}^{(2)}(s)|^2 + \underbrace{\sigma_H(s)}_{\text{tree}} |t_{H,IJ}^{(2)}(s)|^2$$




For *real part*, note that

$$\begin{aligned} \text{Re } t_{IJ}^{(4)} &= a_i\text{-dependent terms} + \text{real part of loop calculation} \\ &\approx a_i\text{-dependent terms} \\ &\quad (\text{for large } s, a_i) \end{aligned}$$

We approximate *real part of loop contribution* with **one-loop Goldstone boson amplitudes** using the **Equivalence Theorem**. In addition, we neglect coupled channels (justified in as much as  $a^2 - b$  is zero or very small in all cases).

*(Results from the Madrid group further justify this assumption)*

# Unitarity checks




Vector channel ( $a_4 = 0.008$ ,  $a_5 = 0$ )



# Are there genuine EWSBS resonances?

Must search for poles in the second Riemann sheet — the phase shift must go through  $+\pi/2$  at the resonance.

Are there any *physically acceptable* resonances?




The blue-shaded area leads to *acausal resonances*. These values for  $a_4$  and  $a_5$  are *unphysical* — they cannot be realized in any effective theory with a meaningful UV completion.

# Are there genuine EWSBS resonances?


Must search for poles in the second Riemann sheet — the phase shift must go through  $+\pi/2$  at the resonance.

Are there any *physically acceptable* resonances?



The blue-shaded area leads to *acausal resonances*. These values for  $a_4$  and  $a_5$  are *unphysical* — they cannot be realized in any effective theory with a meaningful UV completion.

# Are these resonances detectable?





Signal of IAM scalar/vector vs. SM Higgs of *same mass*

The large contribution that the SM Higgs represents leaves little room for additional resonances.



Note: only in  $WW \rightarrow WW$  or  $WW \rightarrow ZZ$  channels!

# Scalar properties



- $M_S \sim 300 - 3000$  GeV
- $\Gamma_S \sim 5 - 120$  GeV

# Vector properties




- $M_V \sim 550 - 2300$  GeV
- $\Gamma_V \sim 2 - 24$  GeV

There are constraints on vector masses from  $S, T, U$  parameter constraints in some models. Typically  $M_V > 1.5$  TeV. *e.g. Pich, Rosell, Sanz-Cillero, 2013.*

# Resonances and bounds on the anomalous QGC $a_4$ and $a_5$


Allowed regions for  $a_i$  (in white) if no resonance below 600 GeV




$$a = 1/0.95/0.9 \text{ with } a^2 = b$$

Current direct limits:  $a_4 \in [0.094, 0.10]$ ;  $a_5 \in [0.23, 0.26]$

# Spectrum of resonances for $a > 1$



'Something' happens when  $a > 1$ ... (Falkowski, Rychkov, Urbano [2012]; Espriu, Mescia [2014]; Bellazzini, Martucci, Torre [2014])



Left: Generating an  $IJ = 02$  resonance in the electroweak sector is possible with adequate values of  $a_4, a_5$ .

Right: positive values of  $a^2 - 1$  generate an isotensor  $I = 2$  resonance in the ET approximation (but not when the 'exact' amplitudes are used).

In hadron physics the isotensor wave is repulsive, and thus, not resonant.



Assuming the strict ET for all  $s$

$$1 - a^2 = \frac{v^2}{6\pi} \int_0^\infty \frac{ds}{s} (2\sigma_{l=0}(s)^{tot} + 3\sigma_{l=1}(s)^{tot} - 5\sigma_{l=2}(s)^{tot}),$$

*(Falkowski, Rychkov and Urbano, 2012)*

However, we have seen that the analytic structure of the real amplitudes is more complex. Then

- LHS is modified to  $3 - a^2 + \mathcal{O}(g^2)$
- The integral along the  $|s| \rightarrow \infty$  does not necessarily vanish.  
*(Bellazzini, Martucci and Torre, 2014)*
- LHS gets renormalized while RHS does not...

Assuming the strict ET for all  $s$

$$1 - a^2 = \frac{v^2}{6\pi} \int_0^\infty \frac{ds}{s} (2\sigma_{l=0}(s)^{tot} + 3\sigma_{l=1}(s)^{tot} - 5\sigma_{l=2}(s)^{tot}),$$

*(Falkowski, Rychkov and Urbano, 2012)*

However, we have seen that the analytic structure of the real amplitudes is more complex. Then

- LHS is modified to  $3 - a^2 + \mathcal{O}(g^2)$
- The integral along the  $|s| \rightarrow \infty$  does not necessarily vanish.  
*(Bellazzini, Martucci and Torre, 2014)*
- LHS gets renormalized while RHS does not...

Forgetting about  $O(g^2)$  corrections the proper SR reads:

$$3 - a^2 = \frac{v^2}{6\pi} \int_0^\infty \frac{ds}{s} (2\sigma_{l=0}(s)^{tot} + 3\sigma_{l=1}(s)^{tot} - 5\sigma_{l=2}(s)^{tot}) + c_\infty$$

The difference between  $1 - a^2$  and  $3 - a^2$  can be traced back to the inclusion of  $W$  exchange in the  $t$ -channel.

If the propagating degrees of freedom remain *unchanged* all the way to  $s = \infty$  (big 'if' !) the  $W$   $t$ -channel contributes to the exterior circuit and gives  $c_\infty = 2$  and restores the  $1 - a^2$  on the LHS of the sum rule obtained when  $W$  propagation is ignored.

*A trivial consequence of Cauchy's theorem*

However, there is no guarantee that the propagating degrees of freedom remain *unchanged*. In fact in strongly interacting theories one might expect  $c_\infty = 0$

Forgetting about  $O(g^2)$  corrections the proper SR reads:

$$3 - a^2 = \frac{v^2}{6\pi} \int_0^\infty \frac{ds}{s} (2\sigma_{l=0}(s)^{tot} + 3\sigma_{l=1}(s)^{tot} - 5\sigma_{l=2}(s)^{tot}) + c_\infty$$

The difference between  $1 - a^2$  and  $3 - a^2$  can be traced back to the inclusion of  $W$  exchange in the  $t$ -channel.

If the propagating degrees of freedom remain *unchanged* all the way to  $s = \infty$  (**big 'if' !**) the  $W$   $t$ -channel contributes to the exterior circuit and gives  $c_\infty = 2$  and restores the  $1 - a^2$  on the LHS of the sum rule obtained when  $W$  propagation is ignored.

*A trivial consequence of Cauchy's theorem*

However, there is no guarantee that the propagating degrees of freedom remain *unchanged*. In fact in strongly interacting theories one might expect  $c_\infty = 0$

Forgetting about  $O(g^2)$  corrections the proper SR reads:

$$3 - a^2 = \frac{v^2}{6\pi} \int_0^\infty \frac{ds}{s} (2\sigma_{l=0}(s)^{tot} + 3\sigma_{l=1}(s)^{tot} - 5\sigma_{l=2}(s)^{tot}) + c_\infty$$

The difference between  $1 - a^2$  and  $3 - a^2$  can be traced back to the inclusion of  $W$  exchange in the  $t$ -channel.



If the propagating degrees of freedom remain *unchanged* all the way to  $s = \infty$  (big 'if' !) the  $W$   $t$ -channel contributes to the exterior circuit and gives  $c_\infty = 2$  and restores the  $1 - a^2$  on the LHS of the sum rule obtained when  $W$  propagation is ignored.

*A trivial consequence of Cauchy's theorem*




However, there is no guarantee that the propagating degrees of freedom remain *unchanged*. In fact in strongly interacting theories one might expect  $c_\infty = 0$

# Implications for the EWSB if $M_R \sim 2$ TeV

Reduction of parameter space if  $1800 \text{ GeV} \leq M_R \leq 2200 \text{ GeV}$



# Blow-up and widths



# Resonance signal

We can estimate how observable these signals are by comparing to a heavy SM Higgs of the same mass  $\rightarrow$  [look at LHC Higgs search data](#)

For a resonance of mass  $M_R$  and width  $\Gamma_R$ , let

$$\begin{aligned}\sigma^{peak} &\equiv \int_{M_R-2\Gamma_R}^{M_R+2\Gamma_R} \left[ dM_{WW} \times \frac{d\sigma}{dM_{WW}} \right] \\ \sigma_{SM}^{peak} &\equiv \int_{M_H-2\Gamma_H}^{M_H+2\Gamma_H} \left[ dM_{WW} \times \frac{d\sigma_{SM}}{dM_{WW}} \right]\end{aligned}$$



Then for a heavy Higgs with  $M_H \rightarrow M_R$  and  $\Gamma_H(M_H \rightarrow M_R)$

$$R \equiv \left( \frac{\sigma^{peak}}{\sigma_{SM}^{peak}} \right)$$

compares the strength of the resonance regions of the same mass.




ATLAS results suggest cross-sections around  $10 \text{ fb}$ .  
Channel identification is dubious.



WW channel can be mediated by  $I=1,2$ . ZZ channel can be mediated by  $I=0,2$ . (But  $I = 2$  .....

*Only resonant contributions to the cross-section is shown.*

# Dependence of $ZZ$ cross-section on $a$



Changes in  $a$  are insufficient to increase substantially the cross-section

# DY: anomalous $\bar{q}_L q_L$ coupling




Production of a pair of Goldstone bosons by  $\bar{q}q'$  annihilation through a  $W$ -meson and anomalous BSM vertex enhancing it

Gauge invariant operator:  $\delta_1 \bar{\psi}_L U \not{D} U^\dagger \psi_L$   
(one of several  $D = 4$  Longhitano's operators)

*Changes the relation between  $G_F$  and the TGB vertex*

# Effect of an anomalous $\bar{q}_L q_L$ coupling



$$\left[ \frac{d\hat{\sigma}}{d\Omega} \right]_{\text{cm}} = \frac{1}{64\pi^2 s} \frac{g^4}{32} \sin^2 \theta \left( 1 + \frac{\delta_1 s}{v^2} \right)^2 |F_V(s)|^2 ,$$


$F_V$  : Form factor (Watson's theorem)

# Introducing form factors

$$\langle 0|V_\mu|\omega(q)\omega(q')\rangle = iF_V^+(q+q')_\mu + iF_V^-(q-q')_\mu$$

CVC  $\Rightarrow F_V^+ = 0$ . Unitarity implies


$$\text{Im}F_V = F_V t^* \quad t = A(\omega\omega \rightarrow \omega\omega) \quad K_V \equiv (s - M_V)^2 F_V$$



Then (symbolically)

$$t = K_V \frac{1}{s - M_V^2} K_V^* = \frac{t^{(2)}}{1 - t^{(4)}/t^{(2)}}$$

(Note: for  $l=1$   $u$  and  $t$  channels cancel each other)



Assuming  $g_V V_\mu^i \partial^\mu w^j w^k \epsilon_{ijk} \Rightarrow g_V(\sqrt{s} = 2 \text{ TeV}) \simeq 1.6$

- Unitarity is a powerful constraint on scattering amplitudes. Its validity is well tested.
- Even in the presence of a light Higgs, it can help constrain anomalous couplings by helping predict heavier resonances.
- An extended EWSBS would typically have such resonances even in the presence of a light 'Higgs'
- However their properties are radically different from the 'naive expectations'
- Current LHC Higgs search results do not yet probe the IAM resonances: at least  $10\times$  statistics is required

Cross sections are at least one order of magnitude too small to explain the 'resonance'

However:

- Near-degeneracy of scalar and vector gives also enhancement (also helps to explain the signal in *all* 3 channels)
- Departures from  $a = 1$  enhances the scalar signal.
  - Ex: going from  $a = 1$  to  $a = 0.95$  doubles the cross-section
  - Current experimental limit (90 % CL):  $a \in [0.67, 1.33]$
  - In some models  $a < 1$  implies breaking of custodial symmetry  $\Rightarrow$  scalar isotriplets can exist, coupling proportionally to the custodial breaking parameter.
- Direct coupling of the resonances to quarks? (Drell-Yan)


All this is under current investigation!




*THANK YOU!*

# Back-up slides


# Comparing ET and 'exact' calculations



(a) Scalar




(b) Vector




(c) Tensor

# Comparison Higgsless/Higgs with $M_H = 125$ GeV



No light Higgs



With light Higgs

Here  $\sqrt{s} = 8$  TeV

Compare before/after for same point (ex: Point D  $a_4 = 0.008$ ,  $a_5 = 0.000$ )

- Different continuum
- Masses have changed positions
- Widths are *much narrower*

# General $hWW$ couplings

For  $a = b = 1$  these results reproduce the **SM** prediction, i.e. no counterterms (renormalizable theory)

$$\delta a_4 = 0, \quad \delta a_5 = 0$$

For  $a = b = 0$  one gets the 'no Higgs' results (**EChL**)


$$\delta a_4 = \Delta_\epsilon \frac{1}{(4\pi)^2} \frac{-1}{12}, \quad \delta a_5 = \Delta_\epsilon \frac{1}{(4\pi)^2} \frac{-1}{24}$$

They should bring 'natural' finite contributions from NP:


$$a_4|_{\text{finite}} \simeq \frac{1}{(4\pi)^2} \frac{-1}{12} (1 - a^2)^2 \log \frac{v^2}{f^2}$$

$$a_5|_{\text{finite}} \simeq \frac{1}{(4\pi)^2} \frac{-1}{24} \left[ (1 - a^2)^2 + \frac{3}{2} ((1 - a^2) - (1 - b))^2 \right] \log \frac{v^2}{f^2},$$

# Bounds on $a_i$ for $a = 0.8$ ( $b = a^2$ )






All Resonances



$M_R < 600$  GeV

# Some diseases



$$\begin{aligned}
 \mathcal{L}_{QGC} = & e^2 \left[ g_1^{\gamma\gamma} A^\mu A^\nu W_\mu^- W_\nu^+ - g_2^{\gamma\gamma} A^\mu A_\mu W^{-\nu} W_\nu^+ \right] \\
 & + e^2 \frac{c_w}{s_w} \left[ g_1^{\gamma Z} A^\mu Z^\nu \left( W_\mu^- W_\nu^+ + W_\mu^+ W_\nu^- \right) - 2g_2^{\gamma Z} A^\mu Z_\mu W^{-\nu} W_\nu^+ \right] \\
 & + e^2 \frac{c_w^2}{s_w^2} \left[ g_1^{ZZ} Z^\mu Z^\nu W_\mu^- W_\nu^+ - g_2^{ZZ} Z^\mu Z_\mu W^{-\nu} W_\nu^+ \right] \\
 & + \frac{e^2}{2s_w^2} \left[ g_1^{WW} W^{-\mu} W^{+\nu} W_\mu^- W_\nu^+ - g_2^{WW} \left( W^{-\mu} W_\mu^+ \right)^2 \right] + \frac{e^2}{4s_w^2 c_w^4} h^{ZZ} (Z^\mu Z_\mu)^2
 \end{aligned}$$

SM values:  $g_1^{\gamma,Z} = \kappa^{\gamma,Z} = 1$ ,  $\lambda^{\gamma,Z} = 0$  and  $\delta_Z = \frac{\beta_1 + g'^2 \alpha_1}{c_w^2 - s_w^2}$ ,  $g_{1/2}^{VV'} = 1$ ,  $h^{ZZ} = 0$

$$\Delta g_1^\gamma = 0$$

$$\Delta \kappa^\gamma = g^2(\alpha_2 - \alpha_1) + g^2 \alpha_3 + g^2(\alpha_9 - \alpha_8)$$

$$\Delta g_1^Z = \delta_Z + \frac{g^2}{c_w^2} \alpha_3$$

$$\Delta \kappa^Z = \delta_Z - g'^2(\alpha_2 - \alpha_1) + g^2 \alpha_3 + g^2(\alpha_9 - \alpha_8)$$

$$\Delta g_1^{\gamma\gamma} = \Delta g_2^{\gamma\gamma} = 0$$

$$\Delta g_2^{ZZ} = 2\Delta g_1^{\gamma Z} - \frac{g^2}{c_w^4}(\alpha_5 + \alpha_7)$$

$$\Delta g_1^{\gamma Z} = \Delta g_2^{\gamma Z} = \delta_Z + \frac{g^2}{c_w^2} \alpha_3$$

$$\Delta g_1^{WW} = 2c_w^2 \Delta g_1^{\gamma Z} + 2g^2(\alpha_9 - \alpha_8) + g^2 \alpha_4$$


$$\Delta g_1^{ZZ} = 2\Delta g_1^{\gamma Z} + \frac{g^2}{c_w^4}(\alpha_4 + \alpha_6)$$

$$\Delta g_2^{WW} = 2c_w^2 \Delta g_1^{\gamma Z} + 2g^2(\alpha_9 - \alpha_8) - g^2(\alpha_4 + 2\alpha_5)$$

$$h^{ZZ} = g^2[\alpha_4 + \alpha_5 + 2(\alpha_6 + \alpha_7 + \alpha_{10})]$$




# Hard to detect!



*Scalar*

$R \sim 0.0001 - 0.30$

*Typically < 0.1*



*Vector*

$R \sim 0.0003 - 0.3$

*Typically  $\sim 0.2$*

They *could still be there*, but would give a small signal.