

# $g - 2$ theory review

Marc Knecht

Centre de Physique Théorique UMR7332,  
CNRS Luminy Case 907, 13288 Marseille cedex 09 - France  
[knecht@cpt.univ-mrs.fr](mailto:knecht@cpt.univ-mrs.fr)

---

4th Workshop on Muon  $g - 2$  and EDM in the LHC Era – CPT-Marseille , May 23 2016




## OUTLINE

- Introduction
- Perturbative contributions: QED and EW
- Non perturbative contributions QCD
  - HVP (LO+NLO+NNLO)
  - HLxL
- Conclusions

# Introduction

General context: looking for new physics by exploring the “three frontiers”



In this presentation, the focus will be on the intensity (or precision) frontier, illustrated through the anomalous magnetic moments of the electron and the muon

## Leptonic sector of the three-family standard model

- charged leptons:  $\ell = e^\pm, \mu^\pm, \tau^\pm$

$$q = \pm 1 \text{ (charge)} \quad s = \frac{1}{2} \text{ (spin)}$$

differ only in their masses

$$m_e = 0.510\,998\,910(13) \text{ MeV}$$

$$m_\mu = 105.658\,367(4) \text{ MeV}$$

$$m_\tau = 1\,776.82(16) \text{ MeV}$$

- neutral leptons:  $\nu_\ell = \nu_e, \nu_\mu, \nu_\tau$

$$q = 0 \quad s = \frac{1}{2}$$

## Response of a charged lepton to an external (and static) electromagnetic field

$$\begin{aligned} \langle \ell; p' | \mathcal{J}_\rho(0) | \ell; p \rangle &\equiv \bar{u}(p') \Gamma_\rho(p', p) u(p) \\ &= \bar{u}(p') \left[ F_1(k^2) \gamma_\rho + \frac{i}{2m_\ell} F_2(k^2) \sigma_{\rho\nu} k^\nu - F_3(k^2) \gamma_5 \sigma_{\rho\nu} k^\nu + F_4(k^2) (k^2 \gamma_\rho - 2m_\ell k_\rho) \gamma_5 \right] u(p) \end{aligned}$$

(uses only the conservation of the electromagnetic current  $\mathcal{J}_\rho$ ,  $k_\mu \equiv p'_\mu - p_\mu$ )

$F_1(k^2)$  → Dirac form factor,  $F_1(0) = 1$

$F_2(k^2)$  → Pauli form factor →  $F_2(0) = a_\ell$

$F_3(k^2)$  →  $P, T$ , electric dipole moment →  $F_3(0) = d_\ell/e_\ell$

$F_4(k^2)$  →  $P$ , anapole moment

$$G_E(k^2) = F_1(k^2) + \frac{k^2}{4m_\ell^2} F_2(k^2), \quad G_M(k^2) = F_1(k^2) + F_2(k^2)$$

$$G_E(0) = 1 \qquad \qquad \qquad G_M(0) = \frac{g_\ell}{2}$$

in the SM,  $F_2(k^2)$ ,  $F_3(k^2)$ ,  $F_4(k^2)$  are only generated by loops → **calculable!**

## Response of a charged lepton to an external (and static) electromagnetic field

For a relativistic, point-like spin 1/2 particle, described by the Dirac equation with the minimal coupling prescription, one has

$$i\hbar \frac{\partial \psi}{\partial t} = \left[ c\boldsymbol{\alpha} \cdot \left( -i\hbar \nabla - \frac{e_\ell}{c} \mathbf{A} \right) + \beta m_\ell c^2 + e_\ell \mathcal{A}_0 \right] \psi$$

In the non relativistic limit, this reduces to the Pauli equation for the two-component spinor  $\varphi$  describing the large components of the Dirac spinor  $\psi$ ,

$$i\hbar \frac{\partial \varphi}{\partial t} = \left[ \frac{(-i\hbar \nabla - (e_\ell/c) \mathbf{A})^2}{2m_\ell} - \underbrace{\frac{e_\ell \hbar}{2m_\ell c} \boldsymbol{\sigma} \cdot \mathbf{B}}_{\mu_\ell \cdot \mathbf{B}} + e_\ell \mathcal{A}_0 \right] \varphi$$

with

$$\mu_\ell = g_\ell \left( \frac{e_\ell}{2m_\ell c} \right) \mathbf{s}, \quad \mathbf{s} = \hbar \frac{\boldsymbol{\sigma}}{2}, \quad g_\ell^{\text{Dirac}} = 2$$


At tree level,  $g_\ell = g_\ell^{\text{Dirac}} \equiv 2$ .

The *anomalous* magnetic moment is induced at loop level:


$$a_\ell \equiv \frac{g_\ell - g_\ell^{\text{Dirac}}}{g_\ell^{\text{Dirac}}} [\equiv F_2(0)]$$

$a_\ell$  probes all the degrees of freedom of the standard model, *and possibly beyond...*

*... provided one can measure it with enough precision !*




*... provided one can measure it with enough precision !*



*And predict its value in the standard model with a comparable accuracy...*

$a_e$  and  $a_\mu$  are among the most precisely measured SM observables:



$$a_e^{\text{exp}} = 1159652180.73(0.28) \cdot 10^{-12}$$

$$\Delta a_e^{\text{exp}} = 2.8 \cdot 10^{-13} \text{ [0.24ppb]}$$

[D. Hanneke et al, Phys. Rev. Lett. 100, 120801 (2008)]

$$\tau_\mu = (2.19703 \pm 0.00004) \times 10^{-6} \text{ s}$$

$$\gamma \sim 29.3, p \sim 3.094 \text{ GeV/c}$$

$$a_\mu^{\text{exp}} = 11659209.1(6.3) \cdot 10^{-10}$$

$$\Delta a_\mu^{\text{exp}} = 6.3 \cdot 10^{-10} \text{ [0.54ppm]}$$

[G. W. Bennett et al, Phys Rev D 73, 072003 (2006)]



Are the SM calculations able to achieve a comparable level of accuracy ? General structure  $a_\ell^{\text{SM}} = a_\ell^{\text{QED}} + a_\ell^{\text{had}} + a_\ell^{\text{weak}}$

Table 1: Summary of  $a_\mu$  results from CERN and BNL, showing the evolution of experimental precision over time. The average is obtained from the 1999, 2000 and 2001 data sets only.

Experiment	Years	Polarity	$a_\mu \times 10^{10}$	Precision [ppm]
CERN I	1961	$\mu^+$	11 450 000(220 000)	4300
CERN II	1962-1968	$\mu^+$	11 661 600(3100)	270
CERN III	1974-1976	$\mu^+$	11 659 100(110)	10
CERN III	1975-1976	$\mu^-$	11 659 360(120)	10
BNL E821	1997	$\mu^+$	11 659 251(150)	13
BNL E821	1998	$\mu^+$	11 659 191(59)	5
BNL E821	1999	$\mu^+$	11 659 202(15)	1.3
BNL E821	2000	$\mu^+$	11 659 204(9)	0.73
BNL E821	2001	$\mu^-$	11 659 214(9)	0.72

$$a_\mu^{exp} = 11 659 209.1(5.4)(3.3) \cdot 10^{-10} [0.54 \text{ ppm}]$$

## The tau case

$$\tau_\tau = (290.6 \pm 1.1) \times 10^{-15} \text{ s}$$

- $e^+e^- \rightarrow \tau^+\tau^-\gamma$

$$-0.052 < a_\tau^{exp} < +0.058 \text{ (L3, 1998, 95% CL)}$$

[Phys. Lett. B 434, 169 (1998)]

$$-0.068 < a_\tau^{exp} < +0.065 \text{ (OPAL, 1998, 95% CL)}$$

[Phys. Lett. B 431, 188 (1998)]

- $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$

$$-0.052 < a_\tau^{exp} < +0.013 \text{ (DELPHI, 2004, 95% CL)}$$

$$a_\tau^{exp} = -0.018(17)$$

[Eur. Phys. J. C 35, 159 (2004)]

- Reanalysis of experiments  $\rightarrow -0.007 < a_\tau^{exp} < +0.005$

[G. A. Gonzalez-Sprinberg, A. Santamaria, J. Vidal, Nucl. Phys. B 582, 3 (2000) [hep-ph/0002203]]

- $\tau^- \rightarrow \ell^-\nu_\tau\bar{\nu}_\ell\gamma$  at B factories  $\rightarrow \Delta a_\tau \sim 0.012$

[S. Eidelman, D. Epifanov, M. Fael, L. Mercolli, M. Passera, arXiv:1601.07987 [hep-ph]]

- theory:  $a_\tau = 117721(5) \cdot 10^{-8}$  [42 ppm]

[S. Eidelman, M. Passera, Mod. Phys. Lett. A 22, 159 (2007)]

[S. Narison, Phys Lett B 513 (2001); err. B 526, 414 (2002)]

# Theory

Useful guidelines:

- Within the framework of a renormalizable quantum field theory,  $F_2(k^2)$ ,  $F_3(k^2)$  and  $F_4(k^2)$  can only arise through loop corrections. These loop contributions have to be finite and calculable, since the possible counterterms correspond to non renormalizable interactions
- $a_\ell$  is dimensionless: the contributions from loops involving only photons and the lepton  $\ell$  (one-flavour QED) are mass independent and thus universal
- Massive degrees of freedom with  $M \gg m_\ell$  contribute to  $a_\ell$  through powers of  $m_\ell^2/M^2$  times logarithms (decoupling)
- Light degrees of freedom with  $m \ll m_\ell$  give logarithmic contributions to  $a_\ell$ , e.g.  $\ln(m_\ell^2/m^2) \left( \pi^2 \ln \frac{m_\mu}{m_e} \sim 50 \right)$

# Theory I: QED ( $a_e$ and $a_\mu$ )

- QED contributions : loops with only photons and leptons

$$a_\ell^{\text{QED}} = C_\ell^{(2)} \left(\frac{\alpha}{\pi}\right) + C_\ell^{(4)} \left(\frac{\alpha}{\pi}\right)^2 + C_\ell^{(6)} \left(\frac{\alpha}{\pi}\right)^3 + C_\ell^{(8)} \left(\frac{\alpha}{\pi}\right)^4 + C_\ell^{(10)} \left(\frac{\alpha}{\pi}\right)^5 + \dots$$


$$C_\ell^{(2n)} = A_1^{(2n)} + A_2^{(2n)}(m_\ell/m_{\ell'}) + A_3^{(2n)}(m_\ell/m_{\ell'}, m_\ell/m_{\ell''})$$

$A_1^{(2n)}$   $\longrightarrow$  mass-independent (universal) contributions (one-flavour QED)

$A_2^{(2n)}(m_\ell/m_{\ell'}), A_3^{(2n)}(m_\ell/m_{\ell'}, m_\ell/m_{\ell''}) \longrightarrow$   
mass-dependent (non-universal) contributions (multi-flavour QED)


$a_\ell$  is finite (no renormalization needed) and dimensionless

Analytic expressions for  $A_1^{(2)}$ ,  $A_1^{(4)}$ ,  $A_1^{(6)}$ ,  $A_2^{(4)}$ ,  $A_2^{(6)}$ ,  $A_3^{(6)}$  known



$$A_1^{(2)} = \frac{1}{2}$$

[J. Schwinger, Phys. Rev. 73, 416L (1948)]



$$A_1^{(4)} = \frac{3}{4}\zeta(3) - \frac{\pi^2}{2} \ln 2 + \frac{\pi^2}{12} + \frac{197}{144} = -0.328\,478\,965\,579\,193\dots$$

[C. M. Sommerfield, Phys. Rev. 107, 328 (1957); Ann. Phys. 5, 26 (1958)]

[A. Petermann, Helv. Phys. Acta 30, 407 (1957)]

$$A_2^{(4)}(m_\ell/m_{\ell'}) = \frac{1}{3} \int_{4m_{\ell'}^2}^\infty dt \sqrt{1 - \frac{4m_{\ell'}^2}{t}} \frac{t + 2m_{\ell'}^2}{t^2} \int_0^1 dx \frac{x^2(1-x)}{x^2 + (1-x) \frac{t}{m_\ell^2}}$$

[H. Suura and E. Wichmann, Phys. Rev. 105, 1930 (1955)]

[A. Petermann, Phys. Rev. 105, 1931 (1955)]

[H. H. Elend, Phys. Lett. 20, 682 (1966); Err. Ibid. 21, 720 (1966)]

[M. Passera, Phys. Rev. D 75, 013002 (2007)]

$$\begin{aligned} A_2^{(4)}(m_\ell/m_{\ell'}) &= \frac{1}{3} \ln \left( \frac{m_\ell}{m_{\ell'}} \right) - \frac{25}{36} + \frac{\pi^2}{4} \frac{m_{\ell'}}{m_\ell} - 4 \left( \frac{m_{\ell'}}{m_\ell} \right)^2 \ln \left( \frac{m_\ell}{m_{\ell'}} \right) \\ &\quad + 3 \left( \frac{m_{\ell'}}{m_\ell} \right)^2 + \mathcal{O} \left[ \left( \frac{m_{\ell'}}{m_\ell} \right)^3 \right], \text{ red } m_\ell \gg m_{\ell'} \end{aligned}$$

[M. A. Samuel and G. Li, Phys. Rev. D 44, 3935 (1991)]

$$A_2^{(4)}(m_\mu/m_e) = 1.094\,258\,312\,0(83)$$

$$m_\mu/m_e = 206.768\,2843(52)$$

P. J. Mohr, B. N. Taylor, D. B. Newell, CODATA 2010, arXiv:1203.5425v1[physics.atom-ph]

$$A_2^{(4)}(m_\ell/m_{\ell'}) = \frac{1}{3} \int_{4m_{\ell'}^2}^\infty dt \sqrt{1 - \frac{4m_{\ell'}^2}{t}} \frac{t + 2m_{\ell'}^2}{t^2} \int_0^1 dx \frac{x^2(1-x)}{x^2 + (1-x) \frac{t}{m_\ell^2}}$$

[H. Suura and E. Wichmann, Phys. Rev. 105, 1930 (1955)]

[A. Petermann, Phys. Rev. 105, 1931 (1955)]

[H. H. Elend, Phys. Lett. 20, 682 (1966); Err. Ibid. 21, 720 (1966)]

[M. Passera, Phys. Rev. D 75, 013002 (2007)]

$$\begin{aligned} A_2^{(4)}(m_\ell/m_{\ell'}) &= \frac{1}{45} \left( \frac{m_\ell}{m_{\ell'}} \right)^2 + \frac{1}{70} \left( \frac{m_\ell}{m_{\ell'}} \right)^4 \ln \left( \frac{m_\ell}{m_{\ell'}} \right) \\ &\quad + \frac{9}{19600} \left( \frac{m_\ell}{m_{\ell'}} \right)^4 + \mathcal{O} \left[ \left( \frac{m_\ell}{m_{\ell'}} \right)^3 \ln \left( \frac{m_\ell}{m_{\ell'}} \right) \right], \text{ red } m_{\ell'} \gg m_\ell \end{aligned}$$

[B.E. Lautrup and E. de Rafael, Phys. Rev. 174, 1835 (1965)]

[M. A. Samuel and G. Li, Phys. Rev. D 44, 3935 (1991)]


$$A_2^{(4)}(m_e/m_\mu) = 5.197\,386\,67(26) \cdot 10^{-7}$$

$$A_2^{(4)}(m_e/m_\tau) = 1.837\,98(34) \cdot 10^{-9}$$

$$A_2^{(4)}(m_\mu/m_\tau) = 7.8079(15) \cdot 10^{-5}$$

$$m_\mu/m_\tau = 5.946\,49(54) \cdot 10^{-2} \quad m_e/m_\tau = 2.875\,92(26) \cdot 10^{-4}$$

order  $(\alpha/\pi)^3$ : 72 diagrams



$$\begin{aligned}
 A_1^{(6)} = & \frac{87}{72} \pi^2 \zeta(3) - \frac{215}{24} \zeta(5) + \frac{100}{3} \left[ \left( a_4 + \frac{1}{24} \ln^4 2 \right) - \frac{1}{24} \pi^2 \ln^2 2 \right] - \frac{239}{2160} \pi^4 \\
 & + \frac{139}{18} \zeta(3) - \frac{298}{9} \pi^2 \ln 2 + \frac{17101}{810} \pi^2 + \frac{28259}{5184} \quad [a_p = \sum_1^\infty 1/(2^n n^p)]
 \end{aligned}$$

[S. Laporta, E. Remiddi, Phys. Lett. B265, 182 (1991); B356, 390 (1995); B379, 283 (1996)]

[S. Laporta, Phys. Rev. D 47, 4793 (1993); Phys. Lett. B343, 421 (1995)]

$$A_1^{(6)} = 1.181\,241\,456\dots$$

numerical evaluations:  $A_1^{(6)}(\text{num}) = 1.181\,259(40)\dots$

[T. Kinoshita, Phys. Rev. Lett. 75, 4728 (1995)]

order  $(\alpha/\pi)^4$ : 891 diagrams

only some diagrams are known analytically →

Automated generation of diagrams

Systematic numerical evaluation of multi-dimensional integrals over Feynman-parameter space

$$A_1^{(8)} = -1.912\,98(84)$$

$$A_2^{(8)}(m_e/m_\mu) = 9.161\,970\,703(373) \cdot 10^{-4} \quad A_2^{(8)}(m_e/m_\tau) = 7.429\,24(118) \cdot 10^{-6}$$

$$A_3^{(8)}(m_e/m_\mu, m_e/m_\tau) = 7.468\,7(28) \cdot 10^{-7}$$

$$A_2^{(8)}(m_\mu/m_e) = 132.685\,2(60) \quad A_2^{(8)}(m_\mu/m_\tau) = 0.042\,34(12)$$

$$A_3^{(8)}(m_\mu/m_e, m_\mu/m_\tau) = 0.062\,72(4)$$

order  $(\alpha/\pi)^4$ : 891 diagrams

only some diagrams are known analytically →

Automated generation of diagrams

Systematic numerical evaluation of multi-dimensional integrals over Feynman-parameter space

order  $(\alpha/\pi)^5$ : 12 672 diagrams...

6 classes, 32 gauge invariant subsets


Five of these subsets are known analytically

[S. Laporta, Phys. Lett. B 328, 522 (1994)]

[J.-P. Aguilar, D. Greynat, E. de Rafael, Phys. Rev. D 77, 093010 (2008)]

Complete numerical results have been published

[T. Kinoshita and M. Nio, Phys. Rev. D 73, 053007 (2006); T. Aoyama et al., Phys. Rev. D 78, 053005 (2008); D 78, 113006 (2008); D 81, 053009 (2010); D 82, 113004 (2010); D 83, 053002 (2011); D 83, 053003 (2011); D 84, 053003 (2011); D 85, 033007 (2012); Phys. Rev. Lett. 109, 111807 (2012); Phys. Rev. Lett. 109, 111808 (2012); Phys. Rev. D 91, 033006 (2015)]



[T. Aoyama et al., Phys. Rev. D 91, 033006 (2015)]

order  $(\alpha/\pi)^4$ : 891 diagrams

only some diagrams are known analytically →

Automated generation of diagrams

Systematic numerical evaluation of multi-dimensional integrals over Feynman-parameter space

Analytic evaluation at  $\mathcal{O}(\alpha^4)$ ? →

Feynamn diagrams contributing to mass-dependent terms have been evaluated using asymptotic expansion techniques

Numerical results for  $A_2^{(8)}$  and  $A_3^{(8)}$  confirmed within the accuracy required in order to make comparison with experimental precision for  $a_\mu$

[A. Kurz, T. Liu, P. Marquard, M. Steinhauser, Nucl. Phys. B 879, 1 (2014)]

[A. Kurz, T. Liu, P. Marquard, A. V. Smirnov, V. A. Smirnov, M. Steinhauser, Phys. Rev. D 92, 073019 (2015); arXiv:1602.02785 [hep-ph]]

order  $(\alpha/\pi)^4$ : 891 diagrams

no independent check for  $A_1^{(8)} = -1.91298(84)$  so far...

$A_1^{(8)}$	$= -1.434(138)$	[Kinoshita and Lindquist (1990)]
	$= -1.557(70)$	[Kinoshita (1995)]
	$= -1.4092(384)$	[Kinoshita (1997)]
	$= -1.5098(384)$	[Kinoshita (2001)]
	$= -1.7366(60)$	[Kinoshita (2005)]
	$= -1.7260(50)$	[Kinoshita (2005)]
	$= -1.7283(35)$	[Kinoshita and Nio, Phys. Rev. D 73, 013003(2006)]
	$= -1.9144(35)$	[Aoyama et al., Phys. Rev. Lett. 99, 110406 (2007)] $\leftarrow$
	$= -1.9106(20)$	[Aoyama et al., Phys. Rev. Lett. 109, 111807 (2012)]
	$= -1.91298(84)$	[Aoyama et al., Phys. Rev. D 91, 033006 (2015)]

$$a_\ell^{\text{QED}} = C_\ell^{(2)} \left(\frac{\alpha}{\pi}\right) + C_\ell^{(4)} \left(\frac{\alpha}{\pi}\right)^2 + C_\ell^{(6)} \left(\frac{\alpha}{\pi}\right)^3 + C_\ell^{(8)} \left(\frac{\alpha}{\pi}\right)^4 + C_\ell^{(10)} \left(\frac{\alpha}{\pi}\right)^5$$

	$\ell = e$	$\ell = \mu$
$C_\ell^{(2)}$	0.5	0.5
$C_\ell^{(4)}$	$-0.328\,478\,444\,00\dots$	$0.765\,857\,425(17)$
$C_\ell^{(6)}$	$1.181\,234\,017\dots$	$24.050\,509\,96(32)$
$C_\ell^{(8)}$	$-1.912\,07(84)$	$130.877\,3(61)$
$C_\ell^{(10)}$	$7.795(336)$	$751.92(93)$

n	1	2	3	4	5
$(\alpha/\pi)^n$	$2.32\dots\cdot 10^{-3}$	$5.39\dots\cdot 10^{-6}$	$1.25\dots\cdot 10^{-8}$	$2.91\dots\cdot 10^{-11}$	$6.76\dots\cdot 10^{-14}$

$$a_\ell^{\text{QED}} = C_\ell^{(2)} \left(\frac{\alpha}{\pi}\right) + C_\ell^{(4)} \left(\frac{\alpha}{\pi}\right)^2 + C_\ell^{(6)} \left(\frac{\alpha}{\pi}\right)^3 + C_\ell^{(8)} \left(\frac{\alpha}{\pi}\right)^4 + C_\ell^{(10)} \left(\frac{\alpha}{\pi}\right)^5$$

	$\ell = e$	$\ell = \mu$
$C_\ell^{(2)}$	0.5	0.5
$C_\ell^{(4)}$	$-0.328\,478\,444\,00\dots$	$0.765\,857\,425(17)$
$C_\ell^{(6)}$	$1.181\,234\,017\dots$	$24.050\,509\,96(32)$
$C_\ell^{(8)}$	$-1.912\,07(84)$	$130.877\,3(61)$
$C_\ell^{(10)}$	$7.795(336)$	$751.92(93)$

n	1	2	3	4	5
$(\alpha/\pi)^n$	$2.32\dots\cdot 10^{-3}$	$5.39\dots\cdot 10^{-6}$	$1.25\dots\cdot 10^{-8}$	$2.91\dots\cdot 10^{-11}$	$6.76\dots\cdot 10^{-14}$

$$\Delta C_e^{(8)} \cdot (\alpha/\pi)^4 \sim 0.2 \cdot 10^{-13} \quad \Delta C_e^{(10)} \cdot (\alpha/\pi)^5 \sim 0.2 \cdot 10^{-13} \quad \Delta a_e^{\text{exp}} = 2.8 \cdot 10^{-13}$$

$$a_\ell^{\text{QED}} = C_\ell^{(2)} \left(\frac{\alpha}{\pi}\right) + C_\ell^{(4)} \left(\frac{\alpha}{\pi}\right)^2 + C_\ell^{(6)} \left(\frac{\alpha}{\pi}\right)^3 + C_\ell^{(8)} \left(\frac{\alpha}{\pi}\right)^4 + C_\ell^{(10)} \left(\frac{\alpha}{\pi}\right)^5$$

	$\ell = e$	$\ell = \mu$
$C_\ell^{(2)}$	0.5	0.5
$C_\ell^{(4)}$	$-0.328\,478\,444\,00\dots$	$0.765\,857\,425(17)$
$C_\ell^{(6)}$	$1.181\,234\,017\dots$	$24.050\,509\,96(32)$
$C_\ell^{(8)}$	$-1.912\,07(84)$	$130.877\,3(61)$
$C_\ell^{(10)}$	$7.795(336)$	$751.92(93)$

n	1	2	3	4	5
$(\alpha/\pi)^n$	$2.32\dots\cdot 10^{-3}$	$5.39\dots\cdot 10^{-6}$	$1.25\dots\cdot 10^{-8}$	$2.91\dots\cdot 10^{-11}$	$6.76\dots\cdot 10^{-14}$

$$\begin{aligned} \Delta C_\mu^{(4)} \cdot (\alpha/\pi)^2 &\sim 0.9 \cdot 10^{-13} & \Delta C_\mu^{(6)} \cdot (\alpha/\pi)^3 &\sim 0.04 \cdot 10^{-13} \\ \Delta C_\mu^{(8)} \cdot (\alpha/\pi)^4 &\sim 1.8 \cdot 10^{-13} & \Delta C_\mu^{(10)} \cdot (\alpha/\pi)^5 &\sim 0.6 \cdot 10^{-13} & \Delta a_\mu^{\text{exp}} &= 6.3 \cdot 10^{-10} \end{aligned}$$

$$C_\mu^{(8)} \cdot (\alpha/\pi)^4 \sim 3.8 \cdot 10^{-9} \quad C_\mu^{(10)} \cdot (\alpha/\pi)^5 \sim 0.5 \cdot 10^{-10}$$

## QED prediction ?

→ requires an input for the fine structure constant  $\alpha$  that matches the experimental accuracy on  $a_e$

$$\frac{\Delta a_e}{a_e} = 0.24 \text{ ppb} \rightarrow \frac{\Delta \alpha}{\alpha} \sim 0.24 \text{ ppb} \rightarrow \Delta \alpha \lesssim 2 \cdot 10^{-12}$$

- quantum Hall effect

$$\alpha^{-1}[qH] = 137.036\,00300(270) \quad [19.7 \text{ ppb}]$$

[P. J. Mohr, B. N. Taylor, D. B. Newell, Rev. Mod. Phys. 80, 633 (2008)]

- atomic recoil velocity through photon absorption

$$\alpha^2 = \frac{2R_\infty}{c} \cdot \frac{M_{\text{atom}}}{m_e} \cdot \frac{h}{M_{\text{atom}}}$$

$$\frac{\Delta R_\infty}{R_\infty} = 5 \cdot 10^{-12} \quad \Delta \left( \frac{M_{\text{Rb}}}{m_e} \right) = 4.4 \cdot 10^{-10}$$

$$\alpha^{-1}[Cs\,02] = 137.036\,0001(11) \quad [7.7\text{ppb}]$$

[A. Wicht, J. M. Hensley, E. Sarajilic, S. Chu, Phys. Scr. T102, 82 (2002)]

$$\alpha^{-1}[Rb\,06] = 137.035\,998\,84(91) \quad [6.7\text{ppb}]$$

[P. Cladé et al, Phys. Rev. A 74, 052109 (2006)]

$$\alpha^{-1}[Rb\,08] = 137.035\,999\,45(62) \quad [4.6\text{ppb}]$$

[M. Cadoret et al, Phys. Rev. Lett. 101, 230801 (2008)]


$$\alpha^{-1}[Rb\,11] = 137.035\,999\,037(91) \quad [0.66\text{ppb}]$$

[R. Bouchendira, P. Cladé, S. Ghelladi-Khélifa, F. Nez, F. Biraben, Phys. Rev. Lett. 106, 080801  
(2011)]

$$a_e^{\text{QED}} = 1\,159\,652\,179.908(25)_{\alpha^4}(23)_{\alpha^5}(763)_{\alpha(Rb11)} \cdot 10^{-12} \quad a_e^{\text{exp}} - a_e^{\text{QED}} = 0.822(813) \cdot 10^{-12}$$

$$\alpha[a_e(HV\,08)] = 137.035\,999\,157\,0(29)_{\alpha^4}(27)_{\alpha^5}(18)_{\text{had}}(331)_{\text{exp}} \quad [0.25\text{ppb}]$$

[Aoyama et al., Phys. Rev. D 91 033006 (2015)]



[R. Bouchendira, P. Cladé, S. Ghelladi-Khélifa, F. Nez, F. Biraben, Phys. Rev. Lett. 106, 080801 (2011)]

$$a_\mu^{\text{QED}}(Rb) = 1\,165\,847\,189.51(9)_\text{mass}(19)_{\alpha^4}(7)_{\alpha^5}(77)_{\alpha(Rb11)} \cdot 10^{-12}$$


$$a_\mu^{\text{QED}}(a_e) = 1\,165\,847\,188.46(9)_\text{mass}(19)_{\alpha^4}(7)_{\alpha^5}(30)_{\alpha(a_e)} \cdot 10^{-12}$$

[Aoyama et al., Phys. Rev. Lett. 109, 111807 (2012)]

$$a_\mu^{\text{exp}} - a_\mu^{\text{QED}} = 737.0(6.3)\cdot 10^{-10}$$

# Theory II: Weak interactions

- Weak contributions :  $W$ ,  $Z$ ,... loops



$$\begin{aligned}
 a_\mu^{\text{weak(1)}} &= \frac{G_F}{\sqrt{2}} \frac{m_\mu^2}{8\pi^2} \left[ \frac{5}{3} + \frac{1}{3} (1 - 4 \sin^2 \theta_W)^2 + \mathcal{O}\left(\frac{m_\mu^2}{M_Z^2} \log \frac{M_Z^2}{m_\mu^2}\right) + \mathcal{O}\left(\frac{m_\mu^2}{M_H^2} \log \frac{M_H^2}{m_\mu^2}\right) \right] \\
 &= 19.48 \times 10^{-10}
 \end{aligned}$$

[W.A. Bardeen, R. Gastmans and B.E. Lautrup, Nucl. Phys. B46, 315 (1972)]

[G. Altarelli, N. Cabibbo and L. Maiani, Phys. Lett. 40B, 415 (1972)]

[R. Jackiw and S. Weinberg, Phys. Rev. D 5, 2473 (1972)]

[I. Bars and M. Yoshimura, Phys. Rev. D 6, 374 (1972)]

[M. Fujikawa, B.W. Lee and A.I. Sanda, Phys. Rev. D 6, 2923 (1972)]

## Two-loop bosonic contributions

$$a_\mu^{\text{weak(2);b}} = \frac{G_F}{\sqrt{2}} \frac{m_\mu^2}{8\pi^2} \frac{\alpha}{\pi} \cdot \left[ -5.96 \ln \frac{M_W^2}{m_\mu^2} + 0.19 \right] = \frac{G_F}{\sqrt{2}} \frac{m_\mu^2}{8\pi^2} \left( \frac{\alpha}{\pi} \right) \cdot (-79.3)$$

[A. Czarnecki, B. Krause and W. J. Marciano, Phys. Rev. Lett. 76, 3267 (1996)]

## Two-loop fermionic contributions

[A. Czarnecki, W.J. Marciano, A. Vainshtein, Phys. Rev. D 67, 073006 (2003). Err.-ibid. D 73, 119901 (2006)]

[M. K., S. Peris, M. Perrottet, E. de Rafael, JHEP11, 003 (2002)]

$$a_\mu^{\text{weak}} = (154 \pm 1) \cdot 10^{-11}$$

$$a_e^{\text{weak}} = (0.0297 \pm 0.0005) \cdot 10^{-12}$$

Recent update:  $a_\mu^{\text{weak}} = (153.6 \pm 1.0) \cdot 10^{-11}$


[C. Gnendinger et al., Phys. Rev. D 88, 053005 (2013)]

$$a_\mu^{\mathrm{exp}} - a_\mu^{\mathrm{QED}} - a_\mu^{\mathrm{weak}} = 721.65(6.38) \cdot 10^{-10}$$

# Theory III: Strong interactions


- Hadronic contributions : quark and gluon loops

$$a_\ell^{\text{had}} = a_\ell^{\text{HVP-LO}} + a_\ell^{\text{HVP-HO}} + a_\ell^{\text{HLxL}}$$



Lowest-order hadronic vacuum polarization (HVP-LO)

Higher-order hadronic vacuum polarization (HVP-HO)



Hadronic light-by-light scattering (HLxL)

## Hadronic vacuum polarization

- Occurs first at order  $\mathcal{O}(\alpha^2)$

- Can be expressed as

$$a_\ell^{\text{HVP-LO}} = \frac{1}{3} \left( \frac{\alpha}{\pi} \right)^2 \int_{4M_\pi^2}^\infty \frac{dt}{t} K(t) R^{\text{had}}(t) \quad K(t) = \int_0^1 dx \frac{x^2(1-x)}{x^2 + (1-x) \frac{t}{m_\ell^2}}$$

[C. Bouchiat, L. Michel, J. Phys. Radium 22, 121 (1961)]

[L. Durand, Phys. Rev. 128, 441 (1962); Err.-ibid. 129, 2835 (1963)]




[M. Gourdin, E. de Rafael, Nucl. Phys. B 10, 667 (1969)]

- $K(s) > 0$  and  $R^{\text{had}}(s) > 0 \implies a_\ell^{\text{HVP-LO}} > 0$

- $K(s) \sim m_\ell^2/(3s)$  as  $s \rightarrow \infty \implies$  the (non perturbative) low-energy region dominates

## Hadronic vacuum polarization

- Can be evaluated using available experimental data



- Some order  $\mathcal{O}(\alpha^3)$  corrections included

- exchange of virtual photons between final state hadrons
- some radiative exclusive modes, e.g.  $\pi^0\gamma$

$$a_\mu^{\pi^0\gamma}(600 \text{ MeV} - 1030 \text{ MeV}) = 4.4(1.9) \cdot 10^{-10}$$

- The two most recent determinations are in good agreement (being based on the same data sets, this should not be a surprise) and give a relative precision of 0.6%

## Latest (published) results

$$\begin{aligned} a_\mu^{\text{HVP-LO}} &= 692.3 \pm 4.2 \cdot 10^{-10} & [\text{M. Davier et al., Eur. Phys. J. C 71, 1515 (2011)}] \\ a_\mu^{\text{HVP-LO}} &= 694.9 \pm 4.3 \cdot 10^{-10} & [\text{K. Hagiwara et al., J. Phys. G 38, 085003 (2011)}] \\ a_e^{\text{HVP-LO}} &= 1.866(11) \cdot 10^{-12} & [\text{D. Nomura, T. Teubner, Nucl. Phys. B 867, 236 (2013)}] \end{aligned}$$

$$a_\mu^{\text{HVP-NLO}} = \frac{1}{3} \left( \frac{\alpha}{\pi} \right)^3 \int_{4M_\pi^2}^\infty \frac{dt}{t} K^{(2)}(t) R^{\text{had}}(t)$$

[J. Calmet, S. Narison, M. Perrottet, E. de Rafael, Phys. Lett. B 61, 283 (1976)]

[B. Krause, Phys. Lett. B 390, 392 (1997)]

$$\begin{aligned} a_\mu^{\text{HVP-NLO}} &= -9.84 \pm 0.07 \cdot 10^{-10} & [\text{K. Hagiwara et al., J. Phys. G 38, 085003 (2011)}] \\ a_e^{\text{HVP-NLO}} &= -0.2234(14) \cdot 10^{-12} & [\text{D. Nomura, T. Teubner, Nucl. Phys. B 867, 236 (2013)}] \\ a_\mu^{\text{HVP-NNLO}} &= 1.24 \pm 0.01 \cdot 10^{-10} & [\text{A. Kurz et al., Phys. Lett. B 734, 144 (2014)}] \\ a_e^{\text{HVP-NNLO}} &= 0.028(1) \cdot 10^{-12} \end{aligned}$$

## Hadronic vacuum polarization

- Some tension between, for instance, the high-precision data collected in the region of the  $\rho$  resonance by BaBar and KLOE/KLOE-2

Experiment	$a_{\mu}^{\text{HVP-LO } 2\pi} (600 - 900 \text{ MeV})$
BaBar	376.7(2.0)(1.9)
KLOE 08	368.9(0.4)(2.3)(2.2)
KLOE 10	366.1(0.9)(2.3)(2.2)
KLOE 12	366.7(1.2)(2.4)(0.8)

- These tensions need to be resolved in order to achieve higher precision  
→ new data (KLOE-2, BaBar, VEPP-2000, BESIII,...)

Experiment	$a_{\mu}^{\text{HVP-LO } 2\pi} (600 - 900 \text{ MeV})$
BESIII	368.2(2.5)(3.3) [BESIII Coll., Phys. Lett. B 753, 629 (2016)]

## Hadronic vacuum polarization

- Update including new data available since 2011

$$a_\mu^{\text{HVP-LO}} = 686.99 \pm 4.21 \cdot 10^{-10}$$

$$a_\mu^{\text{HVP-NLO}} = -9.934 \pm 0.091 \cdot 10^{-10}$$

$$a_\mu^{\text{HVP-NNLO}} = 1.226 \pm 0.012 \cdot 10^{-10}$$

$$a_e^{\text{HVP-LO}} = 1.8464(121) \cdot 10^{-12}$$

$$a_e^{\text{HVP-NLO}} = -0.2210(14) \cdot 10^{-12}$$

$$a_e^{\text{HVP-NNLO}} = 0.0279(2) \cdot 10^{-12}$$

[F. Jegerlehner, arXiv:1511.04473 [hep-ph]]

- Possibility to extract HVP from Bhabha scattering?

[C. M. Carloni-Calame, M. Passera, L. Trentadue, G. Venanzoni, Phys. Lett. B 476, 325 (2015)]


- Alternative for the (near?) future: Lattice QCD

## Hadronic light-by-light: the really complicated thing

- occurs at order  $\mathcal{O}(\alpha^3)$
- not related, as a whole, to an experimental observable...

?


→



- Involves the fourth-rank vacuum polarization tensor

$$\text{F.T. } \langle 0 | T\{VVVV\} | 0 \rangle \longrightarrow \Pi_{\mu\nu\rho\sigma}(q_1, q_2, q_3, q_4) \quad q_1 + q_2 + q_3 + q_4 = 0$$

- Many identifiable contributions...



## Hadronic light-by-light: the really complicated thing

- Need some organizing principle: ChPT, large- $N_c$  (turns out to be most relevant in practice) [E. de Rafael, Phys. Lett. B 322, 239 (1994)]

$$a_\mu^{\text{HLxL}} = N_c \left(\frac{\alpha}{\pi}\right)^3 \frac{N_c}{F_\pi^2} \frac{m_\mu^2}{48\pi^2} \left[ \ln^2 \frac{M_\rho}{M_\pi} + c_\chi \ln \frac{M_\rho}{M_\pi} + \kappa(M_\rho) \right] + \mathcal{O}(N_c^0)$$

[M. Knecht, A. Nyffeler, Phys. Rev. D 65, 073034 (2002)]

[M. Knecht, A. Nyffeler, M. Perrottet, E. de Rafael, Phys. Rev. Lett. 88, 071802 (2002)]

M. J. Ramsey-Musolf, M. B. Wise, Phys. Rev. Lett. 89, 041601 (2002)]

[J. Prades, E. de Rafael, A. Vainshtein, Glasgow White Paper (2008)]

- Impose QCD short-distance properties [K. Melnikov, A. Vainshtein, Phys. Rev. D, 113006 (2004)]

- Only two (so far) attempts at a “complete”, but model-dependent calculation...

$$a_\mu^{\text{HLxL}} = +(8.3 \pm 3.2) \cdot 10^{-10}$$

[J. Bijnens, E. Pallante, J. Prades, Phys. Rev. Lett. 75, 1447 (1995) [Err.-ibid. 75, 3781 (1995)]; Nucl. Phys. B 474, 379 (1995); Nucl. Phys. B 626, 410 (2002)]

$$a_\mu^{\text{HLxL}} = +(89.6 \pm 15.4) \cdot 10^{-11}$$

[M. Hayakawa, T. Kinoshita, A. I. Sanda, Phys. Rev. Lett. 75, 790 (1995); Phys. Rev. D 54, 3137 (1996)]

[M. Hayakawa, T. Kinoshita, Phys. Rev. D 57, 365 (1998) [Err.-ibid. 66, 019902(E) (2002)]]

...after the sign change [M.K. and A. Nyffeler, Phys. Rev. D 65, 073034 (2002)]

units:  $10^{-11}$

Contribution	BPP	HKS	KN	MV	BP	PdRV	N/JN
$\pi^0, \eta, \eta'$	$85 \pm 13$	$82.7 \pm 6.4$	$83 \pm 12$	$114 \pm 10$	—	$114 \pm 13$	$99 \pm 16$
$\pi, K$ loops	$-19 \pm 13$	$-4.5 \pm 8.1$	—	—	—	$-19 \pm 19$	$-19 \pm 13$
$\pi, K$ l. + subl. in Nc	—	—	—	$0 \pm 10$	—	—	—
axial vectors	$2.5 \pm 1.0$	$1.7 \pm 1.7$	—	$22 \pm 5$	—	$15 \pm 10$	$22 \pm 5$
scalars	$-6.8 \pm 2.0$	—	—	—	—	$-7 \pm 7$	$-7 \pm 2$
quark loops	$21 \pm 3$	$9.7 \pm 11.1$	—	—	—	2.3	$21 \pm 3$
total	$83 \pm 32$	$89.6 \pm 15.4$	$80 \pm 40$	$136 \pm 25$	$110 \pm 40$	$105 \pm 26$	$116 \pm 39$

BPP: J. Bijnens, E. Pallante, J. Prades, Phys. Rev. Lett. 75 (1995) 1447 [Erratum-ibid. 75 (1995) 3781]; Nucl. Phys. B 474 (1996) 379; [Erratum-ibid. 626 (2002) 410]

HKS: M. Hayakawa, T. Kinoshita, A. I. Sanda, Phys. Rev. Lett. 75 (1995) 790; Phys. Rev. D 54 (1996) 3137

KN: M. Nech, A. Nyffeler, Phys. Rev. D 65 (2002) 073034

MV: K. Melnikov, A. Vainshtein, Phys. Rev. D 70 (2004) 113006

BP: J. Bijnens, J. Prades, Acta Phys. Polon. B 38 (2007) 2819; J. Prades, Nucl. Phys. Proc. Suppl. 181-182 (2008) 15;  
J. Bijnens, J. Prades, Mod. Phys. Lett. A 22 (2007) 767

BdRV: J. Prades, E. de Rafael, A. Vainshtein, arXiv:0901.0306 [hep-ph]

N/NJ: A. Nyffeler, Phys. Rev. D 79, 073012 (2009); F. Jegerlehner, A. Nyffeler, Phys. Rep. (2009)

units:  $10^{-11}$

Contribution	BPP	HKS	KN	MV	BP	PdRV	N/JN
$\pi^0, \eta, \eta'$	$85 \pm 13$	$82.7 \pm 6.4$	$83 \pm 12$	$114 \pm 10$	—	$114 \pm 13$	$99 \pm 16$
$\pi, K$ loops	$-19 \pm 13$	$-4.5 \pm 8.1$	—	—	—	$-19 \pm 19$	$-19 \pm 13$
$\pi, K$ l. + subl. in Nc	—	—	—	$0 \pm 10$	—	—	—
axial vectors	$2.5 \pm 1.0$	$1.7 \pm 1.7$	—	$22 \pm 5$	—	$15 \pm 10$	$22 \pm 5$
scalars	$-6.8 \pm 2.0$	—	—	—	—	$-7 \pm 7$	$-7 \pm 2$
quark loops	$21 \pm 3$	$9.7 \pm 11.1$	—	—	—	2.3	$21 \pm 3$
total	$83 \pm 32$	$89.6 \pm 15.4$	$80 \pm 40$	$136 \pm 25$	$110 \pm 40$	$105 \pm 26$	$116 \pm 39$

BPP: J. Bijnens, E. Pallante, J. Prades, Phys. Rev. Lett. 75 (1995) 1447 [Erratum-ibid. 75 (1995) 3781]; Nucl. Phys. B 474 (1996) 379; [Erratum-ibid. 626 (2002) 410]

HKS: M. Hayakawa, T. Kinoshita, A. I. Sanda, Phys. Rev. Lett. 75 (1995) 790; Phys. Rev. D 54 (1996) 3137

KN: M. Nech, A. Nyffeler, Phys. Rev. D 65 (2002) 073034

MV: K. Melnikov, A. Vainshtein, Phys. Rev. D 70 (2004) 113006

BP: J. Bijnens, J. Prades, Acta Phys. Polon. B 38 (2007) 2819; J. Prades, Nucl. Phys. Proc. Suppl. 181-182 (2008) 15;  
J. Bijnens, J. Prades, Mod. Phys. Lett. A 22 (2007) 767

BdRV: J. Prades, E. de Rafael, A. Vainshtein, arXiv:0901.0306 [hep-ph]

N/NJ: A. Nyffeler, Phys. Rev. D 79, 073012 (2009); F. Jegerlehner, A. Nyffeler, Phys. Rep. (2009)

Recent reevaluation of single meson exchanges

$$a_\mu(f_1, f'_1) = 6.4(2.0) \cdot 10^{-11} \quad a_\mu(f_0, f'_0, a_0) = (-1 \text{ to } -4) \cdot 10^{-11} \quad a_\mu(f_2, f'_2, a_2, a'_2) = 1.1(0.1) \cdot 10^{-11}$$

[V. Pauk, M. Vanderhaeghen, arXiv:1401.0832 [hep-ph]]

## Hadronic light-by-light: the really complicated thing

### Recent (partial) reevaluations

$$a_\mu^{\text{HLxL}} = (10.5 \pm 2.6) \cdot 10^{-10} \quad [\text{J. Prades, E. de Rafael, A. Vainshtein, arXiv:0901.0306}]$$

“best estimate”

$$a_\mu^{\text{HLxL}} = (11.5 \pm 4.0) \cdot 10^{-10} \quad [\text{A. Nyffeler, Phys. Rev. D 79, 073012 (2009)}]$$


more conservative estimate

$$a_e^{\text{HLxL}} = (0.035 \pm 0.010) \cdot 10^{-12} \quad [\text{J. Prades, E. de Rafael, A. Vainshtein, in } \textit{Lepton Dipole Moments}]$$

## Hadronic light-by-light: the really complicated thing

- More recently: dispersive approaches

- for  $\Pi_{\mu\nu\rho\sigma}$



$$\Pi = \Pi^{\pi^0, \eta, \eta' \text{ poles}} + \Pi^{\pi^\pm, K^\pm \text{ loops}} + \Pi^{\pi\pi} + \Pi^{\text{residual}}$$

[G. Colangelo, M. Hoferichter, M. Procura, P. Stoffer, JHEP09, 091 (2014); arXiv:1506.01386 [hep-ph]]

Needs input from data (transition form factors,...)

[G. Colangelo, M. Hoferichter, B. Kubis, M. Procura, P. Stoffer, Phys. Lett. B 738, 6 (2014)]

[A. Nyffeler, arXiv:1602.03398 [hep-ph]]

Main unanswered issues:

- how will short-distance constraints be imposed?
  - how will  $\Pi^{\text{residual}}$  be estimated? Cf. axial vectors (leading in large- $N_c$ )  $\rightarrow 3\pi$  channel
- 
- for  $F_2^{\text{HLxL}}(k^2)$

only pion pole with VMD form factor (two-loop graph) reconstructed this way so far

[V. Pauk and M. Vanderhaeghen, Phys. Rev. D 90, 113012 (2014) [arXiv:1409.0819 [hep-ph]]]

Try to get further insight or to test new ideas by studying simple models

cf. D. Greynat, E. de Rafael, JHEP1207, 020 (2012) based on a version of the chiral constituent quark model that is renormalizable in the large- $N_C$  limit

[S. Weinberg, Phys. Rev. Lett. 105, 261601 (2010)]

[E. de Rafael, Phys. Lett. B 703, 60 (2011)]

# Conclusions

# Perspectives for the future

- The anomalous magnetic moments of the electron and of the muon are among the most precisely measured observables of the standard model

$$a_e^{\text{exp}} = 1\,159\,652\,180.73(0.28) \cdot 10^{-12} \quad [0.24\text{ppb}]$$

$$a_\mu^{\text{exp}} = 116\,592\,089(63) \cdot 10^{-11} \quad [0.54\text{ppm}]$$

- The anomalous magnetic moments of the electron and of the muon are among the most precisely measured observables of the standard model

$$a_e^{\text{exp}} = 1\,159\,652\,180.73(0.28) \cdot 10^{-12} \quad [0.24\text{ppb}]$$

$$a_\mu^{\text{exp}} = 116\,592\,089(63) \cdot 10^{-11} \quad [0.54\text{ppm}]$$

- In the case of the electron, there is good agreement between theory and experiment:

$$a_e^{\text{exp}} - a_e^{\text{SM}} = (-0.91 \pm 0.82) \cdot 10^{-12}$$

- The anomalous magnetic moments of the electron and of the muon are among the most precisely measured observables of the standard model

$$a_e^{\text{exp}} = 1\,159\,652\,180.73(0.28) \cdot 10^{-12} \quad [0.24\text{ppb}]$$

$$a_\mu^{\text{exp}} = 116\,592\,089(63) \cdot 10^{-11} \quad [0.54\text{ppm}]$$

- At present, the standard model value for  $a_\mu$  misses the experimental determination by about 3.5 (or even 4) standard deviations

- The anomalous magnetic moments of the electron and of the muon are among the most precisely measured observables of the standard model

$$\begin{aligned} a_e^{\text{exp}} &= 1\,159\,652\,180.73(0.28) \cdot 10^{-12} \quad [0.24\text{ppb}] \\ a_\mu^{\text{exp}} &= 116\,592\,089(63) \cdot 10^{-11} \quad [0.54\text{ppm}] \end{aligned}$$

- At present, the standard model value for  $a_\mu$  misses the experimental determination by about 3.5 (or even 4) standard deviations
- It is not obvious to find a straightforward explanation for this persistent discrepancy:

$$a_\mu^{\text{exp}} - a_\mu^{\text{SM}} \sim 30 \cdot 10^{-10} \sim 2 \cdot a_\mu^{\text{weak}} \sim a_\mu^{\text{QED}}(\alpha^4)$$

$$a_\mu^{\text{HVP-LO}} = (692.3 \pm 4.2) \cdot 10^{-10} \quad a_\mu^{\text{HVP-HO}} = (-8.60 \pm 0.07) \cdot 10^{-10}$$

$$a_\mu^{\text{HLxL}} = (10.5 \pm 2.6) \cdot 10^{-10}$$

$$\Delta a_\mu^{\text{exp}} = 6.3 \cdot 10^{-10}$$

- The anomalous magnetic moments of the electron and of the muon are among the most precisely measured observables of the standard model

$$\begin{aligned} a_e^{\text{exp}} &= 1159\,652\,180.73(0.28) \cdot 10^{-12} \quad [0.24\text{ppb}] \\ a_\mu^{\text{exp}} &= 116\,592\,089(63) \cdot 10^{-11} \quad [0.54\text{ppm}] \end{aligned}$$

- At present, the standard model value for  $a_\mu$  misses the experimental determination by about 3.5 (or even 4) standard deviations

- It is not obvious to find a straightforward explanation for this persistent discrepancy:

$$a_\mu^{\text{exp}} - a_\mu^{\text{SM}} \sim 30 \cdot 10^{-10} \sim 2 \cdot a_\mu^{\text{weak}} \sim a_\mu^{\text{QED}}(\alpha^4)$$

Higher order QED effects?

$$A_2^{(12)}(m_\mu/m_e) \sim A_2^{(6)}(m_\mu/m_e; \text{LxL}) \left[ \frac{2}{3} \ln \frac{m_\mu}{m_e} - \frac{5}{9} \right]^3 \cdot 10 \sim 0.6 \cdot 10^4 \longrightarrow \delta a_\mu^{\text{QED}} \sim 1 \cdot 10^{-12}$$

Higher order QCD effects?

$$a_\mu^{\text{HVP-NNLO}} = (1.24 \pm 0.01) \cdot 10^{-10} \qquad a_\mu^{\text{HLxL-HO}} \sim (0.3 \pm 0.2) \cdot 10^{-10}$$

[A. Kurz et al., Phys. Lett. B 734, 144 (2014)]

[G. Colangelo et al., Phys. Lett. B 735, 90 (2014)]

- The anomalous magnetic moments of the electron and of the muon are among the most precisely measured observables of the standard model

$$a_e^{\text{exp}} = 1\,159\,652\,180.73(0.28) \cdot 10^{-12} \quad [0.24\text{ppb}]$$

$$a_\mu^{\text{exp}} = 116\,592\,089(63) \cdot 10^{-11} \quad [0.54\text{ppm}]$$

- At present, the standard model value for  $a_\mu$  misses the experimental determination by about 3.5 (or even 4) standard deviations

- It is not obvious to find a straightforward explanation for this persistent discrepancy:

$$a_\mu^{\text{exp}} - a_\mu^{\text{SM}} \sim 30 \cdot 10^{-10} \sim 2 \cdot a_\mu^{\text{weak}} \sim a_\mu^{\text{QED}}(\alpha^4)$$

Manifestation of BSM degrees of freedom?

→ many proposals, situation remains inconclusive

- The anomalous magnetic moments of the electron and of the muon are among the most precisely measured observables of the standard model

$$a_e^{\text{exp}} = 1159\,652\,180.73(0.28) \cdot 10^{-12} \quad [0.24\text{ppb}]$$


$$a_\mu^{\text{exp}} = 116\,592\,089(63) \cdot 10^{-11} \quad [0.54\text{ppm}]$$

- At present, the standard model value for  $a_\mu$  misses the experimental determination by about 3.5 (or even 4) standard deviations
- Emergence of a pattern?

Observable	Tension wrt SM	Limited by
$B \rightarrow D^{(*)}\tau\nu/B \rightarrow D^{(*)}\ell\nu, \ell = e, \mu$	$3.9\sigma$	experiment
$(g-2)_\mu$	$\gtrsim 3.5\sigma$	exp. and th.
$B^0 \rightarrow K^{*0}\mu\mu$ ang. dist., BR	$3.4\sigma$	exp. and th.
$B_s \rightarrow \phi\mu\mu$ BR	$3.0\sigma$	experiment
Dimuon CP asymmetry	$3.0\sigma$	experiment
$V_{ub}$ exclusive vs. inclusive	$3.0\sigma$	exp. and th.
$B^+ \rightarrow K^+\mu\mu/B^+ \rightarrow K^+ee$	$2.6\sigma$	experiment
$h \rightarrow \tau\mu$	$2.4\sigma$	experiment

[cf. O. Leroy, talk at Lake Louise Winter Institute, Feb. 2016]

# Can lattice QCD contribute?



Strategies are being devised

- twisted boundary conditions

[C. Aubin, T. Blum, M. Golterman and S. Peris, arXiv:1307.4701 [hep-lat]]

- time moments and Padé approximants

[B. Chakraborty et al, arXiv:1403.1778 [hep-lat]]

[ M. Golterman, K. Maltman and S. Peris, arXiv:1405.2389 [hep-lat]]

- Moments of the MB transform

[E. de Rafael, Phys. Lett. B 736, 522 (2014)]



First results also for HLxL...

[T. Blum, S. Chowdhury, M. Hayakawa, T. Izubuchi, Phys. Rev. Lett. 114, 012001 (2015)]

- Two new experiments, with the aim of reducing the experimental uncertainty by a factor of 4, are being prepared, at FNAL and at J-PARC

# Present experimental status and the new project at FNAL

David Hertzog  
*University of Illinois at Urbana-Champaign*



Muon g-2 and EDM in the LHC era - Feb. 2010

D. Hertzog, LPNHE Workshop Paris, Feb. 2010

# Magic vs “New Magic”


## ■Complimentary!

$$\vec{\omega} = -\frac{e}{m} \left[ a_\mu \vec{B} - \left( a_\mu - \frac{1}{\gamma^2 - 1} \right) \frac{\vec{\beta} \times \vec{E}}{c} + \frac{\eta}{2} \left( \vec{\beta} \times \vec{B} + \frac{\vec{E}}{c} \right) \right]$$

BNL/Fermilab Approach

$$a_\mu - \frac{1}{\gamma^2 - 1} = 0$$

$$\eta \approx 0$$



$$\gamma_{\text{magic}} = 29.3$$

$$p_{\text{magic}} = 3.09 \text{ GeV}/c$$


J-PARC Approach

$$\vec{E} = 0$$


$$\vec{\omega} = \vec{\omega}_a + \vec{\omega}_\eta$$



$$\vec{\omega}_a = -\frac{e}{m} a_\mu \vec{B}$$



# Muon storage magnet and detector



T. Mibe, LPNHE Workshop Paris, Dec. 2014

- Two new experiments, with the aim of reducing the experimental uncertainty by a factor of 4, are being prepared, at FNAL and at J-PARC (first results expected in  $\sim 2$  years)

- New high-precision data from VEPP-2000, BESIII,...

- $a_e$  is expected to be about  $(m_\mu/m_e)^2 \sim 40000$  times less sensitive to BSM effects than  $a_\mu$ . But it is known with much better ( $\sim 2300$ ) precision... Possibilities to observe BSM effects through  $a_e$

[G. F. Giudice, P. Paradisi, M. Passera, JHEP 1211, 113 (2012)]

- Needs improvements on the determinations of  $a_e$  [from 0.24ppb to 0.06ppb], of  $R_\infty$ ,  $m_e/m_u$ ,...

$$\alpha^2 = \frac{2R_\infty}{c} \frac{M_{\text{at}}}{m_u} \frac{m_u}{m_e} \frac{h}{M_{\text{at}}}$$



that are within reach on a timescale similar to the one of the new  $(g - 2)_\mu$  experiments.

[F. Terranova, G. M. Tino, Phys. Rev. A 89, 052118 (2014)]

Thanks for your attention!

# Present experimental status and the new project at FNAL

David Hertzog  
*University of Illinois at Urbana-Champaign*



Muon g-2 and EDM in the LHC era - Feb. 2010

D. Hertzog, LPNHE Workshop Paris, Feb. 2010





5 nights sheltering from  
Storm near Norfolk  
Virginia.


Tennessee-Tombigbee Waterway  
Mississippi, Illinois and  
Des Plaines rivers.

25th June - July 20th

S. Maxfield, PhiPsi 2013



S. Maxfield, PhiPsi 2013







S. Maxfield, PhiPsi 2013