

Hard processes in pA collisions

François Arleo

LLR Palaiseau

Théorie LHC France & GDR PH-QCD

Orsay – November 2016

Why hard processes in pA collisions?

Hard processes

- Great variety
 - ▶ W/Z, Drell-Yan, photons, (b-quark) jets, light/heavy hadrons. . .
- Well known in QCD
 - ▶ computed in perturbation theory and systematically compared to pp
 - ▶ caveat: hadron production (especially quarkonia) less understood

Why hard processes in pA collisions?

Hard processes

- Great variety
 - ▶ W/Z, Drell-Yan, photons, (b-quark) jets, light/heavy hadrons. . .
- Well known in QCD
 - ▶ computed in perturbation theory and systematically compared to pp
 - ▶ caveat: hadron production (especially quarkonia) less understood

pA collisions

- 'Simple' medium: static, known density profile
- Easier measurements (than in AA) due to smaller underlying event
- Small influence of the produced medium on hard processes
 - ▶ less true at LHC, less true for (excited) quarkonia

Why hard processes in pA collisions?

Hard processes

- Great variety
 - ▶ W/Z, Drell-Yan, photons, (b-quark) jets, light/heavy hadrons. . .
- Well known in QCD
 - ▶ computed in perturbation theory and systematically compared to pp
 - ▶ caveat: hadron production (especially quarkonia) less understood

pA collisions

- 'Simple' medium: static, known density profile
- Easier measurements (than in AA) due to smaller underlying event
- Small influence of the produced medium on hard processes
 - ▶ less true at LHC, less true for (excited) quarkonia

In short

Closest to (theory/exp) **QCD studies** in pp, yet with a **heavy-ion touch**

From pp to the heavy-ion touch

- pp/pA collisions in collinear factorization
 - ▶ leading twist nPDF analyses
 - ▶ observables: W/Z, Drell-Yan, jets
- Beyond collinear factorization: multiple scattering in nuclei
 - ▶ Momentum broadening and induced gluon radiation
 - ▶ observables: light and heavy-quark hadrons


Collinear factorization in pp collisions

Take a **large momentum transfer process** in pp, scale $Q(= p_{\perp}, M) \gg \Lambda_{\text{QCD}}$

$$pp \rightarrow (h, \gamma, Z, \dots) + X$$

Factorization = approximation

$$\frac{d\sigma_{pp}}{dy dQ} = \sum_{i,j} \int dx_1 f_i^p(x_1, \mu) \int dx_2 f_j^p(x_2, \mu) \frac{d\hat{\sigma}_{ij}(Q, \mu')}{dy dQ} + \mathcal{O}\left(\frac{\Lambda_p^n}{Q^n}\right)$$



Collinear factorization in pp collisions

Take a **large momentum transfer process** in pp, scale $Q(= p_{\perp}, M) \gg \Lambda_{\text{QCD}}$

$$pp \rightarrow (h, \gamma, Z, \dots) + X$$

Factorization = approximation

$$\frac{d\sigma_{pp}}{dy dQ} = \sum_{i,j} \int dx_1 f_i^p(x_1, \mu) \int dx_2 f_j^p(x_2, \mu) \frac{d\hat{\sigma}_{ij}(Q, \mu')}{dy dQ} + \mathcal{O}\left(\frac{\Lambda_p^n}{Q^n}\right)$$

- **Predictive power**
 - ▶ long distance physics encoded into PDF (and FF) which are **universal**
 - ▶ short distance calculable in perturbation theory
- **Power corrections** due to long range soft gluon interaction
 - ▶ process dependent, **not universal**

What about pA collisions ?

Collinear factorization in pA collisions

A nucleus as an ordinary hadron

$$\frac{d\sigma_{pA}}{dy dQ} = \sum_{i,j} \int dx_1 f_i^P(x_1, \mu) \int dx_2 f_j^A(x_2, \mu) \frac{d\hat{\sigma}_{ij}(Q, \mu')}{dy dQ} + \mathcal{O}\left(\frac{\Lambda_A^n}{Q^n}\right)$$

- **Universal** (leading twist) **nuclear PDF**
 - ▶ could be probed in various processes and collision systems (eA, γ A, pA)
- New scale for power corrections ($\Lambda_A > \Lambda_p$)
 - ▶ **higher twist processes enhanced in pA** collisions (wrt pp)
 - ▶ specific processes could spoil the extraction of (universal) nPDF

Collinear factorization in pA collisions

A nucleus as an ordinary hadron

$$\frac{d\sigma_{pA}}{dy dQ} = \sum_{i,j} \int dx_1 f_i^P(x_1, \mu) \int dx_2 f_j^A(x_2, \mu) \frac{d\hat{\sigma}_{ij}(Q, \mu')}{dy dQ} + \mathcal{O}\left(\frac{\Lambda_A^n}{Q^n}\right)$$

- **Universal** (leading twist) **nuclear PDF**
 - ▶ could be probed in various processes and collision systems (eA, γ A, pA)
- New scale for power corrections ($\Lambda_A > \Lambda_p$)
 - ▶ **higher twist processes enhanced in pA** collisions (wrt pp)
 - ▶ specific processes could spoil the extraction of (universal) nPDF

What to expect for f_i^A ? How does it compare to f_i^P ?

PDF of a nucleus

In a super dilute nucleus: f^A given by **incoherent sum** over nucleon PDF

$$f_i^A = Z f_i^p + (A - Z) f_i^n$$

$$d\sigma_{pA} = Z d\sigma_{pp} + (A - Z) d\sigma_{pn} \simeq A d\sigma_{pp} \Rightarrow R_{pA} \equiv \frac{1}{A} \frac{d\sigma^{pA}}{d\sigma^{pp}} \simeq 1$$

PDF of a nucleus

In a super dilute nucleus: f^A given by **incoherent sum** over nucleon PDF

$$f_i^A = Z f_i^p + (A - Z) f_i^n$$

In practice, distance between nucleons **much smaller than coherence time** at high energy

$$\ell_c \sim \frac{E}{Q^2} \sim \frac{1}{2m_N x_2} \gg 1 \text{ fm}$$

- Onset of nuclear shadowing at small $x_2 \lesssim 10^{-1}$
- Working assumption

$$f_i^A = Z R_i^{p/A} f_i^p + (A - Z) R_i^{n/A} f_i^n$$

- ▶ nPDF ratios $R_i^{p/A}(x, Q^2)$ assumed to be universal
- ▶ extracted from (N)LO global fit analysis based on DGLAP evolution
[EKS98, nDS, HKN... EPS09, DSSZ, nCTEQ15, KA15]


- Poor constraints from data, especially at small- x and in the gluon channel
- Strong sensitivity on the parametrization at low scale

[Helenius Paukkunen Armesto, 1606.09003]

- Crucial need to use LHC pPb data to reweight nPDF


[Paukkunen Zurita, 1402.6623]

[nCTEQ15, 1509.00792]



- Poor constraints from data, especially at small- x and in the gluon channel
- Strong sensitivity on the parametrization at low scale
[Helenius Paukkunen Armesto, 1606.09003]
- Crucial need to use LHC pPb data to reweight nPDF
[Paukkunen Zurita, 1402.6623]

What would be the best processes
to reach that goal?



Probing leading twist nPDF at the LHC

Ideally, looking for processes sensitive to PDF **only**

Some requirements (not necessary, but preferable):


- Sufficiently large scale: $Q \gg Q_s \simeq \text{few GeV}$
 - ▶ avoid non-linear evolution and large power corrections
- ... but not too large to keep some sensitivity
 - ▶ $f^A/Af^P \simeq 1$ in the 'Bjorken limit' ($Q^2 \rightarrow \infty$ at fixed x)
- Integrated over all p_\perp (or focus on $p_\perp \gg Q_s$)
 - ▶ avoid multiple scattering effects e.g. Cronin effect
- Favor color-neutral probes
 - ▶ avoid coherent energy loss

Best candidates

Weak bosons, Jets, Drell-Yan

W/Z measured in pPb (and PbPb) by ALICE, ATLAS & CMS



[motivated in Paukkunen Salgado, [1010.5392](#)]




- R_{pA} of Z boson measured by ATLAS
 - ▶ Slight suppression at forward rapidity (smaller x in Pb)

W/Z measured in pPb (and PbPb) by ALICE, ATLAS & CMS

[motivated in Paukkunen Salgado, [1010.5392](https://arxiv.org/abs/1010.5392)]



- W boson rapidity asymmetry measured by CMS
 - ▶ data favor CT10×EPS09




Lepton charge asymmetry

$$\frac{N_l^+ - N_l^-}{N_l^+ + N_l^-}$$


- Tension at negative $\eta \rightarrow$ possible flavour dependence $R_u^A \neq R_d^A$
 - ▶ Isospin symmetry $R_u^A = R_d^A$ often assumed due to lack of data
 - ▶ pPb Run 2 should tell
- Simple scaling relates pPb and PbPb cross sections at $y < 0$

[FA Chapon Paukkunen, [1509.03993](https://arxiv.org/abs/1509.03993)]




- Jets in pPb at LHC sensitive to (gluon & valence quark) nPDF, in the vicinity of the anti-shadowing region

[Paukkunen Eskola Salgado, [1408.4563](#)]





- Tight constraints brought by CMS dijets
- nPDF effects favored... yet no single set reproduces whole dataset
 - ▶ crucial data to further constrain x & Q^2 dependence of nPDF
 - ▶ other effects at the lowest Q^2 ?

A golden probe of sea quark (and gluon) shadowing



- Low scale $Q \sim 10$ GeV can be reached
 - ▶ better than weak bosons, jets, prompt photons
 - ▶ mass can be varied
- Colorless final state at LO
 - ▶ small/negligible energy loss in cold matter
- Very well understood in QCD
 - ▶ better than light or heavy hadrons



[FA Peigné, [1512.01794](#)]


- NLO calculations using DSSZ, EPS09 and nCTEQ15
 - ▶ should reveal sea quark shadowing at low scale
 - ▶ could be computed in the saturation formalism too
- To be measured by LHCb at fwd/bwd rapidity in pPb Run 2 (now!)
 - ▶ reasonable statistics expected

Beyond leading-twist nPDF

What is **not** included in leading-twist nPDF ?

- Non-linear QCD evolution
 - ▶ all nPDF global fits based on (linear) DGLAP evolution
- Momentum broadening of the fast parton in the nucleus

... taken into account in the **saturation formalism** (CGC)



What is **not** included in leading-twist nPDF ?

- Non-linear QCD evolution
 - ▶ all nPDF global fits based on (linear) DGLAP evolution
- Momentum broadening of the fast parton in the nucleus

... taken into account in the **saturation formalism** (CGC)


- Based on JIMWLK or BK (non-linear) evolution
- Rescattering effects resummed in the dipole formalism
- Based on k_{\perp} factorization
 - ▶ working assumption since k_{\perp} factorization not proven in QCD
- Several (semi)hard processes investigated
 - ▶ light hadrons, open heavy-flavour hadrons, quarkonia, photons...

Often used is 'dilute-dense' (hybrid) formalism to model pA collisions

$$\frac{d\sigma_{pp}}{dy d\rho_{\perp}} = \sum_{i=q,g} \int \frac{dz}{z^2} dx_1 f_i^P(x_1, \mu) \mathcal{F}_{x_g}^{F,A}(k_{\perp}) D_{h/i}(z, \mu') + \mathcal{O}(\alpha_s^2)$$



- f_i^P and $D_{h/i}$: ordinary (collinear) PDF and FF – obey DGLAP
- $\mathcal{F}_{x_g}^{F,A}$: unintegrated gluon PDF – obey BFKL, BK, JIMWLK
- In practice, many different implementations and working assumptions
- Important recent development on computing NLO cross section

Light hadrons



- Mid-rapidity light hadrons show almost **no suppression at $p_{\perp} \gtrsim 2$ GeV** but **depletion** below
 - ▶ possible shadowing/saturation at the smallest p_{\perp} values
 - ▶ depletion due to p_{\perp} broadening
- Agrees with some CGC calculations, yet large exp/th uncertainties
 - ▶ **tendency for less suppression** than in theory – also true with EPS09


Heavy hadrons – D meson



- Little or no suppression at mid-rapidity [ALICE 1605.07569]
- forward/backward rapidity asymmetry reported by LHCb [LHCb-CONF-2016-003]

Energy loss-es


On top of momentum broadening, parton multiple scattering in nuclei induces gluon radiation \rightarrow **energy loss in cold nuclear matter**



- not clear if/how taken into account in CGC formalism

Initial/final state energy loss

LPM regime, small formation time $t_f \lesssim L$



$$\Delta E_{\text{LPM}} \propto \alpha_s \hat{q} L^2 \log(E)$$


- Energy dependence at most logarithmic
- Best probed in
 - ▶ Hadron production in nuclear semi-inclusive DIS
 - ▶ Drell-Yan in pA collisions at low energy
 - ▶ Jet in QGP
- Should be negligible in pA at the LHC
 - ▶ fractional energy loss $\Delta E_{\text{LPM}}/E \ll 1$
 - ▶ explains why weak bosons and jets almost unmodified in pPb

Fully coherent energy loss

Interference between initial and final state, large formation time $t_f \gg L$

[FA Peigné Sami 1006.0818]

$$\Delta E_{\text{coh}} \propto \alpha_s \frac{\sqrt{\hat{q}} L}{M_{\perp}} E \quad (\gg \Delta E_{\text{LPM}})$$



Fully coherent energy loss



Interference between initial and final state, large formation time $t_f \gg L$

[FA Peigné Sami 1006.0818]


$$\Delta E_{\text{coh}} \propto \alpha_s \frac{\sqrt{\hat{q} L}}{M_{\perp}} E \quad (\gg \Delta E_{\text{LPM}})$$

- Predicted from first principles
 - ▶ Same spectrum obtained in the opacity expansion and in dipole model
- [FA Kolevatov Peigné, 1402.1671, Peigné Kolevatov 1405.4241]
[Liou Mueller 1402.1647, Munier Peigné Petreska 1603.01028]
- Important at all energies, especially at large rapidity
 - Needs color in both initial & final state
 - ▶ no effect on W/Z nor Drell-Yan, no effect in DIS
 - Hadron production in pA collisions
 - ▶ applied to quarkonia, other processes currently investigated
 - Power suppressed: negligible when $M_{\perp} \gg \sqrt{\hat{q} L} \sim Q_s$
 - ▶ no jet suppression in pA

Simple fully coherent energy loss model able to **solve the longstanding issue** of J/ψ forward suppression pA data [FA Peigné, 1212.0434]



- Good agreement with all (E866, PHENIX...) quarkonium pA data
 - ▶ Wide range in \sqrt{s} and rapidity
- no nPDF calculation can explain these data



- Predictions in excellent agreement with ALICE (and LHCb) data
 - ▶ especially the trend at large y
 - ▶ good agreement also for Υ

Summary

- Hard processes in pA reveal **many facets of QCD processes**
 - ▶ shadowing/saturation, momentum broadening, radiative energy loss. . .
- Impressive data collected at LHC and earlier. And more to come !
 - ▶ . . . pPb Run 2 starting these days and for a month
- A challenge for theorists: **clarify the role of each process** on various observables and at different energies
 - ▶ still a long way to go. . . but very encouraging progress already made
- pA is exciting **in itself**. Not just 'useful for heavy-ion collisions'.

Summary

- Hard processes in pA reveal **many facets of QCD processes**
 - ▶ shadowing/saturation, momentum broadening, radiative energy loss. . .
- Impressive data collected at LHC and earlier. And more to come !
 - ▶ . . . pPb Run 2 starting these days and for a month
- A challenge for theorists: **clarify the role of each process** on various observables and at different energies
 - ▶ still a long way to go. . . but very encouraging progress already made
- pA is exciting **in itself**. Not just 'useful for heavy-ion collisions'.

Thanks!