



CPPM Seminar - Nov. 28th 2016

Claude Vallée (CPPM/DESY)

Status and Prospects of PHYSICS BEYOND COLLIDERS at CERN

Study Group mandated by the CERN Management
to prepare the next European HEP strategy update (2019-20)
(coordination: J. Jäckel, M. Lamont, C.V.)

Excerpt from the mandate:

“Explore the opportunities offered by the CERN accelerator complex to address some of today’s outstanding questions in particle physics through experiments complementary to high-energy colliders and other initiatives in the world.”

Time scale: next 2 decades

KICK-OFF WORKSHOP

held at CERN on Sept. 6-7th

<https://indico.cern.ch/event/523655/>

> 300 registered participants, 3/4 from outside CERN

AGENDA :

1. Theorists wishes


2. Accelerator complex opportunities

3. Potential future of existing programs

Talks on invitation

4. New ideas: Call for abstracts → 33 abstracts submitted,
20 selected for presentations


THE PRESENT CERN ACCELERATOR COMPLEX




PHYSICS BEYOND COLLIDERS...

... builds on a past decade of lively “diversity” physics !

(currently ~1000 physicists on ~20 experiments)



Recent stop of major programs (e.g. CNGS) leaves room to new significant initiatives




CNGS

*CERN ν_μ beam to Gran Sasso (CNGS)
optimized for ν_τ appearance ($E_\nu \sim 17$ GeV)*









*Successful completion of
the beam operation in 2012 !*



OPERA

establishment of
 ν_τ appearance in ν_μ oscillations





Channel	Expected background				Expected signal	Observed
	Charm	Had. re-interac.	Large μ -scat.	Total		
$\tau \rightarrow 1h$	0.017 ± 0.003	0.022 ± 0.006	—	0.04 ± 0.01	0.52 ± 0.10	3
$\tau \rightarrow 3h$	0.17 ± 0.03	0.003 ± 0.001	—	0.17 ± 0.03	0.73 ± 0.14	1
$\tau \rightarrow \mu$	0.004 ± 0.001	—	0.0002 ± 0.0001	0.004 ± 0.001	0.61 ± 0.12	1
$\tau \rightarrow e$	0.03 ± 0.01	—	—	0.03 ± 0.01	0.78 ± 0.16	0
Total	0.22 ± 0.04	0.02 ± 0.01	0.0002 ± 0.0001	0.25 ± 0.05	2.64 ± 0.53	5



ICARUS


Proof of concept of
a large LAr TPC

Limits on sterile neutrinos
from analysis of ν_e events



High LAr Purity
reached

Free electron attenuation length [m]






NEUTRINO PLATFORM

R&D for future beam neutrino programs



ICARUS refurbishment for installation on the FNAL SBL ν beam (sterile ν searches)



1st vessel to be transported soon




1x1x3 m³ Double Phase Lar TPC prototype
Being commissioned for cosmics measurements

NEUTRINO PLATFORM

Large engineering detectors for DUNE




*Single Phase:
ProtoDUNE-SP*




*Double Phase:
ProtoDUNE-DP*

NEUTRINO PLATFORM



*Engineering prototypes to be calibrated
in low energy beams in a North Hall extension*




Hall extension ready
But tight schedule to take beam data before LS2

CERN NEUTRINOS TO GRAN SASSO Underground structures at CERN

AWAKE


- Excavated
- Concreted
- Decay tube (2nd contract)

*R&D for electron acceleration
with a plasma cell excited
by proton bunches*



**AWAKE currently taking first beam data:
goal to establish plasma modulation in 2016
and electron acceleration in 2017**


*A project of interest for future
high E / high I electron beams*



C. Vallee, CERN, 26.11.2010







Physics Beyond Colliders at CERN




DIRAC @ PS

Low E perturbative chiral QCD with mesonic atoms:
Discovery of πK atoms and metastable $\pi\pi$ atoms





***AFTER LS2: wish to perform similar studies at SPS (statistics $\times \sim 20$)
 would allow quantitative test of chiral $SU(3)_L \times SU(3)_R$ symmetry breaking***



NA61/SHINE


Search for QCD Critical Point
by scan in the (T, μ_B) plane

Scan to be completed until LS2
No indication of CP yet




AFTER LS2:

wish to further study
QCD deconfinement
with open charm



S-INE :
2020

REQUIRED FACILITY UPGRADES




Would allow to disentangle statistical/dynamical models
in complement of J/ψ data from NA38/NA50

NB new idea : NA60+


revival of dimuon studies in Heavy Ions

Could a single expt. measure both open and bound charm ?





NA61 large acceptance TPC
also unique to constrain
 ν beam fluxes

Heavily used by T2K with
p-C and p-replica target data



Similar program starting
with the US for LBNF







COMPASS




a large acceptance spectrometer in the intermediate x-domain between H1/ZEUS and HERMES/JLAB




COMPASS I

Data taking completed in 2012, focused on quark spin measurements with muon beams


Longitudinal target polarization \longleftrightarrow Transverse target polarization






Improved precision on g_1 at low x







Non-zero
transversity
in proton
confirmed



COMPASS II (2014-18)




2016-17: DVCS : proton tomography with access to orbital momentum of quarks



2014+15+18: DY : Transverse Momentum Dependent (TMD) QCD effects in the valence regime Measurement complementary to SiDIS : opposite asymmetries expected

AFTER LS2: wish RF separated antiproton and kaon beams (1 x 50)



- High statistics strange meson spectroscopy
- Exotic states spectroscopy complementary to LHCb
- Kaon and antiproton structure

Two body thresholds

Molecules

Gluonic Excitations


Hybrids

Hybrids+Recoil

Glueballs

Glueballs+Recoil

$q\bar{q}$ Mesons




DY statistics


	NH ₃	Al (7cm)	W	NA3	NA10	E537	E615
K^- beam	14,000	2,800	29,600	700			
\bar{p} beam	15,750	2,750	22,500			387	

New idea: direct measurement of the dominant contribution to the theoretical error on $(g-2)_\mu$ from μ -e elastic scattering

High statistics space-like measurement could reduce by factor 2 the current error derived from time-like processes




Vacuum polarisation



Might be feasible with reasonable resources within the (modified) COMPASS setup


New idea: Fixed Target physics with LHC beams

Internal gas target (AFTER)




e.g. SMOG

Upstream
of LHCb
and/or
ALICE




p-p: High precision TMD measurements
(polarized target) and charm at high x
p-A: Nuclear PDFs





New idea: Fixed Target physics with LHC beams

Crystal extraction



Upstream
of LHCb
and/or
ALICE

$$\frac{dN_i}{N_{0i} d\cos\theta_i} = \frac{1}{2} (1 + \alpha P_i \cos\theta_i)$$




Proposed for measurement of
magnetic moments of short lived baryons

*Could test anomalous magnetic
moments of heavy quarks*

ANTIMATTER FACTORY


Antiproton Decelerator




4 running experiments devoted to Antiproton and Antihydrogen properties




BASE



ASACUSA







ATRAP



ALPHA

2.5 more in preparation to test gravity of Antihydrogen: AEGIS/GBAR/ALPHA-g

Antiproton Properties



$$\frac{(-q/m)_{\bar{p}}}{(q/m)_p} - 1 = 1(69) \times 10^{-12}$$

Magnetic moment:

ATRAP gain in precision of ~3 orders of magnitude using new method with single trapped antiproton

Significant improvement expected soon from BASE

Mass:





Regular ASACUSA progress with cold 1- and 2-photon spectroscopy of antiprotonic Helium

Charge/Mass:

High precision BASE measurement with cyclotron frequency


Antihydrogen Properties

Hbar trapping established by 3 experiments



Spin flip with MW causes immediate ejection of Hbar from trap

ALPHA



First Hbar microwave interaction observed by ALPHA



Hbar Laser spectroscopy expected soon

Antihydrogen Properties cont'd: gravitation



2.5 experiments now devoted to a direct measurement

AEGIS

in-flight deviation of
Hbar atoms by gravitation



GBAR: Hbar free fall using ELENA



ALPHA-g

Statistical method
for a first measurement of the sign


AFTER LS2: ELENA

*Further deceleration
of antiprotons from
5 MeV to 100 KeV
kinetic energy*

*Will increase by 2 orders
of magnitude the antiproton
trapping efficiency*

*Under commissioning for
first connection to GBAR in 2017*


**Secures antimatter physics
for the next decade**



$$K \rightarrow \pi \nu \bar{\nu} \quad (BR \sim 10^{-11})$$


NA62

Rare K decays




From concept...

Separated from signal by kinematic cuts :



Signal regions




1 GHz 75 GeV unseparated beam,
11 MHz K⁺ decays in detector

$$K \rightarrow \pi \nu \bar{\nu} \quad (BR \sim 10^{-10})$$


NA62

Rare K decays



...to reality !

After many years of intensive construction and commissioning



Signal regions: ~100 evts expected until LS2



Detector fully operational in 2016, first year of quasi-nominal operation

New idea: $K^0 \rightarrow \pi^0 \nu \bar{\nu}$ rare decay


Both decays are complementary and allow constraining the CKM matrix.

Would require a new high intensity K^0 beam.

~50 events could be collected with a similar but basically new detector.

Competition from starting KOTO at JPARC:


few evts expected in coming years, upgrade to ~100 evts by 2025



C. Vallée, CPPM, 28.11.2016

Detector layout for $K_L \rightarrow \pi^0 \nu \bar{\nu}$

Similar to NA62 but basically new detector



Physics Beyond Colliders at CERN


Intermezzo: the Hidden Sector

$$L = L_{SM} + L_{mediator} + L_{HS}$$

Visible Sector



Mediators or portals to the HS:
vector, scalar, axial, neutrino





Hidden Sector

*Naturally accommodates Dark Matter
(may have rich structure)*


- Long-lived objects
- Interact very weakly with matter

Models	Final states
HNL, SUSY neutralino	$l^+\pi^-, l^+K^-, l^+\rho^-, \rho^+\rightarrow\pi^+\pi^0$
Vector, scalar, axion portals, SUSY sgoldstino	l^+l^-
HNL, SUSY neutralino, axino	$l^+l^-\nu$
Axion portal, SUSY sgoldstino	$\gamma\gamma$
SUSY sgoldstino	$\pi^0\pi^0$

Intermezzo cont'd: the Hidden Sector




2 methods



**Production + decay of new particle:
2 couplings → needs high intensity**

**Invisible decay of new particle:
accommodates lower intensity**

*A similar situation as the search for neutrino oscillations in the 70 – 80's:
do not know if they exist and where they stand !*




Fixed Target searches complementary to others

AFTER LS2 : NA62+


*Wish to run ~1 year in beam dump mode
to look for Heavy Neutral Leptons*

→ possible intermediate step towards
a more ambitious beam dump facility



Compact beam dump: $\sim 11 \lambda_1$ Cu-based beam-defining collimator (TAX)
radioprotection-compliant even if target removed


Decay volume ~ 60 m long (in vacuum):
reasonable acceptance to long-lived states



New idea: SHiP




Similar layout as NA62,
with larger acceptance to
reach the c / b mass range

*Beam Dump Facility
already under study
at CERN*




Flagship program for a comprehensive investigation
of the Hidden Sector in the few GeV domain
Exploits the unique high-E/ high-I SPS features

SHiP physics reach




Significant & mostly unique extension of reach for many channels




NA64

Hidden sector search from invisible decays with missing energy

First implementation in 2016
on an electron test beam





Exclusion of $(g-2)_\mu$ interpretation
with 1 day of data taking
confirms the potential of the method



AFTER LS2: NA64+

Wish to extend the method to $\mu / \pi / K / p$ beams






Process	New Physics	Sensitivity
1. $e^- Z \rightarrow e^- Z + E_{\text{miss}}$		
<ul style="list-style-type: none"> $A' \rightarrow e^+ e^-$ $A' \rightarrow \text{invisible}$ alps milli-q 	Dark Sectors: Dark Photons and DM $(g-2)_\mu$ new particles, Charge Quantization	$10^{-3} < \varepsilon < 10^{-6}$ $M_{A'} \sim \text{sub-GeV}$ $e' < 10^{-5} - 10^{-7}$
2. $\mu^- Z \rightarrow \mu^- Z + E_{\text{miss}}$		
<ul style="list-style-type: none"> $Z_\mu \rightarrow \nu \nu, \mu^+ \mu^-$ $\mu \rightarrow \tau$ conversion 	New gauged symmetry $L_\mu - L_\tau$ and leptonic forces LFV	$\alpha_\mu < 10^{-11} - 10^{-9}$ $\sigma < 10^{-9} - 10^{-8} / \mu$
3. $\pi(K)p \rightarrow M^0 n + E_{\text{miss}}$		
<ul style="list-style-type: none"> $K_L \rightarrow \text{invisible}$ $K_S \rightarrow \text{invisible}$ $\pi^0, \eta, \eta \rightarrow \text{invisible}$ 	CP, CPT symmetry B-S Unitarity, new particles: NHL, $\phi\phi$, VV	$\text{Br} < 10^{-8} - 10^{-6}$, complementary to $K \rightarrow \pi \nu \nu$ $\text{Br} < 10^{-8} - 10^{-7}$
4. $pA \rightarrow X + E_{\text{miss}}$		
<ul style="list-style-type: none"> leptophobic X 	$\sim \text{GeV DM}$	$\sigma < 10^{-7} - 10^{-8} / p$

Another possible source
of hidden particles:

Solar Axions from the sun

CAST: Instrumented LHC magnet pointed
to the sun to convert Axions into X rays





^3He and ^4He scans completed,
start to bite into QCD models

Vacuum runs being continued
together with R&D on
low noise detectors

New idea: IAXO



Next generation Axion Helioscope beyond CAST



Wish to profit from CERN magnet expertise (ATLAS-like large bore toroid)

Laboratory Axions: OSQAR/ALPS

Light shining through a wall





Comparable limits obtained by
OSQAR@CERN and ALPS@DESY

*A combined project ("ALPS III") could benefit
from CERN high field magnet developments*

New idea: DARKSIDE@LNGS



“Ultimate” WIMP search with depleted LAr double phase TPC

Wish to exploit synergies with CERN
on LAr, cryogeny, low noise SiPMs, etc...




New idea: Storage Ring for proton EDM

*incl. electrostatic option
applicable to proton only*



J.M.Pendlebury and E.A. Hinds, NIMA 440 (2000) 471





*10⁻²⁹ e-cm sensitivity
would correspond to 100 TeV
for new physics energy scale*

New idea: NuSTORM


Well controlled ν beam from a μ storage ring.

*Would allow precise $\sigma(\nu)$ measurements.
Also a path towards a ν factory or a μ collider.*



New idea: Gamma Factory

Use LHC beam to convert laser photons into 0.1 - 400 MeV γ rays



NEXT STEPS

Working Groups being set up :

- **Accelerator WG to study possible implementation of the projects at CERN.**
Members: CERN accelerator people + projects proponents
- **Physics WG to study the physics case in worldwide context and optimize detectors including siting options.**
Members: theorists and experimentalists + projects proponents

NB: involvement will be tuned to the level of maturity of the projects

Follow-up PBC workshop foreseen in 2017.

Final deliverable due end 2018:

Summary document as input to the European Strategy Update process (2019-20).
Will gather facts on the projects (no ranking!) to facilitate future orientations from the ESU group.

SPARE SLIDES

ACCELERATOR WORKING GROUP STRUCTURE

Subgroups:

Beam Dump Facility	:	Technical feasibility of BDF as input to SHiP CDS
EDM ring	:	Fully developed feasibility study incl. preliminary costing
Conventional beams	:	Study upgrades for NA62+, NA64+, COMPASS+, DIRAC+... beams
LHC Fixed Target	:	CDR putting together UA9, LHC Collimation, AFTER...
Technology	:	Evaluation of possible CERN contributions to non-acc. projects

Studies:

Complex performance:		Performance plan in LIU era and exploration of new proton driver
AWAKE	:	Exploratory study of possible applications of AWAKE concept
NuSTORM	:	Broad outline of possible implementation at CERN
Gamma Factory	:	Exploratory study incl. initial tests


PHYSICS WORKING GROUP STRUCTURE

Deliverables for each proposed project:

- Evaluation of the physics case in the worldwide context
- Possible further optimization of the detector
- For new projects: investigation of the worldwide siting options

BSM subgroup : SHiP/NA64+/NA62+/IAXO/OSQAR-ALPS-III/EDM

QCD subgroup : COMPASS+/ μ -e/LHC-FT/DIRAC+/NA60+/NA61+



NEUTRINO PLATFORM: BABYMIND

Muon spectrometer for muon charge tagging in WAGASCI experiment at JPARC

Alternance of magnetized iron plates and scintillator bars detection plates



Under assembly at CERN for beam tests and transport to Japan in 2017

Dimeson atom production at proton momentum 450 GeV/c

The dimeson atom production in p -nucleus interaction can be enlarged by more than an order of magnitude if the incident proton momentum is increased from 24 to 450 GeV/c.



With the SPS operation conditions at 450 GeV/c ($\theta_{lab} = 4^\circ$) the yield, i.e. the number of produced dimeson atoms $A_{2\pi}$, $A_{\pi^+ K^-}$ and $A_{\pi^- K^+}$ per time unit, will be 12 ± 2 , 53 ± 11 and 24 ± 5 times higher than in the previous DIRAC experiment (O.Gorchakov, L.Nemenov J. Phys. G: Nucl. Part., 2016).

This significant increase in the $A_{\pi^+ K^-}$ and $A_{\pi^- K^+}$ production allows



- to measure with a new DIRAC setup in a comparable running time, $|a_{1/2} - a_{3/2}|$ with a precision better than 5% and
- to check with the same accuracy predictions of the total $\mathcal{L}(3)$ QCD Lagrangian based on the chiral $SU(3)_L * SU(3)_R$ symmetry breaking

COMPASS: SPECTROSCOPY AND PRIMAKOV

Publication of the new
isovector meson $a_1(1420) 1^{++}$




Publication of the pion polarisability
using 2009 Primakov data
(to be x 5 with 2012 data)



ALPHA-g Phase I: sign measurement in 2017-18

Trap and release:



- Maximum sensitivity requires slow ramping down of mirror coils
- Simulations accounting for real trajectories and longitudinal/transverse energy transfer give an optimum for $\sim 10 \text{ s}$ ramp down duration
- For “normal” gravity this corresponds to **71% of down escape**