Invisible Higgs decays and the GCE in the NMSSM

A. Butter ^{1,2} with T. Plehn, M. Rauch, D. Zerwas, S. Henrot-Versillé, and Rémi Lafaye

¹ITP, Universität Heidelberg, Germany

²LAL, CNRS/IN2P3, Orsay, France

arXiv:1507.02288





There is physics beyond the Standard Model!

- 26.8% dark matter
- strong constraining power on the parameterspace



SFitter likelihood Fit, mSUGRA



There is more ...?

• Since 2009 observation of an excess of gamma ray photons in 3-4 GeV range from the galactic center by Fermi LAT [Hooper et al., Calore et al., Murgia for Fermi Collaboration]



Fermi collaboration, arXiv:1511.02938

NMSSM 0000000

Correlations

γ spectrum for the GCE



D. Hooper, et al., arXiv:1402.6703

NMSSM 0000000 Correlations

Dark matter solutions to the GCE

- GCE can be explained by dark matter around a few GeV
- Observable at a collider?
- Higgs could decay in light dark matter !
- \rightarrow Can we link GCE to an enhanced branching ratio $H_{125} \rightarrow inv$?



D. Hooper, et al., arXiv:1402.6703

Correlations

$\label{eq:what are the main constraints?} What are the main constraints? \\ \bullet \ \Omega h^2 = 0.1187 \pm 0.0017 \ _{st} \pm 0.0120 \ _{th} \qquad \mbox{(Planck)}$

- Dark matter annihilation before the freeze out
- Examples:

. . .

h-funnel, A-funnel, Z-mediated, coannihilation [Drees, Nojiri],



Correlations

What are the main constraints?

- $\Omega h^2 = 0.1187 \pm 0.0017$ _{st} ± 0.0120 _{th}
- $\sigma v|_{v=0} \approx 2 \times 10^{-26} \,\mathrm{cm}^3/\mathrm{s}$

- excess of γ -ray photons in the galactic center observed by Fermi LAT
- well fit by 30 70 GeV dark matter annihilating via a **pseudoscalar** to $b\bar{b}$

$$\sigma v \Big|_{v=0} \approx \frac{3}{2\pi} \frac{g_{A\chi\chi}^2 g_{Abb}^2 m_{\chi}^2}{\left(m_A^2 - 4m_{\chi}^2\right)^2 + m_A^2 \Gamma_A^2}$$



(Planck)

(GCE)

Calore, Cholis, McCabe, Weniger, arXiv:1411.4647

What are the main constraints?

- $\Omega h^2 = 0.1187 \pm 0.0017 \ _{st} \pm 0.0120 \ _{th}$
- $\sigma v|_{v=0} \approx 2 \times 10^{-26} \,\mathrm{cm}^3/\mathrm{s}$
- BR $(H \to inv) < 23\%/36\%$

(Planck) (GCE) (ATLAS/CMS)

- Current measurements
 - CMS: WBF
 - ATLAS: combined associated production and gluon fusion
- Future: probing down to 3% with high luminosity run [Plehn et al.]
- \rightarrow We require BR> 10%

What are the main constraints?

- $\Omega h^2 = 0.1187 \pm 0.0017$ _{st} ± 0.0120 _{th}
- $\sigma v|_{v=0} \approx 2 \times 10^{-26} \,\mathrm{cm}^3/\mathrm{s}$
- BR $(H \to inv) < 23\%/36\%$
- $m_H = (125.1 \pm 0.2 \ _{st} \pm 0.1 \ _{sy} \pm 3.0 \ _{th}) \, \text{GeV}$

(Planck) (GCE) (ATLAS/CMS) (ATLAS/CMS)

What are the main constraints?

- $\Omega h^2 = 0.1187 \pm 0.0017$ _{st} ± 0.0120 _{th}
- $\sigma v|_{v=0} \approx 2 \times 10^{-26} \,\mathrm{cm}^3/\mathrm{s}$
- BR $(H \to inv) < 23\%/36\%$
- $m_H = (125.1 \pm 0.2 \ _{st} \pm 0.1 \ _{sy} \pm 3.0 \ _{th}) \, \text{GeV}$ (ATLAS/CMS)
- $\Gamma_{Z \to Inv} = (-1.9 \pm 1.5 \ _{st} \pm 0.2 \ _{th}) \, \mathrm{MeV}$

```
(Planck)
(GCE)
(ATLAS/CMS)
(ATLAS/CMS)
(LEP)
```

- make sure that the coupling to the Z is small
- we require $\Gamma_{Z \rightarrow Inv} < 2$ MeV

NMSSM 0000000 Correlations

Why SUSY?

Other hints at physics beyond the Standard Model



- \rightarrow promising setup but many models
- \rightarrow Can we connect astronomical observations to LHC measurements in the SUSY framework?

NMSSM •000000 Correlations

The MSSM particle content



NMSSM 0000000 Correlations

From MSSM to NMSSM

- μ -Problem, light pseudoscalar for GCE
- solution:

$$W_{ extsf{NMSSM}} = W_{ extsf{MSSM}} + \lambda SH_uH_d + rac{1}{3}\kappa S^3$$

 \rightarrow additional singlet S

• vev of S generates effective μ -term: $\mu_{\textit{eff}} = \lambda \cdot s$



NMSSM 0000000 Correlations

The NMSSM parameters

• Scale invariant superpotential

$$W_{ extsf{NMSSM}} = W_{ extsf{MSSM}} + \lambda S H_u H_d + rac{1}{3} \kappa S^3$$

• Additional soft SUSY breaking parameters appear:

$$-\mathcal{L}_{soft} \supset \lambda A_{\lambda} H_{u} H_{d} S + rac{1}{3} \kappa A_{\kappa} S^{3}$$

Complete set of input parameters

$$\begin{array}{ll} m_{\tilde{q},\tilde{\ell}}, M_{1,2,3}, A_{q,\ell}, \tan\!\beta, \mu & \to \text{as for MSSM} \\ \lambda, \kappa, A_{\lambda}, A_{\kappa} & \to \text{additional parameters} \end{array}$$

• Decouple gluino and sfermion sector $\rightarrow m_{\tilde{q},\tilde{\ell}} = M_3 = 10$ TeV, $A_{q,\ell} = 0 \rightarrow 8$ free parameters left

Constraints on the spectrum

- Standard Model like Higgs at 125 GeV $[H_{125}]
 ightarrow m_{ ilde{t}} = 6$ TeV
- charged Higgs must be heavy
 - ightarrow MSSM like scalar and pseudoscalar heavier than SM Higgs
- GCE:
 - $ightarrow~m_{ ilde{\chi}_{1}^{0}}pprox$ 30 70 GeV
 - \rightarrow light (singlet like) pseudoscalar $m_{A_1} \approx 2 \cdot m_{\tilde{\chi}_1^0}$
 - $ightarrow A_\kappa = -250 \; {
 m GeV}$
- LEP constraints:
 - $\rightarrow m_{\tilde{\chi}_1^{\pm}} > 103 \text{ GeV} \rightarrow \mu, M_2 > 100 \text{ GeV}$ negligible wino contribution on LSP \rightarrow set $M_2 = 2 \text{ TeV}$
- BR $H_{125} \rightarrow \tilde{\chi}_1^0 + \tilde{\chi}_1^0$: $\rightarrow m_{\tilde{\chi}_1^0} < 64 \text{ GeV}$

Free parameters

 $\mathit{M}_{1}, \mu, \mathit{A}_{\lambda}, \tan\beta, \lambda, \kappa$

NMSSM 0000000 Correlations

Constraints on the couplings

• SM Higgs couplings:

 \rightarrow no tree level mixing between singlet and ${\it H}_{\rm 125}$

$$A_{\lambda} = 2\mu \left(rac{1}{s_{2eta}} - ilde{\kappa}
ight)$$
 define $ilde{\kappa} := rac{\kappa}{\lambda};$ $(m_{ ilde{\chi}s} = 2\mu ilde{\kappa})$

• Z neutralino (Ωh^2 , LEP)

$$g_{Z\tilde{\chi}_{1}\tilde{\chi}_{1}} = \frac{g}{2\cos\theta_{W}}\gamma_{\mu}\gamma_{5}\left(N_{13}^{2} - N_{14}^{2}\right)$$

• A_S neutralino (GCE, Ωh^2)

$$g_{A_{S}\tilde{\chi}_{1}\tilde{\chi}_{1}} = \sqrt{2}\lambda \left(N_{13}N_{14} - \tilde{\kappa}N_{15}^{2} \right)$$

Fit parameters $M_1, \mu, \lambda, ilde{\kappa}$



17 / 25

NMSSM 0000000 Correlations

Relic density and GCE



- m_A increases with $(\lambda \cdot \mu)^{-1}$
- $g_{Z\tilde{\chi}_1^0\tilde{\chi}_1^0}$ increases with $\lambda\cdot\mu^{-1}$ [Higgsino component]
- region compatible with GCE and Ωh² is very narrow!
- Higgsino component too small for an enhanced BR

NMSSM 0000000 Correlations •0000

Annihilation channels in the bino-singlino plane



- choose $\mu = 320$ GeV, variable input parameters: $\lambda, \tilde{\kappa}, M_1$
- criteria: SM Higgs, relic density, Xenon100, BR> 10%
- $\tilde{\kappa} M_1$ plane:
 - asymptotic behaviour of the LSP mass
 - symmetric under bino singlino exchange

NMSSM 0000000 Correlations •0000

Annihilation channels in the bino-singlino plane



- Dark matter annihilation:
 - H_{125} mediated annihilation for $m_{\tilde{\chi}_1^0} \approx 55$ GeV
 - Z mediated for $m_{\chi_1^0} \approx 40$ and 48 GeV
 - cancellation of Higgsino components for $\tilde{\kappa} \approx 0.08$
 - H_1 mediated annihilation for $m_{\tilde{\chi}_1^0} < 40$ GeV

NMSSM 0000000 Correlations •0000

Annihilation channels in the bino-singlino plane



- The galactic center excess [black points]
 - neutralino mass constrained to $m_{\tilde{\chi}_1^0} > 30$ GeV [Weniger et al.]
 - m_{A_1} decreases with $\tilde{\kappa}$ and $1/\lambda$
 - GCE criteria fulfilled for $m_{A_1} \lesssim 2m_{\tilde{\chi}_1^0}$

NMSSM 0000000 Correlations

Branching ratio $H_{125} \rightarrow \tilde{\chi}_1^0 + \tilde{\chi}_1^0$



largest BR for mixed state (45% bino, 45 % singlino, 10% Higgsino)

 \rightarrow 0.45 not compatible with GCE

- BR $(H_{125} \rightarrow \tilde{\chi}^0_1 \tilde{\chi}^0_1) \approx 15\%$ compatible with the GCE
 - \rightarrow requires singlino like dark matter

NMSSM 0000000 Correlations

μ -dependence of the branching ratio



- larger Higgsino component $[1/\mu]$
- annihilation channels no longer well separated
- BR $(H_{125}
 ightarrow ilde{\chi}_1^0 ilde{\chi}_1^0) pprox$ 60% for $\mu=$ 190 GeV

ightarrow scenarios with μ < 250 GeV are excluded by ATLAS and CMS!

NMSSM 0000000 Correlations

The NUH-NMSSM

- Non Unified Higgs mass
- Additional unification criteria: $m_0, m_{1/2}, A_0, A_\lambda, A_\kappa$ at GUT scale $\rightarrow M_1 > 200 \text{ GeV [gluino searches]}$
- Free parameters $\tilde{\kappa}, \lambda, A_{\kappa}, \mu$
- Two regions correspond to Z-mediated annihilation
- GCE compatible with $m_{ ilde{\chi}_1^0}pprox$ 35 GeV
- Invisible branching ratio up to 40 % for $\mu \approx 160~{\rm GeV}$







NMSSM 0000000



Conclusion

- (NUH)-NMSSM is a suitable setup to describe the GCE
- Combining the GCE with the relic density and Xenon100 strongly constrains the parameter space
- BR $(H_{125} \rightarrow \tilde{\chi}_1^0 + \tilde{\chi}_1^0)$ up to 60% is possible in agreement with GCE and is tested by the LHC!

BACK UP

Neutralino mass matrix

$$M_{\tilde{\chi}^{0}} = \begin{pmatrix} M_{1} & 0 & -m_{Z}c_{\beta}s_{w} & m_{Z}s_{\beta}s_{w} & 0\\ 0 & M_{2} & m_{Z}c_{\beta}c_{w} & -m_{Z}s_{\beta}c_{w} & 0\\ -m_{Z}c_{\beta}s_{w} & m_{Z}c_{\beta}c_{w} & 0 & -\mu & -\lambda vs_{\beta}\\ m_{Z}s_{\beta}s_{w} & -m_{Z}s_{\beta}c_{w} & -\mu & 0 & -\lambda vc_{\beta}\\ 0 & 0 & -\lambda vs_{\beta} & -\lambda vc_{\beta} & 2\tilde{\kappa}\mu \end{pmatrix}$$

CP even Higgs mass marix in the basis (H, H_{125}, S)

$$M_{H,H_{125},S}^{2} = m_{Z}^{2} \begin{pmatrix} s_{2\beta}^{2} \left(1 - \tilde{\lambda}^{2}\right) + \frac{2\mu}{c_{2\beta}m_{Z}^{2}} \left(A_{\lambda} + \tilde{\kappa}\mu\right) & c_{2\beta}s_{2\beta} \left(1 - \tilde{\lambda}^{2}\right) & -c_{2\beta}\frac{\tilde{\lambda}}{m_{Z}} \left(A_{\lambda} + \tilde{\kappa}\mu\right) \\ \cdot & c_{2\beta}^{2} + s_{2\beta}^{2}\tilde{\lambda}^{2} & \frac{2\tilde{\lambda}}{m_{Z}} \left(\mu - s_{2\beta}\frac{A_{\lambda}}{2} + s_{2\beta}\tilde{\kappa}\mu\right) \\ \cdot & \cdot & s_{2\beta}\frac{\tilde{\lambda}^{2}A_{\lambda}}{2\mu} + \frac{\tilde{\kappa}\mu}{m_{Z}^{2}} \left(A_{\kappa} + 4\tilde{\kappa}\mu\right) \end{pmatrix}$$

CP odd Higgs mass marix in the basis (A, S)

$$M_{A,S}^{2} = \begin{pmatrix} \frac{2\mu \left(A_{\lambda} + \tilde{\kappa}\mu\right)}{s_{2\beta}} & \tilde{\lambda}m_{Z} \left(A_{\lambda} - 2\tilde{\kappa}\mu\right) \\ \cdot & s_{2\beta}\tilde{\lambda}^{2}m_{Z}^{2} \left(\frac{A_{\lambda}}{2\mu} + 2\tilde{\kappa}\right) - 3\tilde{\kappa}\mu A_{\kappa} \end{pmatrix}$$

Minimal value of μ for a BR less than 0.3



- minimal value of $\mu \approx 200~{
 m GeV}$
- for smaller values the branching ratio becomes too large
- other points with a smaller value of μ are possible, if we accept a BR of 0 %