

# The global Higgs as a signal for compositeness at the LHC

Sylvain Fichet  
ICTP/SAIFR&IFT Sao Paulo

Based on 1607.03125 (JHEP) and 1608.01995 (JHEP) with  
Gero von Gersdorff, Eduardo Pontón, Rogerio Rosenfeld

LAPTh  
09/03/17

1. The composite Higgs paradigm
2. The global Higgs and its properties
3. LHC prospects
4. Outlook

1. The composite Higgs paradigm


# Is the Standard Model composite ?

Motivation 1: Why not ?

Motivation 2:  $m_h \ll M_{\text{Planck}}$  ?

Mass of spin-0 particle not protected by a symmetry. The mass of the Higgs boson is technically unnatural.

A solution: **NO spin-0 particle**. The Higgs is a **composite** state that dissolves above a scale  $\Lambda$ .



# Is the Standard Model composite ?

LHC produced a 125 GeV very SM-like Higgs resonance without revealing any obvious substructure → Binding should be quite strong  
→ **strongly coupled interaction**

But the 125 GeV SM-like Higgs is not accompanied by nearby resonances and has a narrow width... **Doesn't look like a regular hadron...**

# Composite pNGB Higgs

Borrow intuition from QCD: the pions are much lighter than other bound states, because they are **pseudo Nambu-Goldstone bosons** (pNGBs) of a spontaneously broken approximate global symmetry.



# Composite pNGB Higgs

Borrow intuition from QCD: the pions are much lighter than other bound states, because they are **pseudo Nambu-Goldstone bosons** (pNGBs) of a spontaneously broken approximate global symmetry.



# Composite pNGB Higgs


Borrow intuition from QCD: the pions are much lighter than other bound states, because they are **pseudo Nambu-Goldstone bosons** (pNGBs) of a spontaneously broken approximate global symmetry.






# Composite pNGB Higgs

**Idea:** The Higgs boson and the three longitudinal polarizations of the electroweak (EW) gauge bosons are pNGBs, the 'pions' of a global group  $G$  broken into  $G'$ . [Georgi/Kaplan/... '83]



The pNGB gap is not arbitrarily large without tuning.  
Hence  $f$  is expected to be  $O(\text{TeV}) \rightarrow$  resonances accessible at LHC


# Composite Higgs paradigm



2. The global Higgs and its properties

# Radius fluctuation


The  $G/G'$  coset has a finite radius [in most of constructions], which is assumed to be stable.



The fluctuation of this radius is a massive, CP-even scalar (just like the SM Higgs boson for the  $SU(2) \times U(1)/U(1)$  coset).


# Radius fluctuation

The  $G/G'$  coset has a finite radius [in most of constructions], which is assumed to be stable.



There is another scalar in the composite Higgs picture: the **GLOBAL HIGGS**, noted  $\phi$ .

The **fluctuation** of this radius is a massive, CP-even scalar (just like the SM Higgs boson for the  $SU(2) \times U(1)/U(1)$  coset).



Conceptual step : Global Higgs and pNGBs could be embedded into a multiplet of the unbroken group  $G$  (noted  $\Phi$ ).

# Qualitative features

- The global Higgs is expected to couple to the **NGB**'s parametrising the coset. Couplings should be derivative by pNGB shift symmetry  
 → One expects a coupling  $\phi(\partial_\mu h)^2$ , and thus  $\phi(Z_\mu)^2$  and  $\phi W_\mu^+ W^{\mu-}$  (by gauge invariance)

- should couple to **fermion resonances**, as one needs to write proto-Yukawa operators of the form

$$-\xi_{u_i} \mathcal{O}_u(\Phi) \bar{Q}_i U_i - \xi_{d_i} \mathcal{O}_d(\Phi) \bar{Q}_i D_i - \xi_{l_i} \mathcal{O}_l(\Phi) \bar{L}_i E_i + \text{h.c.}$$

- May couple to **SM fermions** proportionally to their mass,

$$\phi m_\psi (\psi_{\text{SM}} \psi'_{\text{SM}} + \text{h.c.})$$

# The $SO(5)/SO(4)$ Higgs

- The  $G=SO(5)$ ,  $G'=SO(4)$ , choose  $\Phi \sim \mathbf{5}$  of  $SO(5)$ .

$$\underbrace{\Phi = U_5 \mathcal{H}}_{\text{pNGBs}} \quad \underbrace{\mathcal{H} = (\hat{f} + \phi) e_5}_{\text{VEV}} \quad \underbrace{\phantom{(\hat{f} + \phi)} e_5}_{\text{global Higgs}}$$

radius

- Consider quartic expansion of the radius potential:

$$V(\mathcal{H}) = \frac{1}{4} \lambda \left( \mathcal{H}^2 - \hat{f}^2 \right)^2 \rightarrow m_\phi = \sqrt{2\lambda} \hat{f}$$

- General bosonic Lagrangian:

$$\mathcal{L}_{\text{bos}} = \frac{1}{2} (\nabla_\mu \mathcal{H})^2 - V(\mathcal{H}) + \frac{1}{4} f_\rho^2 \left( \mathcal{A}_\mu^A - i [U_5^\dagger D_\mu U_5]^A \right)^2 + \dots$$

The pNGBs mix with the coset resonances,  $f^{-2} = \hat{f}^{-2} + f_\rho^{-2}$



# Global Higgs interactions

- Bosonic couplings:

- SM Higgs and EW gauge bosons:  $2 \frac{f^2}{\hat{f}^3} \phi (D_\mu H)^2$

- Spin-1 (coset) resonances:  $\frac{2}{\hat{f}} \phi \mathcal{A}_\mu^{\hat{a}^2}$

- Fermionic couplings:

- SM fermions:  $-\frac{m_\psi}{\hat{f}} \phi \bar{\psi} \psi$

Only two  
free parameters !

And  $\hat{f}$  should be minimized  
to reduce EW fine-tuning.

- fermion resonances: completely model dependent

# Complete realizations

To learn further about the global Higgs properties, one needs to define **full realizations** of the fermion sector.

We consider four typical models,  $\text{MCHM}_{Q,U,D}$ , with

$$(Q, U, D) = (5_{2/3}, 1_{2/3}, 10_{2/3})$$

$$(5_{2/3}, 14_{2/3}, 10_{2/3})$$

$$(14_{2/3}, 14_{2/3}, 10_{2/3})$$

$$(5_{2/3}, 1_{2/3}, \times)$$

← Higgs in the 14 of  $SO(5)$

← No partial compositeness

**IN SHORT:** Yukawa operators and thus couplings of the global Higgs to fermion resonances differ in each scenario (multiplicities, group theoretical factors...)

# Renormalization



- Yukawa couplings  $\xi$ :

$$\xi_{\text{eff}}^2 = 4N_c \left( N^U [\xi_U \xi_U^T + \xi'_U \xi'^T_U] + N^D [\xi_D \xi_D^T + \xi'_D \xi'^T_D] \right)$$

$$\mu \frac{d\xi_{\text{eff}}^2}{d\mu} \approx \frac{\xi_{\text{eff}}^4}{16\pi^2}$$


- global Higgs quartic  $\lambda$ :

$$\mu \frac{d\lambda}{d\mu} \approx \frac{1}{16\pi^2} (26\lambda^2 + 2\lambda\xi_{\text{eff}}^2 - \epsilon \xi_{\text{eff}}^4)$$



# One-loop effective couplings

**We can now go ahead:** The global Higgs couples to many non SM-singlet fermion and vector resonances, which induce loop couplings to SM gauge bosons.



The fermion loops completely depend on the fermion sector (representation, loops, mass matrices). However, for heavy enough vector like masses, compact expressions are obtained.


# One-loop effective couplings

For example : gluon coupling

$$\mathcal{L} \supset -\phi \frac{a_{gg}}{\hat{f}} (G_{\mu\nu}^a)^2$$

$$a_{gg} = c_{gg} \frac{\hat{f}^2}{M_\psi^2} A_{1/2} \left( \frac{m_\phi^2}{4M_\psi^2} \right)$$

$$c_{gg} = \begin{pmatrix} 0.013 \\ 0.014 \\ 0.011 \\ 0.01 \end{pmatrix} \begin{matrix} \text{MCHM}_{5,1,10} \\ \text{MCHM}_{5,14,10} \\ \text{MCHM}_{14,14,10} \\ \text{MCHM}_{5,1} \end{matrix}$$



Very similar for each scenario !

3. Some LHC prospects

# Global Higgs parameters

- Global Higgs mass  $m_\Phi$
- Global Higgs quartic  $\lambda$
- Fermion resonances mass scale  $M_\Psi$
- pNGB decay constant  $f$

# Global Higgs parameters

- Global Higgs mass  $m_\Phi$
- Global Higgs quartic  $\lambda \in [0.2, 3]$  from RG
- Fermion resonances mass scale  $M_\Psi$
- pNGB decay constant  $f$




# Global Higgs parameters

- global Higgs mass  $m_\Phi$
- global Higgs quartic  $\lambda \in [0.2, 3]$  from RG
- Fermion resonances mass scale  $M_\Psi$
- pNGB decay constant  $f = 800 \text{ GeV}$

from standard LE bounds on composite models. In this work we don't focus on LE effects from the global Higgs.

# Production at the 13 TeV LHC

I



Red and purple :  $M_\psi = m_\phi$ ,  $M_\psi = 2m_\phi$   
 Plain, dashed, dotted :  $\lambda = 0.2, 1, 3$

# Scenarios



The LHC signals of the global Higgs can be split into two broad cases:

- **Case I:** All decays involving fermion resonances are closed. The phenomenology is then largely independent of the details of the heavy fermion sector.  
Leading decays are  $\phi \rightarrow \text{pNGBs}$ ,  $\phi \rightarrow t\bar{t}$
- **Case II:** Some decays involving fermion resonances  $\phi \rightarrow \bar{\psi}\psi'$  are also open. The phenomenology depends strongly on the realization of the fermion sector.

Case I

# Decays


If no decays to fermion resonances are allowed, the global Higgs mainly decays into Higgs, EW bosons and top quarks.



Total width doesn't exceed  $\Gamma/m_\phi \sim 0.1$ , the global Higgs is a narrow state and NWA applies.

# Decays


If no decays to fermion resonances, the branching ratios are



# Prospects for decays into SM particles

## Sensitivities :

- $4l$ ,  $t\bar{t}$  directly obtained from exp. sensitivity
- $JJ$  obtained by extrapolating a 8 TeV  $JJ$  background from ATLAS



Case II



# Decays

If decays to fermion resonances are **open**, the global Higgs width grows. In the extreme limit where all decays are open and masses negligible, one has simply

$$\frac{\Gamma_{\phi \rightarrow \bar{\psi}\psi}}{m_\phi} = \begin{pmatrix} 27/4\pi \\ 54/5\pi \\ 117/20\pi \\ 3/4\pi \end{pmatrix} |\xi|^2 \approx \begin{pmatrix} 0.8 \\ 0.9 \\ 0.7 \\ 0.6 \end{pmatrix},$$

MCHM<sub>5,1,10</sub>


MCHM<sub>5,14,10</sub>

MCHM<sub>14,14,10</sub>

MCHM<sub>5,1</sub>


# Decays into top partners

When the global Higgs decays into fermion partners, there is a larger model-dependence. We consider the case of decays into top partners ( $t'$ ) only, like



# Decays into top partners


When the global Higgs decays into fermion partners, there is a larger model-dependence. We consider the case of decays into top partners ( $t'$ ) only, like



- A new production mode for  $t'$
- Resonant topology is useful to reject background (not studied)
- If hadronic decays, boosted  $t'$  may produce large-radius jets

# Fat $t'$ jets

Can these actually happen ?



## 4. Conclusions

# Summary (1)

- Composite Higgs models contain another scalar: the excitation of global-symmetry breaking vacuum  $G/G'$  -dubbed the **global Higgs**, coupled in a predictive way to Higgs, Z, W, top.
- We considered complete fermion sector realizations, taking into account renormalization of the composite sector. Sizeable couplings to **gauge bosons** arise from resonance loops.
- We evaluated the LHC sensitivity to global Higgs resonant production. Our results suggest that these channels can compete with the standard searches for compositeness via SM-produced  $t'$ .

## Summary (2)

- If the global Higgs decays mostly into **NGBs** and **top quarks**, sensitivity in boosted hadronic channels gives a reach up to  **$m \sim 2-2.5 \text{ TeV}$**  for  $300 \text{ fb}^{-1}$ . This case is very predictive.
- If global Higgs also decay also into  $t'$ , such channel can **compete** with standard  $t'$  production. Moreover we find these boosted  $t'$  have a sizeable probability to produce  **$t'$ -jets**.

# Outlook

A number of exciting directions can be pursued, including:

- Complete analyses including jet substructures for the resonantly produced, boosted  $t'$  channels
- If global Higgs light enough, main signals for  $t'$  production can be a **multi-top** final state
- Investigating possible charged partners of the global Higgs. For ex, if global Higgs embedded in the 14 of  $SO(5)$ , a heavy  $\eta$  of  $SO(4)$  is also present.



Thanks!


More

# Prospects for decays into SM particles

When  $\phi \rightarrow \bar{\psi}\psi$  closed,

(leading decays are  
 $\phi \rightarrow \text{pNGBs}$   $\phi \rightarrow t\bar{t}$  )

Decays depends on  
 2 parameters, chosen to  
 be the global Higgs  
 mass and quartic.



Thanks!