

Exclusive photoproduction of a $\gamma\rho$ pair with a large invariant mass

Samuel Wallon

Université Pierre et Marie Curie

and

Laboratoire de Physique Théorique

CNRS / Université Paris Sud

Orsay

Nucleon and Resonance Structure with Hard Exclusive Processes

Orsay, 29 May 2017

in collaboration with


B. Pire (CPHT, Palaiseau), R. Boussarie (LPT Orsay), L. Szymanowski (NCBJ, Warsaw)

JHEP 1702 (2017) 054 [arXiv:1609.03830](https://arxiv.org/abs/1609.03830) [hep-ph]

Transversity of the nucleon using hard processes: using a two body final state process?

How to get access to transversity GPDs?

- the dominant DA of ρ_T is of twist 2 and chiral-odd ($[\gamma^\mu, \gamma^\nu]$ coupling)
- unfortunately $\gamma^* N^\uparrow \rightarrow \rho_T N' = 0$
 - This cancellation is true at any order : such a process would require a helicity transfer of 2 from a photon.
 - lowest order diagrammatic argument:



$$\gamma^\alpha [\gamma^\mu, \gamma^\nu] \gamma_\alpha \rightarrow 0$$

[Diehl, Gousset, Pire], [Collins, Diehl]


Transversity of the nucleon using hard processes: using a two body final state process?

Can one circumvent this vanishing?


- This vanishing only occurs at **twist 2**
- At twist 3 this process does not vanish [Ahmad, Goldstein, Liuti], [Goloskokov, Kroll]
- However processes involving **twist 3 DAs** may face problems with factorization (end-point singularities)
can be made safe in the high-energy k_T -factorization approach [Anikin, Ivanov, Pire, Szymanowski, S.W.]
- One can also consider a 3-body final state process [Ivanov, Pire, Szymanowski, Teryaev], [Enberg, Pire, Szymanowski], [El Beiyad, Pire, Segond, Szymanowski, S. W.]

Probing GPDs using ρ meson + photon production

- We consider the process $\gamma N \rightarrow \gamma \rho N'$
- Collinear factorization of the amplitude for $\gamma + N \rightarrow \gamma + \rho + N'$ at large $M_{\gamma\rho}^2$




large angle factorization
à la Brodsky Lepage



Probing chiral-even GPDs using ρ meson + photon production

Processes with 3 body final states can give access to chiral-even GPDs




chiral-even twist 2 DA

chiral-even twist 2 GPD

Probing chiral-odd GPDs using ρ meson + photon production

Processes with 3 body final states can give access to chiral-odd GPDs




chiral-odd twist 2 GPD

Probing **chiral-odd** GPDs using ρ meson + photon production

Processes with **3 body final states** can give access to **chiral-odd GPDs**

How did we manage to circumvent the no-go theorem for $2 \rightarrow 2$ processes?




Typical non-zero diagram for a **transverse** ρ meson

the σ matrices (from DA and GPD sides) do not kill it anymore!

Master formula based on leading twist 2 factorization

$$\mathcal{A} \propto \int_{-1}^1 dx \int_0^1 dz T(x, \xi, z) \times H(x, \xi, t) \Phi_\rho(z) + \dots$$

- Both the DA and the GPD can be either **chiral-even** or **chiral-odd**.
- At twist 2 the **longitudinal ρ DA** is **chiral-even** and the **transverse ρ DA** is **chiral-odd**.
- Hence we will need both **chiral-even** and **chiral-odd** non-perturbative building blocks and hard parts.



Kinematics

Kinematics to handle GPD in a 3-body final state process

- use a **Sudakov** basis :
light-cone vectors p , n with $2p \cdot n = s$
- assume the following kinematics:
 - $\Delta_{\perp} \ll p_{\perp}$
 - $M^2, m_{\rho}^2 \ll M_{\gamma\rho}^2$

- initial state particle momenta:


$$q^{\mu} = n^{\mu}, \quad p_1^{\mu} = (1 + \xi) p^{\mu} + \frac{M^2}{s(1+\xi)} n^{\mu}$$

- final state particle momenta:

$$p_2^{\mu} = (1 - \xi) p^{\mu} + \frac{M^2 + \vec{p}_t^2}{s(1 - \xi)} n^{\mu} + \Delta_{\perp}^{\mu}$$

$$k^{\mu} = \alpha n^{\mu} + \frac{(\vec{p}_t - \vec{\Delta}_t/2)^2}{\alpha s} p^{\mu} + p_{\perp}^{\mu} - \frac{\Delta_{\perp}^{\mu}}{2},$$

$$p_{\rho}^{\mu} = \alpha_{\rho} n^{\mu} + \frac{(\vec{p}_t + \vec{\Delta}_t/2)^2 + m_{\rho}^2}{\alpha_{\rho} s} p^{\mu} - p_{\perp}^{\mu} - \frac{\Delta_{\perp}^{\mu}}{2},$$



Non perturbative **chiral-even** building blocks

- Helicity conserving GPDs at twist 2 :

$$\int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p_2, \lambda_2 | \bar{\psi}_q \left(-\frac{1}{2}z^- \right) \gamma^+ \psi \left(\frac{1}{2}z^- \right) | p_1, \lambda_1 \rangle$$

$$= \frac{1}{2P^+} \bar{u}(p_2, \lambda_2) \left[H^q(x, \xi, t) \gamma^+ + E^q(x, \xi, t) \frac{i\sigma^{\alpha+} \Delta_\alpha}{2m} \right]$$

$$\int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p_2, \lambda_2 | \bar{\psi}_q \left(-\frac{1}{2}z^- \right) \gamma^+ \gamma^5 \psi \left(\frac{1}{2}z^- \right) | p_1, \lambda_1 \rangle$$

$$= \frac{1}{2P^+} \bar{u}(p_2, \lambda_2) \left[\tilde{H}^q(x, \xi, t) \gamma^+ \gamma^5 + \tilde{E}^q(x, \xi, t) \frac{\gamma^5 \Delta^+}{2m} \right]$$

- We will consider the simplest case when $\Delta_\perp = 0$.
- In that case and in the forward limit $\xi \rightarrow 0$ only the H^q and \tilde{H}^q terms survive.
- Helicity conserving (vector) DA at twist 2 :

$$\langle 0 | \bar{u}(0) \gamma^\mu u(x) | \rho^0(p, s) \rangle = \frac{p^\mu}{\sqrt{2}} f_\rho \int_0^1 du e^{-iup \cdot x} \phi_{\parallel}(u)$$

Non perturbative **chiral-odd** building blocks

- Helicity flip GPD at twist 2 :

$$\begin{aligned}
 & \int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p_2, \lambda_2 | \bar{\psi}_q \left(-\frac{1}{2}z^- \right) i\sigma^{+i} \psi \left(\frac{1}{2}z^- \right) | p_1, \lambda_1 \rangle \\
 = & \frac{1}{2P^+} \bar{u}(p_2, \lambda_2) \left[H_T^q(x, \xi, t) i\sigma^{+i} + \tilde{H}_T^q(x, \xi, t) \frac{P^+ \Delta^i - \Delta^+ P^i}{M_N^2} \right. \\
 + & \left. E_T^q(x, \xi, t) \frac{\gamma^+ \Delta^i - \Delta^+ \gamma^i}{2M_N} + \tilde{E}_T^q(x, \xi, t) \frac{\gamma^+ P^i - P^+ \gamma^i}{M_N} \right] u(p_1, \lambda_1)
 \end{aligned}$$

- We will consider the simplest case when $\Delta_\perp = 0$.
- In that case and in the forward limit $\xi \rightarrow 0$ **only the H_T^q term survives.**
- Transverse ρ DA at twist 2 :

$$\langle 0 | \bar{u}(0) \sigma^{\mu\nu} u(x) | \rho^0(p, s) \rangle = \frac{i}{\sqrt{2}} (\epsilon_\rho^\mu p^\nu - \epsilon_\rho^\nu p^\mu) f_\rho^\perp \int_0^1 du e^{-iup \cdot x} \phi_\perp(u)$$

Models for DAs

Asymptotical DAs

We take the simplistic asymptotic form of the (normalized) DAs:

$$\phi_{\parallel}(z) = 6z(1-z),$$

$$\phi_{\perp}(z) = 6z(1-z).$$

Model for GPDs: based on the Double Distribution ansatz

Realistic Parametrization of GPDs

- GPDs can be represented in terms of **Double Distributions** [Radyushkin] based on the **Schwinger** representation of a toy model for GPDs which has the structure of a triangle diagram in scalar ϕ^3 theory

$$H^q(x, \xi, t = 0) = \int_{-1}^1 d\beta \int_{-1+|\beta|}^{1-|\beta|} d\alpha \delta(\beta + \xi\alpha - x) f^q(\beta, \alpha)$$

- ansatz for these Double Distributions [Radyushkin]:

- chiral-even sector:

$$f^q(\beta, \alpha, t = 0) = \Pi(\beta, \alpha) q(\beta) \Theta(\beta) - \Pi(-\beta, \alpha) \bar{q}(-\beta) \Theta(-\beta),$$

$$\tilde{f}^q(\beta, \alpha, t = 0) = \Pi(\beta, \alpha) \Delta q(\beta) \Theta(\beta) + \Pi(-\beta, \alpha) \Delta \bar{q}(-\beta) \Theta(-\beta).$$

- chiral-odd sector:

$$f_T^q(\beta, \alpha, t = 0) = \Pi(\beta, \alpha) \delta q(\beta) \Theta(\beta) - \Pi(-\beta, \alpha) \delta \bar{q}(-\beta) \Theta(-\beta),$$

- $\Pi(\beta, \alpha) = \frac{3}{4} \frac{(1-\beta)^2 - \alpha^2}{(1-\beta)^3}$: profile function

- simplistic factorized ansatz for the t -dependence:

$$H^q(x, \xi, t) = H^q(x, \xi, t = 0) \times F_H(t)$$

with $F_H(t) = \frac{C^2}{(t-C)^2}$ a standard **dipole form factor** ($C = .71$ GeV)

Model for GPDs: based on the Double Distribution ansatz



Sets of used PDFs

- $q(x)$: unpolarized PDF [GRV-98]
and [MSTW2008lo, MSTW2008nnlo, ABM11nnlo, CT10nnlo]
- $\Delta q(x)$ polarized PDF [GRSV-2000]
- $\delta q(x)$: transversity PDF [Anselmino *et al.*]

Model for GPDs: based on the Double Distribution ansatz



Typical sets of chiral-even GPDs ($C = -1$ sector)

$$\xi = .1 \leftrightarrow S_{\gamma N} = 20 \text{ GeV}^2 \text{ and } M_{\gamma\rho}^2 = 3.5 \text{ GeV}^2$$



$$H^{q(-)}(x, \xi, t) = H^q(x, \xi, t) + H^q(-x, \xi, t)$$

five Ansätze for $q(x)$: GRV-98, MSTW2008lo, MSTW2008nnlo, ABM11nnlo, CT10nnlo





$$\tilde{H}^{q(-)}(x, \xi, t) = \tilde{H}^q(x, \xi, t) - \tilde{H}^q(-x, \xi, t)$$

“valence” and “standard”: two GRSV Ansätze for $\Delta q(x)$

Model for GPDs: based on the Double Distribution ansatz

Typical sets of chiral-odd GPDs ($C = -1$ sector)

$$\xi = .1 \leftrightarrow S_{\gamma N} = 20 \text{ GeV}^2 \text{ and } M_{\gamma\rho}^2 = 3.5 \text{ GeV}^2$$




$$H_T^{q(-)}(x, \xi, t) = H_T^q(x, \xi, t) + H_T^q(-x, \xi, t)$$

“valence” and “standard”: two GRSV Ansätze for $\Delta q(x)$

⇒ two Ansätze for $\delta q(x)$

Computation of the hard part

20 diagrams to compute



The other half can be deduced by $q \leftrightarrow \bar{q}$ (anti)symmetry

Red diagrams cancel in the chiral-odd case

Final computation

Final computation


$$\mathcal{A} \propto \int_{-1}^1 dx \int_0^1 dz T(x, \xi, z) H(x, \xi, t) \Phi_\rho(z)$$

- One performs the z integration **analytically** using an asymptotic DA $\propto z(1-z)$
- One then plugs our GPD models into the formula and performs the integral w.r.t. x numerically.
- Differential cross section:

$$\left. \frac{d\sigma}{dt du' dM_{\gamma\rho}^2} \right|_{-t=(-t)_{min}} = \frac{|\overline{\mathcal{M}}|^2}{32S_{\gamma N}^2 M_{\gamma\rho}^2 (2\pi)^3}.$$

$|\overline{\mathcal{M}}|^2 =$ averaged amplitude squared

- Kinematical parameters: $S_{\gamma N}^2$, $M_{\gamma\rho}^2$ and $-u'$




Fully differential cross section

Chiral even cross section


at $-t = (-t)_{\min}$

$$\frac{d\sigma_{\text{even}}}{dM_{\gamma\rho}^2 d(-u') d(-t)} \quad (\text{nb} \cdot \text{GeV}^{-6})$$



proton

$$\frac{d\sigma_{\text{even}}}{dM_{\gamma\rho}^2 d(-u') d(-t)} \quad (\text{nb} \cdot \text{GeV}^{-6})$$



neutron

$$S_{\gamma N} = 20 \text{ GeV}^2$$


$$M_{\gamma\rho}^2 = 3, 4, 5, 6 \text{ GeV}^2$$

solid: "valence" model

dotted: "standard" model


Fully differential cross section

Chiral odd cross section

at $-t = (-t)_{\min}$ 

proton

“valence” and “standard” models,
each of them with $\pm 2\sigma$ [S. Melis]



neutron

“valence” model only

$$S_{\gamma N} = 20 \text{ GeV}^2$$

$$M_{\gamma\rho}^2 = 3, 4, 5, 6 \text{ GeV}^2$$







Phase space integration

Evolution of the phase space in $(-t, -u')$ plane

large angle scattering: $M_{\gamma\rho}^2 \sim -u' \sim -t'$

in practice: $-u' > 1 \text{ GeV}^2$ and $-t' > 1 \text{ GeV}^2$ and $(-t)_{\min} \leq -t \leq .5 \text{ GeV}^2$
 this ensures large $M_{\gamma\rho}^2$

example: $S_{\gamma N} = 20 \text{ GeV}^2$



Variation with respect to $S_{\gamma N}$

$$\text{Mapping } (S_{\gamma N}, M_{\gamma\rho}) \mapsto (\tilde{S}_{\gamma N}, \tilde{M}_{\gamma\rho})$$

One can save a lot of CPU time:

- $\mathcal{M}(\alpha, \xi)$ and $GPDs(\xi, x)$
- In the generalized Bjorken limit:
 - $\alpha = \frac{-u'}{M_{\gamma\rho}^2}$
 - $\xi = \frac{M_{\gamma\rho}^2}{2(S_{\gamma N} - M^2) - M_{\gamma\rho}^2}$

Given $S_{\gamma N}$ ($= 20 \text{ GeV}^2$), with its grid in $M_{\gamma\rho}^2$, choose another $\tilde{S}_{\gamma N}$

One can get the corresponding grid in $\tilde{M}_{\gamma\rho}$ by just keeping the same ξ 's:

$$\tilde{M}_{\gamma\rho}^2 = M_{\gamma\rho}^2 \frac{\tilde{S}_{\gamma N} - M^2}{S_{\gamma N} - M^2},$$


From the grid in $-u'$, the new grid in $-\tilde{u}'$ is given by just keeping the same α 's:

$$-\tilde{u}' = \frac{\tilde{M}_{\gamma\rho}^2}{M_{\gamma\rho}^2} (-u').$$


\Rightarrow a single set of numerical computations is required (we take $S_{\gamma N} = 20 \text{ GeV}^2$)

Single differential cross section

Chiral even cross section



proton




neutron

“valence” scenario

$S_{\gamma N}$ vary in the set 8, 10, 12, 14, 16, 18, 20 GeV² (from left to right)

Single differential cross section

Chiral odd cross section




Various ansätze for the PDFs Δq used to build the GPD H_T :

- *dotted curves*: “standard” scenario
- *solid curves*: “valence” scenario
- *deep-blue* and *red* curves: central values
- *light-blue* and *orange*: results with $\pm 2\sigma$.

Single differential cross section

Chiral odd cross section




proton, "valence" scenario


$S_{\gamma N}$ vary in the set 8, 10, 12, 14, 16, 18, 20 GeV² (from left to right)

Integrated cross-section

Chiral even cross section

 σ_{even} (nb)

proton

 σ_{even} (nb)


neutron

solid red: "valence" scenario


dashed blue: "standard" one

Integrated cross-section

Chiral odd cross section

 σ_{odd} (pb)

proton

 σ_{odd} (pb)

neutron

solid red: "valence" scenario

dashed blue: "standard" one


Counting rates for 100 days

example: JLab Hall B

- untagged incoming $\gamma \Rightarrow$ Weizsäcker-Williams distribution
- With an expected luminosity of $\mathcal{L} = 100 \text{ nb}^{-1} \text{ s}^{-1}$, for 100 days of run:
 - Chiral even case : $\simeq 6.8 \cdot 10^6 \rho_L$.
 - Chiral odd case : $\simeq 7.5 \cdot 10^3 \rho_T$


Effects of an experimental angular restriction for the produced γ Angular distribution of the produced γ (chiral-even cross section)

after boosting to the lab frame




$$S_{\gamma N} = 10 \text{ GeV}^2$$

$$M_{\gamma\rho}^2 = 3, 4 \text{ GeV}^2$$



$$S_{\gamma N} = 15 \text{ GeV}^2$$

$$M_{\gamma\rho}^2 = 3, 4, 5 \text{ GeV}^2$$




$$S_{\gamma N} = 20 \text{ GeV}^2$$

$$M_{\gamma\rho}^2 = 3, 4, 5 \text{ GeV}^2$$


JLab Hall B detector equipped between 5° and 35° \Rightarrow this is safe!

Effects of an experimental angular restriction for the produced γ


Angular distribution of the produced γ (chiral-even cross section)



$$S_{\gamma N} = 10 \text{ GeV}^2$$



$$S_{\gamma N} = 15 \text{ GeV}^2$$



$$S_{\gamma N} = 20 \text{ GeV}^2$$

$$\theta_{\text{max}} = 35^\circ, 30^\circ, 25^\circ, 20^\circ, 15^\circ, 10^\circ$$


JLab Hall B detector equipped between 5° and 35°

\Rightarrow this is safe!

Effects of an experimental angular restriction for the produced γ ?


Angular distribution of the produced γ (chiral-odd cross section)

after boosting to the lab frame




$$S_{\gamma N} = 10 \text{ GeV}^2$$

$$M_{\gamma\rho}^2 = 3, 4 \text{ GeV}^2$$



$$S_{\gamma N} = 15 \text{ GeV}^2$$

$$M_{\gamma\rho}^2 = 3.5, 5, 6.5 \text{ GeV}^2$$



$$S_{\gamma N} = 20 \text{ GeV}^2$$


$$M_{\gamma\rho}^2 = 4, 6, 8 \text{ GeV}^2$$

JLab Hall B detector equipped between 5° and 35°


\Rightarrow this is safe!

Effects of an experimental angular restriction for the produced γ


Angular distribution of the produced γ (chiral-even cross section)



$$S_{\gamma N} = 10 \text{ GeV}^2$$



$$S_{\gamma N} = 15 \text{ GeV}^2$$



$$S_{\gamma N} = 20 \text{ GeV}^2$$

$$\theta_{\text{max}} = 35^\circ, 30^\circ, 25^\circ, 20^\circ, 15^\circ, 10^\circ$$

JLab Hall B detector equipped between 5° and 35°

\Rightarrow this is safe!

Conclusion (1)

- High statistics for the chiral-even component: enough to extract H (\tilde{H} ?) and **test the universality of GPDs**
- In this chiral-even sector: analogy with **Timelike Compton Scattering**, the $\gamma\rho$ pair playing the role of the γ^* .
- Strong dominance of the chiral-even component w.r.t. the chiral-odd one:
 - In principle the separation ρ_L/ρ_T can be performed by an angular analysis of its decay products, but this could be very challenging. Cuts in θ_γ might help
 - Future: **study of polarization observables** \Rightarrow sensitive to the interference of these two amplitudes
- The **Bethe Heitler** component (outgoing γ emitted from the incoming lepton) is:
 - zero for the chiral-odd case
 - suppressed for the chiral-even case
- Our result can also be applied to **electroproduction** ($Q^2 \neq 0$) after adding **Bethe-Heitler** contributions and interferences.
- Possible measurement at **JLAB** (Hall B, C, D)
- A similar study could be performed at **COMPASS**. **EIC**, **LHC** in UPC?

Conclusion (2)


Collaboration with Goran Duplančić, Kornelija Passek-Kumerički (IRB, Zagreb), Hervé Moutarde (SPhN), Bernard Pire (CPhT), Lech Szymanowski (NCBJ)

- We are now investigating the process $\gamma N \rightarrow \gamma \pi^{\pm,0} N'$
 - at Born order
 - at one loop
- the processes $\gamma N \rightarrow \gamma \pi^0 N'$ and $\gamma N \rightarrow \gamma \eta^0 N'$ are of particular interest: they give an access to the gluonic GPDs at Born order.

Chiral-even cross section


Contribution of u versus d

$$\frac{d\sigma_{\text{even}}}{dM_{\gamma\rho}^2 d(-u')d(-t)} \quad (\text{nb} \cdot \text{GeV}^{-6})$$



proton

$$\frac{d\sigma_{\text{even}}}{dM_{\gamma\rho}^2 d(-u')d(-t)} \quad (\text{nb} \cdot \text{GeV}^{-6})$$



neutron

$M_{\gamma\rho}^2 = 4 \text{ GeV}^2$. Both vector and axial GPDs are included.

$u + d$ quarks u quark d quark

Solid: "valence" model


dotted: "standard" model

- u -quark contribution dominates due to the charge effect
- the interference between u and d contributions is important and negative.

Chiral-even cross section


Contribution of vector versus axial amplitudes

$$\frac{d\sigma_{\text{even}}}{dM_{\gamma\rho}^2 d(-u') d(-t)} \quad (\text{nb} \cdot \text{GeV}^{-6})$$



proton

$$\frac{d\sigma_{\text{even}}}{dM_{\gamma\rho}^2 d(-u') d(-t)} \quad (\text{nb} \cdot \text{GeV}^{-6})$$



neutron

$M_{\gamma\rho}^2 = 4 \text{ GeV}^2$. Both u and d quark contributions are included.

vector + axial amplitudes / vector amplitude / axial amplitude

solid: "valence" model

dotted: "standard" model

- dominance of the vector GPD contributions
- no interference between the vector and axial amplitudes