# COSMIC ANISOTROPIES FROM QUASARS

# **VINCENT PELGRIMS**

Seminar @ Centre de Physique des Particules de Marseille

Marseille, September 25, 2017

# Past and Current Works

Strong gravitational lensing Master thesis – Liege University, Belgium

Asymptotic solutions for the case of nearly symmetric gravitational lens system
 [0, Wertz, V.P., J. Surdei 2012, MNRAS, 424 1543]

Large-scale alignments of quasar polarization vectors Doctoral thesis – Liege University, Belgium

- A new analysis of quasar polarization alignments
  [V.P., J.R. Cudell 2014, MNRAS, 442 1239]; [V.P., Proc. 2014 IAUS, S306 276]
- Polarization alignments of quasars from JVAS/CLASS 8.4-GHz surveys [V.P., D. Hutsemékers 2015, MNRAS, 450 4161]
- Alignment of quasar polarizations with large-scale structures [D. Hutsemékers, L. Braibant, V.P., D. Sluse 2014, A&A, 472 A18]
- Evidence for the alignment of quasar radio polarizations with large quasar group axes [V.P., D. Hutsemékers 2016, A&A, 590 A53]
- Cosmological-scale coherent orientations of quasar optical polarization vectors in the Planck era – Surviving to Galactic dust contamination scenario [V.P. 2017, A&A submitted]

#### Radio Foregrounds and Galactic Magnetic Field Postdoc – LPSC, Grenoble, France



 Constraints on regular Galactic magnetic field models from 353-GHz polarized sky [V.P., J.F. Macías-Pérez et al. 2017, in preparation]

### **Quasars and Cosmology**


Large-scale alignments of quasar polarization vectors

- Cosmological principle: Isotropy and Homogeneity
- Quasars: general properties and polarization
- Quasars and extreme-scale correlations
- Quasars and large-scale structures

**Cosmic Anisotropies from Quasars** from polarization to structural-axis alignments V.P. 2016, astro-ph: [arXiv:1604.05141]

# **Cosmological Principle**

### The ACDM: successful concordance model of cosmology



Cosmological principle + General Relativity → FLRW Universes [e.g Trodden & Carroll 2004]

Though, some anomalies:

- Low-I deficit in the TT angular power spectrum
- Small temperature variance
- Dipole and quadrupole alignment of moments
- ...
- Departure from isotropic H<sub>0</sub> from SNIa
- Extreme-scale alignments of quasar (optical) polarization vectors

• ...

# **Cosmological Principle**

The Universe has to be **homogeneous** and **isotropic** when it is viewed at *sufficiently* large scale.

It requires/implies that the part of the Universe that we observe and study is a statistically representative sample of its entirety.

- Homogeneity = same observation can be made from wherever
- Isotropy = same observation can be made by looking in whatever direction
  - Isotropy for all observers implies homogeneity
  - Homogeneity for all does not imply isotropy


e.g. Bianchi cosmological models that are homogeneous and anisotropic

Resurgent interests to explain some anomalies such as:

- Low-*I* deficit in the TT angular power spectrum
- Small temperature variance
- Dipole and quadrupole alignment of moments
- ...
- Departure from isotropic H<sub>0</sub> from SNIa
- Extreme-scale alignments of quasar (optical) polarization vectors

• ...

Isotropy appears to be questionable ...



# **Cosmological Principle**

The Universe has to be **homogeneous** and **isotropic** when it is viewed at *sufficiently* large scale.

It requires/implies that the part of the Universe that we observe and study is a statistically representative sample of its entirety.

- Homogeneity = same observation can be made from wherever
- > Isotropy = same observation can be made by looking in whatever direction
  - > Isotropy for all observers implies homogeneity
  - Homogeneity for all does not imply isotropy


e.g. Bianchi cosmological models that are homogeneous and anisotropic

Resurgent interests to explain some anomalies such as:

- Low-*I* deficit in the TT angular power spectrum
- Small temperature variance
- Dipole and quadrupole alignment of moments
- ...
- Departure from isotropic H<sub>0</sub> from SNIa
- Extreme-scale alignments of quasar (optical) polarization vectors

• ...

Isotropy appears to be questionable ... Homogeneity as well, at least the value of the *homogeneity scale* has long been debated and most recently with quasars.




### **Quasars and Cosmology**

Large-scale alignments of quasar polarization vectors

- Cosmological principle: Isotropy and Homogeneity
- Quasars: general properties and polarization
- Quasars and extreme-scale correlations
- Quasars and large-scale structures


**Cosmic Anisotropies from Quasars** from polarization to structural-axis alignments V.P. 2016, astro-ph: [arXiv:1604.05141]


# Polarization of light













# Polarization of light



#### Quasars: some properties


- Most luminous Active Galactic Nuclei
- Ultra-bright point-like sources
- Emit light in the whole spectrum
- Observable at the far reaches of the Universe
- Tiny region at the center of a Galaxy (~ 10<sup>-3</sup> – 10<sup>-4</sup> pc)
   → matter accretion onto a Super Massive Black Hole (>10<sup>8</sup> M<sub>☉</sub>)
- Light is polarized at various wavelengths
   → no spherical symmetry
- Very-high resolution observations of a few showed the optical polarization orientation relates to structural axis of the source (blue/UV continuum or radio jet) [Borguet et al. 2005]



[QSO 1229+204; Hutchings et al. 1994 (HST)]

#### Quasars: some properties


- Most luminous Active Galactic Nuclei
- Ultra-bright point-like sources
- Emit light in the whole spectrum
- Observable at the far reaches of the Universe
- Tiny region at the center of a Galaxy (~ 10<sup>-3</sup> - 10<sup>-4</sup> pc)
   → matter accretion onto a Super Massive Black Hole (>10<sup>8</sup> M<sub>☉</sub>)
- Light is polarized at various wavelengths
   → no spherical symmetry
- Very-high resolution observations of a few showed the optical polarization orientation relates to structural axis of the source (blue/UV continuum or radio jet) [Borguet et al. 2005]



[Urry & Padovani; unified model]

#### Quasars: some properties

- Most luminous Active Galactic Nuclei
- Ultra-bright point-like sources
- Emit light in the whole spectrum
- Observable at the far reaches of the Universe
- Tiny region at the center of a Galaxy (~ 10<sup>-3</sup> - 10<sup>-4</sup> pc)
   → matter accretion onto a Super Massive Black Hole (>10<sup>8</sup> M<sub>☉</sub>)
- Light is polarized at various wavelengths
   → no spherical symmetry
- Very-high resolution observations of a few showed the optical polarization orientation relates to structural axis of the source (blue/UV continuum or radio jet) [Borguet et al. 2005]
- Similar relation at other wavelength




[QSO 1229+204; Hutchings et al. 1994 (HST)]

Extreme-scale alignments of quasar optical polarization vectors

Originally discovered: [Hutsemékers 1998]

Confirmed with:

- new observations [Hutsemékers & Lamy 2001 ; Sluse et al. 2005]
- independent analyses [Hutsemékers & Lamy 2001; Jain et al. 2004; Cabanac et al. 2005; Hutsemékers et al. 2005; Pelgrims & Cudell 2012; Pelgrims 2017]




Probability of uniformity  $\sim 6 \ 10^{-5}$ 

Extreme-scale alignments of quasar optical polarization vectors

#### [Hutsemékers et al. 2005]

- Current sample:
  355 quasars with *reliable* opt. pol.
- Significant orientation correlations within few Gpc scale regions




Probability of uniformity ~  $6 \ 10^{-5}$ 

Extreme-scale alignments of quasar optical polarization vectors

#### [Hutsemékers et al. 2005]


- Current sample:
  355 quasars with *reliable* opt. pol.
- Significant orientation correlations within few Gpc scale regions
- Redshift dependence of preferred orientation



Extreme-scale alignments of quasar optical polarization vectors

#### [Hutsemékers et al. 2005]

- Current sample:
  355 quasars with *reliable* opt. pol.
- Significant orientation correlations within few Gpc scale regions
- Redshift dependence of preferred orientation




Extreme-scale alignments of quasar optical polarization vectors

#### [Hutsemékers et al. 2005]

- Current sample:
  355 quasars with *reliable* opt. pol.
- Significant orientation correlations within few Gpc scale regions
- Redshift dependence of preferred orientation
- Statistically significant inside the whole sample (including LEE)

 $\rightarrow$  probability of randomness <0.1% using dedicated statistical tests



Extreme-scale alignments of quasar optical polarization vectors

#### [Hutsemékers et al. 2005]

- Current sample:
  355 quasars with *reliable* opt. pol.
- Significant orientation correlations within few Gpc scale regions
- Redshift dependence of preferred orientation
- Statistically significant inside the whole sample (including LEE)

 $\rightarrow$  probability of randomness <0.1% using dedicated statistical tests

#### Still not understood today !

#### No satisfactory explanation despite the various investigated scenarios

- Cosmic strings/loops
- Cosmological-scale magnetic field
- Axion-like Dark Matter particle
- Birefringence of the Universe
- Anisotropic cosmological expansion
- ...

#### [V.P. & Cudell 2014 ; V.P. 2014]


- Confirmation of alignments with new and statistically independent methods
- Confirmation of redshift dependence but with no smooth and continuous rotation as suggested before

#### [V.P. 2017]

 Robustness of alignments regarding interstellar polarization contamination evaluated from *Planck* map

Extreme-scale alignments of quasar optical polarization vectors


What can cause the polarization alignments ?



- Photon path effects
  - Modulation of the polarization state
  - Asymptotic rotation of the polarization vectors
- Structural axis alignment


Extreme-scale alignments of quasar optical polarization vectors

What can cause the polarization alignments ?



- Photon path effects
  - Modulation of the polarization state
  - Asymptotic rotation of the polarization vectors
- Structural axis alignment

Extreme-scale alignments of quasar optical polarization vectors




What can cause the polarization alignments ?

- Photon path effects
  - Modulation of the polarization state
  - Asymptotic rotation of the polarization vectors
- Structural axis alignment

Extreme-scale alignments of quasar optical polarization vectors

What can cause the polarization alignments ?



- Photon path effects
  - Modulation of the polarization state
  - Asymptotic rotation of the polarization vectors
- Structural axis alignment

These scenario have different observational signatures

Wavelength dependence of the alignments ?


#### Quasar polarization alignments in JVAS/CLASS 8.4 GHz surveys

Based on JVAS/CLASS 8.4-GHz surveys [Jackson et al. 2007]

- Situation unclear from previous studies
  [Joshi et al. 2007 ; Tiwari & Jain 2013 ; Shurtleff 2014]
- Lack of consideration of the intrinsic properties of the sources (redshift, type, ...)

#### [V.P. & Hutsemékers 2015]

- > Clear identification of 4155 Flat Spectrum Radio Sources with reliable polarization measurements  $(f_{pol} > 1 \text{ mJy}; \sigma_{\psi} \le 14^{\circ})$
- Nasa Extragalactic Database
  - $\rightarrow$  redshift for 1531 sources
  - $\rightarrow$  Classification in Object Types




Quasar polarization alignments in JVAS/CLASS 8.4 GHz surveys

#### [V.P. & Hutsemékers 2015]

- > Evidence for alignment in one of the region of optical pol. alignment (~ $3\sigma$ )
- Stat. significant alignment features within the whole sample

Dedicated global statistical tests:

- comparison of polarizations in groups of nearest neighbors and averaged with the whole sample
- 10<sup>4</sup> Monte Carlo simulations for random distribution
- For any given size of groups of neighboring sources



Quasar polarization alignments in JVAS/CLASS 8.4 GHz surveys


10<sup>0</sup>

#### [V.P. & Hutsemékers 2015]

- > Evidence for alignment in one of the region of optical pol. alignment (~ $3\sigma$ )
- Stat. significant alignment features within the whole sample

Dedicated global statistical tests:

- comparison of polarizations in groups of nearest neighbors and averaged with the whole sample
- 10<sup>4</sup> Monte Carlo simulations for random distribution
- For any given size of groups of neighboring sources



- ightarrow 2D analysis with no restriction on the redshift
- $\rightarrow$  For a wide range of size of groups of neighboring sources
- $\rightarrow$  For all subsamples at hand

Quasar polarization alignments in JVAS/CLASS 8.4 GHz surveys


#### [V.P. & Hutsemékers 2015]

- > Evidence for alignment in one of the region of optical pol. alignment (~ $3\sigma$ )
- Stat. significant alignment features within the whole sample


Only for quasars!

Identification of aligned groups

clustered towards regions where quasar polarization vectors are aligned at optical wavelengths!




Quasar polarization alignments in JVAS/CLASS 8.4 GHz surveys



#### Radio wavelengths

- Photon path effects
  - Modulation of the polarization state
  - Asymptotic rotation of the polarization vectors
- Structural axis alignment

Quasar polarization alignments in JVAS/CLASS 8.4 GHz surveys




### **Quasars and Cosmology**


Large-scale alignments of quasar polarization vectors

- Cosmological principle: Isotropy and Homogeneity
- Quasars: general properties and polarization
- Quasars and extreme-scale correlations
- Quasars and large-scale structures

**Cosmic Anisotropies from Quasars** from polarization to structural-axis alignments V.P. 2016, astro-ph: [arXiv:1604.05141]



[Clowes et al. 2013] → discovery of a big inhomogeneity in the quasar distribution the Huge-LQG, next to the CCLQG



#### Huge-LQG


- ≻ z ~ 1.3
- > 73 quasars
- elongation ~ 1 Gpc !

*much* bigger than the homogeneity scale of the Universe ...

A problem that has finally been solved:

[see: Nadathur 2013 ; Einasto et al. 2014 ; Parkes et al. 2015 and finally [Marinello et al. 2016]


[Clowes et al. 2013]  $\rightarrow$  discovery of a big inhomogeneity in the quasar distribution the Huge-LQG, next to the CCLQG



The Huge-LQG (and the CCLQG) is at the outskirt (3D) of one of the regions of optical polarization alignments of quasars

[Hutsemékers, Braibant, V.P., Sluse 2014]


Polarization in the Huge-LQG and CCLQG





[Hutsemékers, Braibant, V.P., Sluse 2014]


Polarization in the Huge-LQG and CCLQG




- ≻ z ~ 1.3
- > 73 + 20 observed quasars
- > 19 with  $p_{\text{lin}} \ge 0.6\%$

[Hutsemékers, Braibant, V.P., Sluse 2014]

Polarization in the Huge-LQG and CCLQG






# Quasars and optical polarization

Within Quasar Unification Scheme [e.g. Antonucci 1993; Urry & Padovani 1995] observables depend on the inclination of the system w.r.t. the line of sight


- Optical polarization result from two competing components  $\rightarrow$  either parallel or perpendicular to quasar morphological axis [e.g. Smith et al. 2004 ; Borget et al. 2008]
- Width low-ionization emission lines depends on inclination [e.g. Wills & Brown 1986; Brotherton 1996; Jarvis & McLure 2006]



Large emission line width of MgII


[Hutsemékers, Braibant, V.P., Sluse 2014]

Polarization in the Huge-LQG and CCLQG




[Hutsemékers, Braibant, V.P., Sluse 2014]

Polarization in the Huge-LQG and CCLQG




[Hutsemékers, Braibant, V.P., Sluse 2014]

Polarization in the Huge-LQG and CCLQG



#### [V.P. & Hutsemékers 2016]

Radio polarization in a large LQG sample




- > SDSS DR7 : 22.381 quasars
  - with 1.0 < z < 1.8 ;  $i_{mag}$  < 19.1
- Large sample of large quasar

groups by [Einasto et al. 2014]

#### [V.P. & Hutsemékers 2016]

Radio polarization in a large LQG sample




- > SDSS DR7 : 22.381 quasars
  - with 1.0 < z < 1.8;  $i_{mag} < 19.1$
- Large sample of large quasar

groups by [Einasto et al. 2014]

[V.P. & Hutsemékers 2016]


Radio polarization in a large LQG sample



- SDSS DR7 : 22.381 quasars
  - with 1.0 < z < 1.8 ;  $i_{mag}$  < 19.1
- Large sample of large quasar groups by [Einasto et al. 2014]
- Radio polarization from JVAS/CLASS 8.4-GHz surveys
   [Jackson et al. 2007]
- > 19 quasars with pol. in LQG having > 20 members

[V.P. & Hutsemékers 2016]


Radio polarization in a large LQG sample



- SDSS DR7 : 22.381 quasars
  - with 1.0 < z < 1.8 ;  $i_{mag}$  < 19.1
- Large sample of large quasar groups by [Einasto et al. 2014]
- Radio polarization from JVAS/CLASS 8.4-GHz surveys
   [Jackson et al. 2007]
- > 19 quasars with pol. in LQG having > 20 members
- LQG orientations from inertia tensors

#### [V.P. & Hutsemékers 2016]

- Radio polarization in a large LQG sample
- > Polarization (synchrotron) is preferentially ⊥ to quasar spin axis [Joshi et al. 2007]




Quasar spin-axes preferentially parallel to the major axes of rich large quasar groups at *high redshifts* and *over large scales* !

[Hutsemékers, Braibant, V.P., Sluse 2014; V.P. & Hutsemékers 2016]

- Made use of optical and radio polarization of quasars to infer their spin axes at high redshift
- Show quasar spin-axes correlate to the major axes of their host LQG


Corroborated by

- > degree scale radio-jet axis correlations
  - [Taylor & Jagannathan 2016]
    → alignments in 1.4 deg<sup>2</sup> ELAIS N1 field
  - [Contigiani et al. 2017]
    - $\rightarrow$  alignments at scale 1.5–2.5 deg in 7000 deg<sup>2</sup> FIRST+RadioGalaxyZoo sample (30 059 sources)
- > degree scale radio-polarization correlations
  - · [V.P. & Hutsemékers 2015]
    - $\rightarrow$  alignments < 5 deg found in JVAS/CLASS 8.4GHz



#### Explained through coevolution of galaxy spin axes within the cosmic web?

Involved scales seem too large ...



[Artist view of the "spooky" alignment Credit: ESO/M. Kornmesser]

#### Take away

#### Quasar polarization alignments

There are evidences for extreme-scale alignments of the polarization of quasars when measured at optical and at radio wavelengths

- Origin is still to be found
- Could indicate departure from isotropy of the Universe given the characteristic size of the correlation
- > Difference between optical and radio signatures needs to be clarified

The large-scale correlation of quasar spin axes with and within large quasar groups

- Could be due to coevolution of black hole spins in LSS
- > Typical size involved are way larger than expected

 $\rightarrow$  large-scale intrinsic alignments of galaxies ???

If true...

the two types of alignments could find the same origin assuming anisotropy in large-scale structure orientations

Thank you

# Васкир