$pp \to ZH \to \nu\bar{\nu}b\bar{b}$, @14 TeV, $m_H = 115$ GeV

SIGNAL

Herwig 6.510 + Jimmy 4.31 + FastJet 2.3



Zbb BACKGROUND

Cluster event, C/A, R=1.2

Butterworth, Davison, Rubin & GPS '08

arbitrary norm.

$pp \to ZH \to \nu\bar{\nu}b\bar{b}$, @14 TeV, $m_H = 115$ GeV

SIGNAL

Herwig 6.510 + Jimmy 4.31 + FastJet 2.3



Zbb BACKGROUND

Fill it in, $\to$ show jets more clearly

Butterworth, Davison, Rubin & GPS '08

arbitrary norm.

$pp \to ZH \to \nu\bar{\nu}b\bar{b}$, @14 TeV, $m_H = 115$ GeV

Herwig 6.510 + Jimmy 4.31 + FastJet 2.3



SIGNAL



Zbb BACKGROUND



Consider hardest jet, $m = 150$ GeV

Butterworth, Davison, Rubin & GPS '08

arbitrary norm.

$pp \to ZH \to \nu\bar{\nu}b\bar{b}$, @14 TeV, $m_H = 115$ GeV

Herwig 6.510 + Jimmy 4.31 + FastJet 2.3



SIGNAL



Zbb BACKGROUND



split: $m = 150$ GeV, $\frac{\max(m_1, m2)}{m} = 0.92 \to$ repeat

Butterworth, Davison, Rubin & GPS '08

arbitrary norm.

$pp \to ZH \to \nu\bar{\nu}b\bar{b}$, @14 TeV, $m_H = 115$ GeV

Herwig 6.510 + Jimmy 4.31 + FastJet 2.3



SIGNAL



Zbb BACKGROUND



split: $m = 139$ GeV, $\frac{\max(m_1, m2)}{m} = 0.37 \to$ mass drop

Butterworth, Davison, Rubin & GPS '08

arbitrary norm.

$pp \to ZH \to \nu\bar{\nu}b\bar{b}$, @14 TeV, $m_H = 115$ GeV

Herwig 6.510 + Jimmy 4.31 + FastJet 2.3



SIGNAL



Zbb BACKGROUND



check: $y_{12} \simeq \frac{p_{t2}}{p_{t1}} \simeq 0.7 \to$ OK + 2 b-tags (anti-QCD)

Butterworth, Davison, Rubin & GPS '08

arbitrary norm.

$pp \to ZH \to \nu\bar{\nu}b\bar{b}$, @14 TeV, $m_H = 115$ GeV

Herwig 6.510 + Jimmy 4.31 + FastJet 2.3



SIGNAL



Zbb BACKGROUND



$R_{filt} = 0.3$

Butterworth, Davison, Rubin & GPS '08

arbitrary norm.

$pp \to ZH \to \nu\bar{\nu}b\bar{b}$, @14 TeV, $m_H = 115$ GeV

Herwig 6.510 + Jimmy 4.31 + FastJet 2.3



SIGNAL



Zbb BACKGROUND



$R_{filt} = 0.3$: take 3 hardest, **m = 117 GeV**

Butterworth, Davison, Rubin & GPS '08

arbitrary norm.

$pp \to ZH \to \nu\bar{\nu}b\bar{b}$, @14 TeV, $m_H = 115$ GeV

Herwig 6.510 + Jimmy 4.31 + FastJet 2.3





The declustering and cuts do two things

- **sharpen signals**
- **reduce backgrounds**

$R_{filt} = 0.3$: take 3 hardest, **m =**

Butterworth, Davison, Rubin & GPS '08

arbitrary norm.