Workshop on Active Targets and Time Projection Chambers for High-intensity and Heavy-ion beams in Nuclear Physics

ID de Contribution: 27

Type: Non spécifié

Study of giant resonances with active targets

jeudi 18 janvier 2018 10:20 (20 minutes)

The presence of coherent motions of particles in many-body systems, i.e. collective motions, is a common feature in several branches of physics. In atomic nuclei, a particular case of nuclear collective motion is represented by the giant resonances (GR) [1], which are the subject of this presentation. These resonance states play a key role in the understanding of the nuclear structure because of their connection with the bulk properties of atomic nuclei.

Giant resonances can be macroscopically viewed as a quantum oscillation of two fermionic liquids (neutron and protons) involving spatial (L), spin (S) and isospin (I) degree of freedom. In the case of an isoscalar oscillation (Δ I=0) neutron and protons move together in phase. On the other hand, in an isovector oscillation (Δ I =1) neutron and protons move in opposite direction.

The isoscalar giant monopole resonance (ISGMR) measures the collective response of the nucleus to density fluctuations (Δ I, Δ S, Δ L=0) [1]. The ISGMR is particularly interesting for its connection with the incom- pressibility of the nucleus KA, which, in turn, can be linked to the incompressibility of nuclear matter K ∞ , an important ingredient of the nuclear-matter equation-of-state (EOS). The EOS, essentially, describes the binding energy per nucleon as a function of nuclear density and it plays an important role in the description of heavy-ion nuclear collision, the collapse of the heavy stars in super novae explosion and the description of neutron stars [2]. In order to improve the understanding of this nuclear mechanism, new experimental data in unstable nuclei far from the stability are needed [3].

The reaction mechanism used to excite the ISGMR is the inelastic scattering of the nuclei of interest on an hadron isosclar probe, typically an α particle. The use of an active target coupled with silicon detectors allows to measure the α particles at forward angles (where the maximum of the cross section is located) and with a very small kinetic energy [4].

In addition, the (α, α') reaction can be also used to excite isoscalar dipole states (L=1, I=0) around the neutron separation energy [5]. These states, also called pygmy dipole resonance (PDR), are of great interest for the impact on astrophysical phenomena, such as r-process nucleosynthesis [6]. The nature of the PDR is largely debated. SpecMAT [7], an active target placed in a high magnetic field and coupled with scintillation detectors, will be a powerful detector to observe PDR in unstable nuclei.

In this presentation the use of active targets to study ISGMR and PDR will be shown.

[1] M. N. Harakeh and A. van der Woude, Giant Resonances, Fundamental High-Frequency Modes of Nuclear Excitation, Oxford Science Publications, 2001.

[2] J.M. Lattimer and M. Prakash, Astrophys. J. 550, 426 (2001) and Science 304, 5670 (2004).

[3] E. Khan, J. Margueron and I. Vidana, Phys. Rev. Lett. 109, 092501 (2012).

[4] M. Vandebrouck, et al., Phys. Rev. Lett. 113, 032504 (2014).

[5] D. Savran, T. Aumann, and A. Zilges, Prog. Part. Nucl. Phys. 70, 210 (2013). [6] S. Goriely, Phys. Lett. B 436, 10 (1998).

[7] R. Raabe, SpecMAT ERC Consolidator Grant (2014).

Authors: Dr CERUTI, Simone (KU Leuven); M. RAJ, Alex (KU Leuven); RAABE, Riccardo (KU Leuven); Dr DE WITTE, Hilde (KU Leuven); POLESHCHUK, Oleksii (KU Leuven, Institute for Nuclear and Radiation Physics); M. RENAUD, Maxim (KU Leuven); M. YANG, Jiecheng (KU Leuven)

Orateur: Dr CERUTI, Simone (KU Leuven)