


# Very rare decays at LHC

**Moriond EW 2018**

La Thuile, 10-17 March, 2018



Maarten van Veghel<sup>1</sup> (Nikhef, the Netherlands)  
*on behalf of the ATLAS, CMS and LHCb<sup>1</sup> collaborations*



# Why look for very rare decays?

## Rare decays

- Small Standard Model (SM) branching fractions  $\rightarrow$  relatively large hypothetical **new physics** (NP) contributions
- Sensitive to higher energy than direct searches due to virtual contributions
- Often Flavour Changing Neutral Currents (FCNCs), which are not possible at tree level in SM
- **Anomalies** seen in  $b \rightarrow s \ell^+ \ell^-$  FCNCs
  - ▶ See e.g. F. Dettori's and C. Langenbruch's talks for  $b \rightarrow s \ell^+ \ell^-$


## Forbidden decays

- If you see SM forbidden decays, obviously new physics

*Note: will only discuss a selection of recent results*

# $B_{(s)}^0 \rightarrow \mu^+ \mu^-$ : the flagship of very rare decays


## Standard Model



- $b \rightarrow s l^+ l^-$  and  $b \rightarrow d l^+ l^-$  like FCNC
- Both loop diagrams and helicity suppression, so **heavily suppressed**
  - ▶ In SM<sup>a</sup>:  $\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) \approx 4 \times 10^{-9}$   
and  $\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) \approx 1 \times 10^{-10}$
- **Theoretically clean** and experimentally favourable

## New physics scenarios

- New particles in loops
- Sensitive to (pseudo)scalars (no helicity suppression)
- Specific: e.g. certain **SUSY** models, multiple Higgs models





<sup>a</sup>[Phys. Rev. Lett. 112, 101801 (2014)]

# $B_{(s)}^0 \rightarrow \mu^+ \mu^-$ : CMS+LHCb, [Nature 522, 68-72 (04 June 2015)]

- **First observation** of  $B_s^0 \rightarrow \mu^+ \mu^-$  with CMS+LHCb **combined analysis** from full LHC Run 1 dataset published in 2015
- First evidence of  $B^0 \rightarrow \mu^+ \mu^-$  (slight enhancement from SM)
- Although **compatible** with SM

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = 2.8_{-0.6}^{+0.7} \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) = 3.9_{-1.4}^{+1.6} \times 10^{-10}$$




# $B_s^0 \rightarrow \mu^+ \mu^-$ : ATLAS, [Eur.Phys.J. C76 (2016) no.9, 513]

- Data from full Run 1 dataset
- No significant signal, but **consistent** with SM with  $2.0\sigma$
- Result in **agreement** with CMS+LHCb results


$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = 0.9_{-0.8}^{+1.1} \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 4.2 \times 10^{-10} \text{ [95\% CL]}$$

## High BDT<sup>1</sup> region



## Limits




<sup>1</sup>[Multivariate classifier (Boosted Decision Tree)]

# $B_{(s)}^0 \rightarrow \mu^+ \mu^-$ : LHCb, update [Phys. Rev. Lett. 118, 191801 (2017)]

## Updated $\mathcal{B}$ measurement

- Run 1 + part Run 2 (2015/2016;  $1.4 \text{ fb}^{-1}$ ) data
- Improved BDT and  $B \rightarrow h^+ h^-$  rejection
- Result **compatible** with SM
- $B^0 \rightarrow \mu^+ \mu^-$  fluctuated down

## High BDT region




$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (3.0 \pm 0.6^{+0.3}_{-0.2}) \times 10^{-9}$$

## Effective lifetime measurement

- Possible **NP orthogonal to  $\mathcal{B}$**  contribution in effective lifetime
- Disentangle CP-odd versus CP-even contribution
- First measurement, as proof of principle, compatible with SM

## Decay time distribution



$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 3.4 \times 10^{-10} \text{ [95\% CL]}$$

$$B_{(s)}^0 \rightarrow \tau^+ \tau^-$$

- **Lepton Flavour Universality tensions** hint to new physics with  $\tau$ 's
- Similar to  $B_{(s)}^0 \rightarrow \mu^+ \mu^-$ , but **less helicity suppressed**
  - ▶ In SM<sup>1</sup>:  $\mathcal{B}(B_s^0 \rightarrow \tau^+ \tau^-) \approx 8 \times 10^{-7}$  and  $\mathcal{B}(B^0 \rightarrow \tau^+ \tau^-) \approx 2 \times 10^{-8}$
- Experimentally much more **difficult**
  - ▶ Neutrinos in final state

**Previous limit** from BaBar [Phys. Rev. Lett. 96, 241802 (2006)]

$$\mathcal{B}(B^0 \rightarrow \tau^+ \tau^-) < 4.1 \times 10^{-3} \text{ [90\% CL]}$$


---

<sup>1</sup>[Phys. Rev. Lett. 112, 101801 (2014)]

$B_{(s)}^0 \rightarrow \tau^+ \tau^-$ : LHCb, [Phys. Rev. Lett. 118, 251802 (2017)]

## Neural net for signal region

- Run 1 data
- $\tau$  reconstruction with  $\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \nu$
- No clear mass peak due to neutrinos, **fit neural net output<sup>a</sup>**
- Different Dalitz regions of  $\tau$  decay and control decay channels to determine/verify neural net response



## New best limits

$$\mathcal{B}(B^0 \rightarrow \tau^+ \tau^-) < 2.1 \times 10^{-3} \text{ [95\% CL]}$$

$$\mathcal{B}(B_s^0 \rightarrow \tau^+ \tau^-) < 6.8 \times 10^{-3} \text{ [95\% CL]}$$


<sup>a</sup>[Neural net used as multivariate classifier]

$B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ : forbidden lepton flavour violating decays

*Lepton flavour violation linked to lepton universality<sup>2</sup>*

- SM has no charged lepton flavour violation
- For example **leptoquark** model could **explain** seen **tensions**
- Opens up possibilities for lepton flavour violation (like  $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ )
- Previous limits for  $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$  at about  $10^{-8}$  to  $10^{-9}$

$$B_s^0 \rightarrow \ell'^+ \ell^- \quad (\text{like } B_{(s)}^0 \rightarrow e^\pm \mu^\mp)$$




<sup>2</sup>[Phys.Rev.Lett. 114 (2015) 091801]

# $B_{(s)}^0 \rightarrow e^\pm \mu^\mp$ : LHCb, [LHCb-PAPER-2017-031] (accepted by JHEP)

- Previous best limit set by LHCb with 2011 data [Phys.Rev.Lett. 111 (2013) 141801]
- **New limits** factor **2 to 3 improvement** on limit

## High BDT region



## Improvements

- Increased BDT performance
- Higher trigger efficiency
- Full Run 1 dataset

## Limits

$$\mathcal{B}(B^0 \rightarrow e^\pm \mu^\mp) < 1.3 \times 10^{-9} \text{ [95\% CL]}$$

$$\mathcal{B}(B_s^0 \rightarrow e^\pm \mu^\mp) < 6.3 \times 10^{-9} \text{ [95\% CL]}$$

$$K_S^0 \rightarrow \mu^+ \mu^-$$


- Similar to  $B_{(s)}^0 \rightarrow \mu^+ \mu^-$ : also a FCNC, but  $s \rightarrow d\ell^+\ell^-$
- $K_S^0 \rightarrow \mu^+ \mu^-$  requires  $CP$  violation for s-wave SM component, which is small, so more suppressed, at level  $5 \times 10^{-12}$  in SM
- Model-independent bounds on  $CP$  violation phase of  $s \rightarrow d\ell^+\ell^-$

$$K_L^0 \rightarrow \mu^+ \mu^-$$

- Observed:  $(6.84 \pm 0.11) \times 10^{-9}$  [PDG 2017]
- Excellent agreement with SM:  $(6.85 \pm 0.32) \times 10^{-9}$  [arXiv:hep-ph/9411439]

## Example of New Physics range


[arXiv:1711.11030]




# $K_S^0 \rightarrow \mu^+ \mu^-$ : LHCb, Eur. Phys. J. C, 77 10 (2017) 678

- Challenge for LHCb: high backgrounds and low trigger efficiencies
- **Improved trigger and more data**: full Run 1
- Main background:  $K_S^0 \rightarrow \pi^+ \pi^-$
- Specific BDTs for muon ID and against combinatorial

$$K_S^0 \rightarrow \pi^+ \pi^-$$



## Part signal region (high BDT bin)



$$\mathcal{B}(K_S^0 \rightarrow \mu^+ \mu^-) < 1.0 \times 10^{-9} \text{ [95\% CL]}$$



- **Improvement of factor 11** from previous result on limit

$$\Sigma^+ \rightarrow p\mu^+\mu^-$$

- Similar  $s \rightarrow d\ell^+\ell^-$  transition as  $K_S^0 \rightarrow \mu^+\mu^-$
- SM  $\mathcal{B}$  prediction higher, due to long-distance contributions, at  $10^{-8}$  to  $10^{-7}$
- Non-resonant SM  $\mathcal{B}$  at about  $10^{-12}$

## HyperCP

- Evidence from HyperCP experiment [Phys.Rev.Lett.94:021801,2005]
- **All 3 events at dimuon mass of  $\approx 214 \text{ MeV}/c^2$**



$$\mathcal{B}(\Sigma^+ \rightarrow pX^0(\rightarrow \mu^+\mu^-)) = (3.1_{-1.9}^{+2.4} \pm 1.5) \times 10^{-8}$$




# $\Sigma^+ \rightarrow p\mu^+\mu^-$ : LHCb, [LHCb-PAPER-2017-049] (submitted to PRL)

- Full Run 1 dataset
- **Evidence** for the decay! ( $4.0\sigma$ )
- Consistent with SM
- **No visible structure in dimuon mass**

$$\mathcal{B}(\Sigma^+ \rightarrow p\mu^+\mu^-) = 2.1_{-1.2}^{+1.6} \times 10^{-8}$$


## Limit on resonance at $214 \text{ MeV}/c^2$

$$\mathcal{B}(\Sigma^+ \rightarrow pX^0(\rightarrow \mu^+\mu^-)) < 1.2 \times 10^{-8} \text{ [95\% CL]}$$




$$\Lambda_c^+ \rightarrow p \mu^+ \mu^- \text{ and } D^0 \rightarrow h^+ h^- \mu^+ \mu^-$$

- $c \rightarrow u \ell^+ \ell^-$  FCNC
- Both  $\Lambda_c^+ \rightarrow p \mu^+ \mu^-$  and  $D^0 \rightarrow h^+ h^- \mu^+ \mu^-$  have this transition
- SM predicts both at around  $10^{-9}$  for non-resonant contribution




$\Lambda_c^+ \rightarrow p\mu^+\mu^-$ : LHCb, [LHCb-PAPER-2017-039] (submitted to PRL)

$\Lambda_c^+ \rightarrow p\phi(\rightarrow \mu^+\mu^-)$



Dimuon mass



- Previous limit at about  $4 \times 10^{-5}$  [90% CL] from BaBar

**This analysis**

- Full Run 1 dataset
- Contributions seen from  $\phi$  and  $\omega$
- No signal seen in non-resonant region
- Near  **$10^3$  improvement** in limit!


**Limit on non-resonant contribution**

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) < 9.6 \times 10^{-8} \text{ [95\% CL]}$$


# $D^0 \rightarrow h^+ h^- \mu^+ \mu^-$ : LHCb, [Phys. Rev. Lett. 119, 181805 (2017)]

- Part Run 1 (2012, 2 fb<sup>-1</sup>) dataset
- **First observation!** of both decays
- Rarest charm decay so far!
- Both in agreement with SM (long-distance contributions)

$$D^0 \rightarrow K^+ K^- \mu^+ \mu^-$$



$$D^0 \rightarrow \pi^+ \pi^- \mu^+ \mu^-$$



$$\mathcal{B}(D^0 \rightarrow \pi^+ \pi^- \mu^+ \mu^-) = (9.64 \pm 0.48(\text{stat.}) \pm 0.51(\text{syst.}) \pm 0.97(\text{norm.})) \times 10^{-7}$$

$$\mathcal{B}(D^0 \rightarrow K^+ K^- \mu^+ \mu^-) = (1.54 \pm 0.27(\text{stat.}) \pm 0.09(\text{syst.}) \pm 0.16(\text{norm.})) \times 10^{-7}$$

# Summary

- Very rare decays **good probe for new physics**
- Rich field, many results not shown
- Many improved limits and observations of decays with suppressed FCNC  $b \rightarrow sl^+l^-$ ,  $b \rightarrow dl^+l^-$ ,  $s \rightarrow dl^+l^-$  and  $c \rightarrow ul^+l^-$  transitions
- Improved limits on lepton flavour violating channels
- Help to **constrain possible new physics in B-anomalies**
- Stay tuned for (Run 2) updates and new analyses!