



ID de Contribution: 29

Type: **Orale**

Light superfluidity in hot atomic vapors

mardi 9 juillet 2019 09:15 (15 minutes)

A laser field propagating through a hot atomic vapor leads to a third-order nonlinear Kerr susceptibility. In the paraxial approximation the evolution of the transverse electric field is described by a 2D Gross-Pitaevskii equation. As such, this system is a promising platform to study phenomena related to Bose-Einstein condensation and superfluidity of light.

In this talk, I will report on the measurement of the dispersion relation of small amplitude density waves propagating on top of a photon fluid. We find a dispersion relation of Bogoliubov type: linear at small wave vector as expected in the superfluid regime and “particle-like” (quadratic) at larger wave vectors ¹. In the superfluid regime, we characterize the dependence of the sound velocity with intensity (photon density) and compare our results with theoretical predictions.

¹ Observation of the Bogoliubov Dispersion in a Fluid of Light, Q. Fontaine, T. Bienaimé, S. Pigeon, E. Giacobino, A. Bramati, and Q. Glorieux, Phys. Rev. Lett. 121, 183604 (2018).

Indico rendering error

Could not include image: [404] Error fetching image

Choix de session parallèle

2.3 Fluides quantiques et lumière

Auteurs principaux: Dr BIENAIMÉ, Tom (Laboratoire Kastler Brossel); M. FONTAINE, Quentin (Laboratoire Kastler Brossel); Dr PIGEON, Simon (Laboratoire Kastler Brossel); Prof. GIACOBINO, Élisabeth (Laboratoire Kastler Brossel); Prof. BRAMATI, Alberto (Laboratoire Kastler Brossel); Prof. GLORIEUX, Quentin (Laboratoire Kastler Brossel)

Orateur: Dr BIENAIMÉ, Tom (Laboratoire Kastler Brossel)

Classification de Session: Séance Parallèle