

# A new light on rare B decays with holographic light-front wavefunctions

Mohammad Ahmady

Department of Physics  
Mount Allison University

LIO International Conference on Flavour Physics: From Flavour to New Physics

April 19, 2018

- 1 Holographic Light-Front Wavefunction
- 2 vector mesons
- 3 Light cone distribution amplitudes
- 4 Light cone DAs in rare B decays
- 5  $B \rightarrow V$  transition form factors
- 6 Summary and outlook

# Holographic Schrödinger equation

An important equation in light-front holographic QCD is the holographic Schrödinger equation (HSE) for mesons:

$$\left( -\frac{d^2}{d\zeta^2} - \frac{1-4L^2}{4\zeta^2} + U_{\text{eff}}(\zeta) \right) \phi(\zeta) = M^2 \phi(\zeta)$$

- Derived within a semiclassical approximation of light-front QCD, where quantum loops and quark masses are neglected.
- The holographic variable  $\zeta = \sqrt{x\bar{x}b}$  with  $\bar{x} \equiv 1 - x$  where  $b$  is the transverse separation of the quark and antiquark and  $x$  is the light-front momentum fraction carried by the quark.
- $M$  is the meson mass.

G. F. de Teramond and S. J. Brodsky,  
PRL94,201601(2005),PRL96,201601(2006),PRL102,081601(2009)

## 3 mechanisms to break conformal symmetry

- 1 Spontaneous
- 2 Explicit
- 3 de Alfaro, Fubini, Furlan (dAFF) ✓ V. Alfaro, S. Fubini and G. Furlan. Nuovo. Cim. A34 (1976) 569

in conformal QM, changing the evolution parameter allows the introduction of a mass scale in the Hamiltonian while preserving the conformal invariance of the underlying action.

- 4 dAFF/LF mapping  $\Rightarrow$  Semiclassical QCD LF Hamiltonian

$$H_{\text{scQCD}}^{\text{LF}} = \left( -\frac{d^2}{d\zeta^2} - \frac{1 - 4L^2}{4\zeta^2} + U(\zeta) \right)$$

with  $U(\zeta) = \kappa^4 \zeta^2$

## Confining potential $U_{\text{eff}}$

A remarkable feature in light-front holography is that the form of the confinement potential is uniquely determined to be that of a harmonic oscillator, i.e.  $U_{\text{eff}} = \kappa^4 \zeta^2$  where  $\kappa$  is the fundamental of the model.

S. J. Brodsky, G. F. De Tramond, and H. G. Dosch, *Phys. Lett. B* **729**, 3 (2014), 1302.4105.

$\zeta \rightarrow z$  (the fifth dimension of anti-de Sitter (AdS) space), the HSE also described the propagation of weakly-coupled spin- $J$  modes in a modified AdS space with the confining QCD potential then determined by the form of the dilaton field,  $\varphi(z)$ , which modifies the pure AdS geometry.

$$U_{\text{eff}}(\zeta) = \frac{1}{2}\varphi''(z) + \frac{1}{4}\varphi'(z)^2 + \frac{2J-3}{2z}\varphi'(z)$$

To recover this harmonic potential, the dilaton field has to be quadratic, i.e.  $\varphi(z) = \kappa^2 z^2$

$$U_{\text{eff}}(\zeta) = \kappa^4 \zeta^2 + 2\kappa^2(J-1)$$

where  $J = L + S$ .

# Eigenvalues and eigenfunctions

Solving the holographic LF Schrödinger Equation

$$\left( -\frac{d^2}{d\zeta^2} - \frac{1 - 4L^2}{4\zeta^2} + U(\zeta) \right) \phi(\zeta) = M^2 \phi(\zeta)$$

with

$$U(\zeta) = \kappa^4 \zeta^2 + 2\kappa^2(J - 1)$$

gives

$$M^2 = (4n + 2L + 2)\kappa^2 + 2\kappa^2(J - 1) = 4\kappa^2\left(n + L + \frac{S}{2}\right)$$

$$\phi_{n,L}(\zeta) = \kappa^{1+L} \sqrt{\frac{2n!}{(n+L)!}} \zeta^{1/2+L} \exp\left(\frac{-\kappa^2 \zeta^2}{2}\right) L_n^L(x^2 \zeta^2)$$

- Lightest bound state ( $n = L = J = 0$ ) is massless ( $M = 0$ )
- $M^2 = 4\kappa^2 L \Rightarrow \kappa = 0.54 \text{ GeV}$  from Regge slope

# Light front Wavefunction for $\rho$ , $K^*$ and $\phi$

For the vector mesons  $\rho$  and  $\phi$ , we set  $n = 0, L = 0$  to obtain

$$\Psi_{0,0}(z, \zeta) = \frac{\kappa}{\sqrt{\pi}} \sqrt{z(1-z)} \exp \left[ -\frac{\kappa^2 \zeta^2}{2} \right]$$

Allowing for small quark masses, the wavefunction becomes

$$\Psi_\lambda(z, \zeta) = \mathcal{N}_\lambda \sqrt{z(1-z)} \exp \left[ -\frac{\kappa^2 \zeta^2}{2} \right] \exp \left[ -\frac{(1-z)m_q^2 + zm_{\bar{q}}^2}{2\kappa^2 z(1-z)} \right]$$

$m_{u,d} = 0.046$  GeV and  $m_s = 0.357$  GeV are fixed from the y-intercepts of the Regge trajectories. Decay constant provides the first test of the wave function:

$$f_V P^+ = \langle 0 | \bar{q}(0) \gamma^+ q(0) | V(P, L) \rangle$$

$$f_V = \sqrt{\frac{N_c}{\pi}} \int_0^1 dz \left[ 1 + \frac{m_q m_{\bar{q}} - \nabla_r^2}{z(1-z)M_V^2} \right] \Psi_L(r, z) |_{r=0}$$

# Predictions for leptonic decay width

We can use this decay constant to predict the experimentally measured electronic decay width  $\Gamma_{V \rightarrow e^+e^-}$  of the vector meson:

$$\Gamma_{V \rightarrow e^+e^-} = \frac{4\pi\alpha_{em}^2 C_V^2}{3M_V} f_V^2$$


where  $C_\phi = 1/3$  for the  $C_\rho = 1/\sqrt{2}$ .

Meson	$f_V$ [GeV]	$\Gamma_{e^+e^-}$ [KeV]	$\Gamma_{e^+e^-}$ [KeV] (PDG)
$\rho$	0.210, 0.211	6.355, 6.383	$7.04 \pm 0.06$
$\phi$	0.191, 0.205	0.891, 1.024	$1.251 \pm 0.021$

**Table:** Predictions for the electronic decay widths of the  $\rho$  and  $\phi$  vector mesons using the holographic wavefunction with  $m_{u,d} = 0.046, 0.14$  GeV and  $m_s = 0.357, 0.14$  GeV.



# Diffractive vector meson production



- $ep \rightarrow epV$  or  $\gamma^* p \rightarrow pV$
- Sensitivity to non-perturbative physics
- Can be used to fine tune the vector meson wavefunction

N. Sharma, R. Sandapen, MA, PhysRevD.94.074018(2016)

# $K^*$ decay constants

$f_{K^*}$  and "transverse decay constant"  $f_{K^*}^\perp$  defined as:

$$\langle 0 | \bar{q} [\gamma^\mu, \gamma^\nu] s | K^*(P, \epsilon) \rangle = 2f_{K^*}^\perp (\epsilon^\mu P^\nu - \epsilon^\nu P^\mu)$$

$$f_{K^*}^\perp(\mu) = \sqrt{\frac{N_c}{2\pi}} \int_0^1 dx (xm_{\bar{q}} + (1-x)m_s) \int db \mu J_1(\mu b) \frac{\Psi_T(\zeta, x)}{x(1-x)}$$

Approach	Scale $\mu$	$m_{\bar{q}}$ [MeV]	$m_s$ [MeV]	$f_{K^*}$ [MeV]	$f_{K^*}^\perp(\mu)$ [MeV]	$f_{K^*}^\perp / f_{K^*}(\mu)$
AdS/QCD	$\sim 1$ GeV	140	280	200	118	0.59
AdS/QCD	$\sim 1$ GeV	195	300	200	132	0.66
AdS/QCD	$\sim 1$ GeV	250	320	200	142	0.71
Experiment				$205 \pm 6$		
Lattice	2 GeV					$0.780 \pm 0.008$
Lattice	2 GeV					$0.74 \pm 0.02$

Comparison between AdS/QCD predictions for the decay constant of the  $K^*$  meson with experiment (obtained from  $\Gamma(\tau^- \rightarrow K^{*-} \nu_\tau)$ ), and the ratio of couplings with lattice data.

# Light cone distribution amplitudes

Light cone coordinates:  $x^\mu = (x^+, x^-, x_\perp)$ , where  $x^\pm = x^0 \pm x^3$  and  $x_\perp$  any combinations of  $x_1$  and  $x_2$ .

At equal light-front time  $x^+ = 0$  and in the light-front gauge  $A^+ = 0$ ,

$$\begin{aligned}\langle 0 | \bar{q}(0) \gamma^\mu q(x^-) | \rho(P, \epsilon) \rangle &= f_\rho M_\rho \frac{\epsilon \cdot X}{P^+ X^-} P^\mu \int_0^1 du e^{-iuP^+ x^-} \phi_\rho^\parallel(u, \mu) \\ &+ f_\rho M_\rho \left( \epsilon^\mu - P^\mu \frac{\epsilon \cdot X}{P^+ X^-} \right) \int_0^1 du e^{-iuP^+ x^-} g_\rho^\perp(\nu)(u, \mu)\end{aligned}$$

$$\langle 0 | \bar{q}(0) [\gamma^\mu, \gamma^\nu] q(x^-) | \rho(P, \epsilon) \rangle = 2f_\rho^\perp (\epsilon^\mu P^\nu - \epsilon^\nu P^\mu) \int_0^1 du e^{-iuP^+ x^-} \phi_\rho^\perp(u, \mu)$$

$$\langle 0 | \bar{q}(0) \gamma^\mu \gamma^5 q(x^-) | \rho(P, \epsilon) \rangle = -\frac{1}{4} \epsilon_{\nu\rho\sigma}^\mu \epsilon^\nu P^\rho X^\sigma f_\rho M_\rho \int_0^1 du e^{-iuP^+ x^-} g_\rho^{\perp(a)}(u, \mu)$$

Vector meson's polarization vectors  $\epsilon$  are chosen as

$$\epsilon_L = \left( \frac{P^+}{M_\rho}, -\frac{M_\rho}{P^+}, 0_\perp \right) \quad \text{and} \quad \epsilon_{T(\pm)} = \frac{1}{\sqrt{2}} (0, 0, 1, \pm i)$$

R. Sandapen, MA, PRD87.054013(2013)

$$\phi_{\rho}^{\parallel}(z, \mu) = \frac{N_c}{\pi f_{\rho} M_{\rho}} \int dr \mu J_1(\mu r) [M_{\rho}^2 z(1-z) + m_f^2 - \nabla_r^2] \frac{\phi_L(r, z)}{z(1-z)},$$

$$\phi_{\rho}^{\perp}(z, \mu) = \frac{N_c m_f}{\pi f_{\rho}^{\perp}} \int dr \mu J_1(\mu r) \frac{\phi_T(r, z)}{z(1-z)},$$


$$g_{\rho}^{\perp(\nu)}(z, \mu) = \frac{N_c}{2\pi f_{\rho} M_{\rho}} \int dr \mu J_1(\mu r) [m_f^2 - (z^2 + (1-z)^2) \nabla_r^2] \frac{\phi_T(r, z)}{z^2(1-z)^2}$$

$$\frac{dg_{\rho}^{\perp(a)}}{dz}(z, \mu) = \frac{\sqrt{2} N_c}{\pi f_{\rho} M_{\rho}} \int dr \mu J_1(\mu r) (1-2z) [m_f^2 - \nabla_r^2] \frac{\phi_T(r, z)}{z^2(1-z)^2}.$$


Distribution amplitudes are normalized:

$$\int_0^1 du \phi_{\rho}^{\perp, \parallel}(u, \mu) = \int_0^1 du g_{\rho}^{\perp(a, \nu)}(u, \mu) = 1$$

# AdS/QCD DAs for $\rho$ : comparison to Sum Rules




(a) Twist-2 DA for the longitudinally polarized  $\rho$  meson



(b) Twist-2 DA for the transversely polarized  $\rho$  meson

**Figure:** Twist-2 DAs for the  $\rho$  meson. Solid Red: AdS/QCD DA at  $\mu \sim 1$  GeV; Dashed Blue: Sum Rules DA at  $\mu = 2$  GeV.

# DAs for $K^*$ : comparison to Sum Rules



**Figure:** Twist-2 DAs predicted by AdS/QCD (graphs on the left) and SR (graphs on the right). The uncertainty band is due to the variation of the quark masses for AdS/QCD and the error bar on Gegenbauer coefficients for SR.

# Isospin asymmetry in $B \rightarrow K^* \gamma$

Isospin asymmetry defined as:

$$\Delta_{0-} = \frac{\Gamma(\bar{B}^0 \rightarrow \bar{K}^* \gamma) - \Gamma(B^- \rightarrow K^{*-} \gamma)}{\Gamma(\bar{B}^0 \rightarrow \bar{K}^* \gamma) + \Gamma(B^- \rightarrow K^{*-} \gamma)}.$$

Branching ratio	BABAR	BELLE	CLEO	PDG
$\mathcal{B}(B^0 \rightarrow K^{*0} \gamma) \times 10^6$	$44.7 \pm 1.0 \pm 1.6$	$45.5^{+7.2}_{-6.8} \pm 3.4$	$40.1 \pm 2.1 \pm 1.7$	$43.3 \pm 1.5$
$\mathcal{B}(B^+ \rightarrow K^{*+} \gamma) \times 10^6$	$42.2 \pm 1.4 \pm 1.6$	$42.5 \pm 3.1 \pm 2.4$	$37.6^{+8.9}_{-8.3} \pm 2.8$	$42.1 \pm 1.8$
$\Delta_{0-}$	$6.6 \pm 2.1 \pm 2.2$	$1.2 \pm 4.4 \pm 2.6$		$5.2 \pm 2.6$

# Isospin calculation

Based on original work by Kagan and Neubert: Phys.Lett. B539, 227 (2002)

$$F_{\perp}(\mu_h) = \int_0^1 dz \frac{\phi_{K^*}^{\perp}(z, \mu_h)}{3(1-z)}$$

$$G_{\perp}(s_c, \mu_h) = \int_0^1 dz \frac{\phi_{K^*}^{\perp}(z, \mu_h)}{3(1-z)} G(s_c, \bar{z})$$

$$X_{\perp}(\mu_h) = \int_0^1 dz \phi_{K^*}^{\perp}(z, \mu_h) \left( \frac{1+\bar{z}}{3\bar{z}^2} \right)$$

and

$$H_{\perp}(s_c, \mu_h) = \int_0^1 dz \left( g_{K^*}^{\perp(v)}(z, \mu_h) - \frac{1}{4} \frac{dg_{K^*}^{\perp(a)}}{dz}(z, \mu_h) \right) G(s_c, \bar{z})$$



R. Sandapen, MA, PRD88.014042(2013)

Integral	SR	AdS/QCD
$X_{\perp}$	$\infty$	26.9
$F_{\perp}$	1.14	1.38
$G_{\perp}$	$2.55 + 0.43i$	$2.89 + 0.30i$
$H_{\perp}$	$2.48 + 0.50i$	$2.12 + 0.21i$

Branching ratio for  $B \rightarrow K^* \gamma$ :  $44.3 \times 10^{-6}$  from AdS/QCD compared with  $45.9 \times 10^{-6}$  from Sum Rules

$\Delta_{0-} = 3.3\%$  from AdS/QCD

# $B \rightarrow \rho$ transition form factors





Form factors are defined as:

$$\begin{aligned}\langle \rho(k, \varepsilon) | \bar{q} \gamma^\mu (1 - \gamma^5) b | B(p) \rangle &= \frac{2iV(q^2)}{m_B + m_\rho} \epsilon^{\mu\nu\rho\sigma} \varepsilon_\nu^* k_\rho p_\sigma - 2m_\rho A_0(q^2) \frac{\varepsilon^* \cdot q}{q^2} q^\mu \\ &- (m_B + m_\rho) A_1(q^2) \left( \varepsilon^{\mu*} - \frac{\varepsilon^* \cdot q q^\mu}{q^2} \right) \\ &+ A_2(q^2) \frac{\varepsilon^* \cdot q}{m_B + m_\rho} \left[ (p + k)^\mu - \frac{m_B^2 - m_\rho^2}{q^2} q^\mu \right]\end{aligned}$$


$$\begin{aligned}q_\nu \langle \rho(k, \varepsilon) | \bar{d} \sigma^{\mu\nu} (1 - \gamma^5) b | B(p) \rangle &= 2T_1(q^2) \epsilon^{\mu\nu\rho\sigma} \varepsilon_\nu^* p_\rho k_\sigma \\ &- iT_2(q^2) [(\varepsilon^* \cdot q)(p + k)_\mu - \varepsilon_\mu^* (m_B^2 - m_\rho^2)] \\ &- iT_3(q^2) (\varepsilon^* \cdot q) \left[ \frac{q^2}{m_B^2 - m_\rho^2} (p + k)_\mu - q_\mu \right]\end{aligned}$$

# AdS/QCD prediction for $B \rightarrow \rho$ transition form factors

R. Campbell, S. Lord, R. Sandapen, MA, PRD88.074031(2013)



# Differential decay rate for semileptonic $B \rightarrow \rho l \bar{\nu}$



(a) Differential decay rate for the semileptonic  $B \rightarrow \rho l \bar{\nu}$  decay. The lattice data points are from UKQCD Collaboration.

# Numerical predictions

BaBar collaboration has measured partial branching fractions in  $q^2$  bins: PRD83, 032007 (2011)

$$\Delta B_{\text{low}} = \int_0^8 \frac{dB}{dq^2} dq^2 = (0.564 \pm 0.166) \times 10^{-4}$$

$$\Delta B_{\text{mid}} = \int_8^{16} \frac{dB}{dq^2} dq^2 = (0.912 \pm 0.147) \times 10^{-4}$$

$$\Delta B_{\text{high}} = \int_{16}^{20.3} \frac{dB}{dq^2} dq^2 = (0.268 \pm 0.062) \times 10^{-4}$$

$$R_{\text{low}} = \frac{\Delta B_{\text{low}}}{\Delta B_{\text{mid}}} = 0.618 \pm 0.207$$

$$R_{\text{high}} = \frac{\Delta B_{\text{high}}}{\Delta B_{\text{mid}}} = 0.294 \pm 0.083$$


Our predictions for  $m_f = 0.14, 0.35$ :

$$R_{\text{low}} = 0.580, 0.424$$

$$R_{\text{high}} = 0.427, 0.503$$


# AdS/QCD prediction for $B \rightarrow K^*$ transition form factors

R. Campbell, S. Lord, R. Sandapen, MA, PRD89.074021(2014)




# Isospin asymmetry in dileptonic $B \rightarrow K^* \mu^+ \mu^-$

S. Lord, R. Sandapen, MA, PRD90.074010(2014)



(b) Isospin asymmetry in  $B \rightarrow K^* \mu^+ \mu^-$  decay vs dileptonic invariant mass. The data points are from LHCb. The dashed red curve is the prediction of the QCD sum rules.

# Differential decay rate for $B \rightarrow K^* \nu \bar{\nu}$



(c) The AdS/QCD (Solid line) and SR (Dashed line) predictions for the differential Branching Ratio for  $B \rightarrow K^* \nu \bar{\nu}$ . The shaded band represents the uncertainty coming from the form factors.



# Summary and outlook

- AdS/QCD LFWF is used to obtain  $\rho$  and  $K^*$  DAs.
- DAs are essential ingredients for the calculation of the  $B \rightarrow \rho, K^*$  transition form factors via LCSR.
- We are in the process of calculating the form factors directly from LFWF.
- We have looked into the proper LFWF for pseudoscalar mesons (F. Chishtie, R. Sandapen, MA PRD95,074008(2017)) and working to predict  $B \rightarrow \pi, K$  transition form factors.