

Dark Matter Direct Detection (XENON1T world best sensitivity)

Julien Masbou
Subatech – Université de Nantes

Summer School in **Particle and Astroparticle physics** of
Annecy-le-Vieux
19-25 July 2018



What Dark Matter it not



What Dark Matter it not



➔ Barnard 68 : cold molecular cloud ~ 500 ly.
Transparent in infrared

Definition


By « Dark Matter » we mean non-luminous matter :
no associated emission of light (visible, UV, IR, radio, etc...)

... But we assume its existence by its gravitational effect in:


- 1) Galaxies
- 2) Galaxy clusters
- 3) Cosmology

Galaxies


In galaxies, stars are not static but turn around the galactic center. Thanks to the rotation, the centrifugal force compensates the gravitational force, which prevents stars from collapsing in the core.



Galaxies




Vitesse observée




Distance du centre

Solid rotation






Vitesse observée



Distance du centre

Planetary rotation


Galaxies




Rotation velocity almost constant at all radius !

\rightarrow Presence of a halo of invisible matter, 5-10 times heavier than standard matter

Vera Rubin ~1970



Gravitational lenses




Gravitational lenses

Astronetti.com


Abell 2218
Credit: NASA/STScI/AURA




Dark Matter 3D-map



Colliding clusters




Energy composition of the universe





Characteristics of Dark Matter Particles

- Weak interaction
- Stable
- Non-baryonic Matter
- Non relativistic



Direct dark matter detection principle



Nuclear
Recoil
(NR)



- Direct detection
- Indirect detection
- Production

Direct dark matter detection principle


Nuclear
Recoil
(NR)



Electronic
Recoil
(ER)

γ and β particles
interact with the atomic electrons
 \rightarrow background

Cinematic



$$\frac{m_\chi}{2} v^2 = \frac{m_\chi}{2} v'^2 + E_r \quad \left(E_r = \frac{1}{2} m_N w^2 \right)$$

$$m_\chi v = m_\chi v' \cos \varphi + m_N w \cos \theta_r$$
$$m_\chi v' \sin \varphi = m_N w \sin \theta_r$$





$$E_r = \left(\frac{m_\chi}{2} v^2 \right) \times \frac{4 m_N m_\chi}{(m_N + m_\chi)^2} \times \cos^2 \vartheta_r$$


$\sim 1 - 100 \text{ keV}$

Expected rate for terrestrial detector

$$\frac{dR}{dE_R} = N_N \frac{\rho_\odot}{m_\chi} \int_{v_{min}}^{v_{max}} f(v) v \frac{d\sigma}{dE_R} dv$$




$$v_{min} = \sqrt{\frac{m_N E_{th}}{2\mu}}$$




How is evolving the field of Direct Detection ?

$$R \sim 0.13 \frac{\text{events}}{\text{kg} \cdot \text{year}} \left[\frac{A}{100} \times \frac{\sigma_{\chi N}}{10^{-38} \text{ cm}^2} \times \frac{\langle v \rangle}{220 \text{ km.s}^{-1}} \times \frac{\rho_{\odot}}{0.3 \text{ GeV.cm}^{-3}} \right]$$




Direct detection : progress over time



Detectors needs

- ultra-low background experimental environment
- low energy threshold to detect small recoil energy signals
- good discrimination power against particle that might mimic WIMP collision
- large detector mass to enhance the interaction probability inside the target



The fight against the background


- **Avoid background**
- **External γ 's** from natural radioactivity
 - Material screening
 - Self shielding (fiducialization)
- **External neutrons**
muon-induced (α, n) and fission reaction
 - Material screening (low U and Th)
 - Underground experiments
 - Shield & active veto
- **Internal contamination**
 - ^{85}Kr : removed by cryogenic distillation
 - ^{222}Rn : removed by cryogenic distillation
 - ^{136}Xe : $\beta\beta$ decay, long lifetime ($T_{1/2} = 2.2 \times 10^{21}$ years)

- **Use WIMP properties**
 - No double scatter
 - Homogeneously distributed
→ *Position reconstruction*
 - Nuclear recoils
→ *ER/NR Discrimination*



Cosmic Rays

To increase the sensitivity of the experiments, we need:

- To hide under a mountain to be protected from cosmic rays (100 per second across our body),
- To be protected from natural radioactivity from rocks
- To purify from materials of the detector



XENON1T experiment site



PERIODIC TABLE OF ELEMENTS

Relative Atomic Mass: 12.0107
 Oxidation State: -4, +2, +4
 Atomic Number: 6
 Element Symbol: C
 Atomic Electron Configuration: 2-4



"The chemistry must be respected."
 -Walter "Walt" Hartwell White Sr. (1959-2013)

METALLOIDS

Si
 14
 28.0855

d-Block

p-Block w/o He

s-Block + He

1	1 1.00794 +1 H	2 2 4.002602 +2 He
2	3 6.941 +1 Li	4 9.01218 +2 Be
3	11 22.98977 +1 Na	12 24.305 +2 Mg
4	19 39.0983 +1 K	20 40.08 +2 Ca
5	37 85.468 +1 Rb	38 87.62 +2 Sr
6	55 132.905 +1 Cs	56 137.33 +2 Ba
7	87 [223] +1 Fr	88 226.025 +2 Ra

3	4	5	6	7	8	9	10	11	12
IIIB	IVB	VB	VIB	VII B	VIII	VIII	VIII	IB	IIB
21 44.9559 +3 Sc	22 47.88 +2 Ti	23 50.9415 +2 V	24 51.996 +2 Cr	25 54.9390 +2 Mn	26 55.845 +2 Fe	27 58.9332 +2 Co	28 58.693 +2 Ni	29 63.546 +2 Cu	30 65.39 +2 Zn
39 88.9059 +3 Y	40 91.224 +4 Zr	41 92.90636 +3 Nb	42 95.94 +3 Mo	43 [98] +7 Tc	44 101.07 +3 Ru	45 102.906 +1 Rh	46 106.42 +2 Pd	47 107.868 +1 Ag	48 112.411 +2 Cd
57 138.906 +3 La	58 178.49 +4 Hf	59 180.945 +5 Ta	60 183.85 +6 W	61 186.207 +7 Re	62 190.23 +4 Os	63 192.217 +4 Ir	64 196.078 +2 Pt	65 196.967 +3 Au	66 200.59 +2 Hg
89 [223] +1 Ac	90 [261] +2 Rf	91 [262] +3 Db	92 [263] +4 Sg	93 [262] +5 Bh	94 [265] +6 Hs	95 [266] +7 Mt	96 [268] +8 Ds	97 [269] +9 Rg	98 [277] +10 Cn

13	14	15	16	17	18
IIIA	IVA	VA	VIA	VIIA	VIIIA
5 10.811 +3 B	6 12.0107 +4 C	7 14.0067 +3 N	8 15.999 +2 O	9 18.998 +1 F	10 4.003 0 He
13 26.982 +3 Al	14 28.0855 +4 Si	15 30.974 +3 P	16 32.065 +2 S	17 35.453 +1 Cl	18 39.948 0 Ar
31 69.723 +3 Ga	32 72.64 +4 Ge	33 74.922 +3 As	34 78.96 +2 Se	35 79.904 +1 Br	36 83.80 0 Kr
49 114.818 +3 In	50 118.710 +2 Sn	51 121.75 +3 Sb	52 127.60 +2 Te	53 126.904 +1 I	54 131.293 0 Xe
81 204.383 +3 Tl	82 207.2 +2 Pb	83 208.980 +3 Bi	84 [209] +2 Po	85 [210] +1 At	86 [222] 0 Rn
113 [288] +3 Uut	114 [289] +2 Fl	115 [288] +3 Uup	116 [289] +2 Lv	117 [289] +1 Uus	118 [289] 0 Uuo


Breaking
 Bad

f-Block

58 140.116 +3 Ce	59 140.907 +4 Pr	60 144.24 +3 Nd	61 [147] +3 Pm	62 150.36 +3 Sm	63 151.964 +3 Eu	64 157.25 +3 Gd	65 158.925 +3 Tb	66 162.50 +3 Dy	67 164.930 +3 Ho	68 167.259 +3 Er	69 168.934 +3 Tm	70 173.04 +3 Yb	71 174.967 +3 Lu
90 [232] +3 Th	91 [231] +5 Pa	92 [238] +6 U	93 [237] +5 Np	94 [244] +4 Pu	95 [243] +3 Am	96 [247] +3 Cm	97 [247] +3 Bk	98 [251] +3 Cf	99 [252] +3 Es	100 [257] +3 Fm	101 [258] +3 Md	102 [258] +3 No	103 [262] +3 Lr


Why Xenon ?

- Large mass number A (131) (Interaction cross section $\propto A^2$)
- 50% odd isotopes (^{129}Xe , ^{131}Xe) for Spin-Dependent interactions
- Kr can be reduced to ppt levels
- High stopping power, i.e. active volume is self-shielding
- Efficient scintillator (178 nm)
- Scalable to large target masses
- Electronic recoil discrimination with simultaneous measurement of scintillation and ionization



Dual phase TPC: principle

TPC = Time Projection Chamber



S1:

→ Photon ($\lambda = 178 \text{ nm}$)
from Scintillation process

→ Detected by PMTs
(mainly bottom array)

S2:

→ Electrons drift



→ Extraction in gaseous phase

→ Proportional scintillation light

3D reconstruction :


→ X,Y from top array

→ Z from Drift time






$$(S2/S1)_{WIMP,n} < (S2/S1)_{\gamma,\beta}$$

Dual phase TPC: real life



X and Y position from S2 hit pattern on the top PMTs



XENON World

25 Institutions
11 Countries
165 Scientists



Phases of the XENON Program



XENON10

2005 – 2007
15 cm drift TPC
Total: 25 kg
Target: **14** kg
Fiducial: 5.4 kg


Achieved (2007)
 $\sigma_{SI} = 8.8 \cdot 10^{-44} \text{ cm}^2$
@ 100 GeV/c²



XENON100

2008 – 2016
30 cm drift TPC
Total: 161 kg
Target: **62** kg
Fiducial: 34/48 kg


Achieved (2016)
 $\sigma_{SI} = 1.1 \cdot 10^{-45} \text{ cm}^2$
@ 55 GeV/c²



XENON1T

2012 – 2019
100 cm drift TPC
Total: 3 200 kg
Target: **2 000** kg
Fiducial: 1 000 kg

Achieved (2018)
 $\sigma_{SI} = 4.1 \cdot 10^{-47} \text{ cm}^2$
@ 30 GeV/c²



XENONnT

2017 (R&D) – 2023
144 cm drift TPC
Total: 8 000 kg
Target: **6 000** kg
Fiducial: 4 500 kg

Projected (2022)
 $\sigma_{SI} = 1.6 \times 10^{-48} \text{ cm}^2$
@ 50 GeV/c²

XENON1T facility

Water shield: deionized water as passive radiation shield


Muon veto: Active muon veto against muon induced neutrons (84 PMTs)

Cryogenics: Stable conditions (3.2t LXe)
Purification: LXe flow through getters, remove impurities


DAQ: Each channel has its own threshold, Flexible software algorithms
Readout: Up to 300MB/s for high rate calibrations

ReStoX: Emergency recovery up to 7.6 tons of LXe
Passive: No active cooling required to keep Xe contained

Kr Distillation: Remove Kr from system during fill or online
Rn Distillation: Initial tests show promising reduction for Rn



XENON1T Data Taking






- DM total exposure SR0+SR1: 278.8 Live days
- Calibration Data:
 - 83mKr → Spatial Response (position correction)
 - 220Rn → ER-Bands
 - 241AmBe & NG → NR-Bands
 - LED → PMT gain monitoring

Calibrations


Electronic Recoils

- ^{228}Th source emanates ^{220}Rn into LXe
- β -decay of ^{212}Pb to ^{212}Bi \rightarrow low energy events (2–20 keV)
- Decay of activity dominated by ^{212}Pb half-life (10.6 h)

Internal source



External source




Nuclear Recoils

- External $^{241}\text{AmBe}$ source mounted on a belt
- The α particles emitted by the decay of the Am collide with the light Be nuclei producing fast neutrons
- Neutron Generator

Dark Matter Search Data


- **Blinding** → to avoid biases in event selection and signal/background modeling
- **Salting** (addition of fake events) → to protect against post-unblinding tuning of the cuts and background models



Fiducial Volume Optimization

Optimize fiducial volume before unblinding by using improved understanding

- position reconstruction
- detector response
- correlations between spectral and spacial distribution
- include knowledge on background distributions in statistical framework
- MC simulations




Aim at
optimal S/B

larger FV
1 t → 1.3 t

XENON1T Expectations

	1.3 t	0.65 t	Mass
278.8 days live-time	Full ROI	NR Reference	(S2,S1) region
ER	627 ± 18	0.60 ± 0.13	
neutron	1.43 ± 0.66	0.14 ± 0.07	
CEvNS	0.05 ± 0.01	0.01	
AC	$0.47^{+0.27}$	$0.04^{+0.02}$	
Surface	106 ± 8	0.01	
TOTAL BKG	735 ± 20	0.80 ± 0.14	



WIMP
50 GeV/c²

► Background models


In 4-dimensional space: S1, S2, r, z

► Statistical inference


Done with PLR analysis in 1.3 t fiducial volume and full (S1,S2) space, corresponding to [4.9, 40.9] keV_{nr} and [1.4, 10.6] keV_{ee}.

► NR reference region

Between NR median and -2σ quantile. Numbers in table are for illustration; final results from complete PLR statistical inference.




Dark Matter Search Results



- Results interpreted with unbinned profile likelihood analysis in cs1, cs2, R space
- Piechart indicate the relative probabilities of this event to be of a certain class for a best fit to a 200 GeV/c² WIMPs with a cross-section of 4.6×10^{-47} cm²

Spacial Distribution of Dark Matter Search Results




- Core volume to distinguish WIMPs over neutron background
- Yellow shaded regions display the 1σ (dark), and 2σ (light) probability density percentiles of the radiogenic neutron background component

XENON1T Results

► Spin-independent WIMP-nucleon cross section

Strongest exclusion limits (at 90% CL) on WIMPs $> 6 \text{ GeV}/c^2$.



arxiv 1805.12562

7 times better sensitivity compared to previous experiments (LUX, PANDAX-II)

World best limit:


First 1 ton x years exposition !

$$\sigma_{SI} < 4.1 \cdot 10^{-47} \text{ cm}^2 \text{ at } 30 \text{ GeV}/c^2$$


► 1 sigma upper fluctuation at higher WIMP masses

Local p-value ~ 0.2 (at $200 \text{ GeV}/c^2$). No significant excess (> 3 sigma) is observed.

From XENON1T to XENONnT



Conclusion & Perspectives



And other analysis already published or to come:

- Axions / ALP
- 2ν double electron capture on ^{124}Xe
- Low mass
- Effective field theories
- Calibration
- Neutrinos
- ...
- Stay tuned !

- **Dark matter is highly searched**
- Solution to an astrophysics / particle physics / Cosmology problem