

4 september 2009  
IPHC Strasbourg

# Seeing Dark Matter in cosmic rays?!?

Marco Cirelli  
(CNRS, IPhT-CEA/Saclay)

in collaboration with:

A.Strumia (Pisa)  
N.Fornengo (Torino)  
M.Tamburini (Pisa)  
R.Franceschini (Pisa)  
M.Raidal (Tallin)  
M.Kadastik (Tallin)  
Gf.Bertone (IAP Paris)  
M.Taoso (Padova)  
C.Bräuninger (Saclay)  
P.Panci (Saclay)  
F.Iocco (Saclay + IAP Paris)

Nuclear Physics B 753 (2006)  
Nuclear Physics B 787 (2007)  
Nuclear Physics B 800 (2008)  
0808.3867 [astro-ph]  
Nuclear Physics B 813 (2009)  
JCAP 03 009 (2009)  
Physics Letters B 678 (2009)  
Nuclear Physics B 821 (2009)  
0907.0719

*and work in progress*

4 september 2009  
IPHC Strasbourg

# Seeing Dark Matter in cosmic rays?!?

Marco Cirelli  
(CNRS, IPhT-CEA/Saclay)

in collaboration with:

A.Strumia (Pisa)  
N.Fornengo (Torino)  
M.Tamburini (Pisa)  
R.Franceschini (Pisa)  
M.Raidal (Tallin)  
M.Kadastik (Tallin)  
Gf.Bertone (IAP Paris)  
M.Taoso (Padova)  
C.Bräuninger (Saclay)  
P.Panci (Saclay)  
F.Iocco (Saclay + IAP Paris)

Nuclear Physics B 753 (2006)  
Nuclear Physics B 787 (2007)  
Nuclear Physics B 800 (2008)  
0808.3867 [astro-ph]  
Nuclear Physics B 813 (2009)  
JCAP 03 009 (2009)  
Physics Letters B 678 (2009)  
Nuclear Physics B 821 (2009)  
0907.0719

*and work in progress*

# Questions


1. Are we seeing Dark Matter  
in cosmic rays?

# Questions

1. Are we seeing Dark Matter  
in cosmic rays?
  
2. Why  $\gtrsim 300$  new DM models have  
been proposed in one year?

# The Evidence for DM

1) galaxy rotation curves




$$\Omega_M \gtrsim 0.1$$

2) clusters of galaxies



$$\Omega_M \sim 0.2 \div 0.4$$

3) CMB+LSS(+SNIa:)



$$\Omega_M \approx 0.26 \pm 0.05$$




DM exists.

It consists of a particle.  
Permeates galactic haloes.

# The Evidence for DM

1) galaxy rotation curves




$$\Omega_M \gtrsim 0.1$$

2) clusters of galaxies



$$\Omega_M \sim 0.2 \div 0.4$$

3) CMB+LSS(+SNIa:)



$$\Omega_M \approx 0.26 \pm 0.05$$




What is the DM??  
It consists of a particle.  
Permeates galactic haloes.

# A thermal relic from the Early Universe

Boltzmann equation  
in the Early Universe:

$$\Omega_X \approx \frac{6 \cdot 10^{-27} \text{ cm}^3 \text{s}^{-1}}{\langle \sigma_{\text{ann}} v \rangle}$$

Relic  $\Omega_{\text{DM}} \simeq 0.23$  for  
 $\langle \sigma_{\text{ann}} v \rangle = 3 \cdot 10^{-26} \text{ cm}^3/\text{sec}$



Weak cross section:

$$\langle \sigma_{\text{ann}} v \rangle \approx \frac{\alpha_w^2}{M^2} \approx \frac{\alpha_w^2}{1 \text{ TeV}^2} \Rightarrow \Omega_X \sim \mathcal{O}(\text{few } 0.1) \quad (\text{WIMP})$$

# DM detection

direct detection

Xenon, CDMS (Dama/Libra?)

production at colliders

LHC

$\gamma$  from annihil in galactic halo or center  
(line + continuum) Fermi

indirect

$e^+$  from annihil in galactic halo or center  
PAMELA, ATIC, Fermi

$\bar{p}$  from annihil in galactic halo or center

$\bar{D}$  from annihil in galactic halo or center

GAPS

$\nu, \bar{\nu}$  from annihil in massive bodies

Icecube, Km3Net

# DM detection

direct detection

production at colliders

$\gamma$  from annihil in galactic halo or center  
(line + continuum)

indirect  $e^+$  from annihil in galactic halo or center  
PAMELA, ATIC, Fermi

$\bar{p}$  from annihil in galactic halo or center

$\bar{D}$  from annihil in galactic halo or center

$\nu, \bar{\nu}$  from annihil in massive bodies

# DM detection

direct detection

production at colliders

$\gamma$  from annihil in galactic halo or center  
(line + continuum)

indirect  $e^+$  from annihil in galactic halo or center

PAMELA, ATIC, Fermi


$\bar{p}$  from annihil in galactic halo or center

$\bar{D}$  from annihil in galactic halo or center

$\nu, \bar{\nu}$  from annihil in massive bodies


# Indirect Detection

$\bar{p}$  and  $e^+$  from DM annihilations in halo




# Indirect Detection

$\bar{p}$  and  $e^+$  from DM annihilations in halo




# Indirect Detection

$\bar{p}$  and  $e^+$  from DM annihilations in halo




# Indirect Detection

$\bar{p}$  and  $e^+$  from DM annihilations in halo




# Indirect Detection

$\bar{p}$  and  $e^+$  from DM annihilations in halo




# Indirect Detection

$\bar{p}$  and  $e^+$  from DM annihilations in halo



# Indirect Detection

$\bar{p}$  and  $e^+$  from DM annihilations in halo




$$\frac{\partial f}{\partial t} - K(E) \cdot \nabla^2 f - \frac{\partial}{\partial E} (b(E)f) + \frac{\partial}{\partial z} (V_c f) = Q_{\text{inj}} - 2h\delta(z)\Gamma_{\text{spall}}f$$

diffusion      energy loss      convective wind      source      spallations

Salati, Chardronnat, Barrau,  
Donato, Tallet, Fornengo,  
Maurin, Brun... '90s, '00s

# Indirect Detection

$\bar{p}$  and  $e^+$  from DM annihilations in halo




What sets the overall expected flux?

$$\text{flux} \propto n^2 \sigma_{\text{annihilation}}$$

# Indirect Detection

$\bar{p}$  and  $e^+$  from DM annihilations in halo




What sets the overall expected flux?

$$\text{flux} \propto n^2 \sigma_{\text{annihilation}} \text{particle}$$

astro&cosmo

# Indirect Detection

$\bar{p}$  and  $e^+$  from DM annihilations in halo



What sets the overall expected flux?

$$\text{flux} \propto n^2 \sigma_{\text{annihilation}} \text{particle}$$

astro&  
cosmo

reference cross section:  
 $\sigma v = 3 \cdot 10^{-26} \text{ cm}^3/\text{sec}$


# From N-body numerical simulations:

$$\rho(r) = \rho_{\odot} \left[ \frac{r_{\odot}}{r} \right]^{\gamma} \left[ \frac{1 + (r_{\odot}/r_s)^{\alpha}}{1 + (r/r_s)^{\alpha}} \right]^{(\beta-\gamma)/\alpha}$$

Halo model	$\alpha$	$\beta$	$\gamma$	$r_s$ in kpc
Cored isothermal	2	2	0	5
Navarro, Frenk, White	1	3	1	20
Moore	1	3	1.16	30

At small r:  $\rho(r) \propto 1/r^\gamma$

$$\rho(r) = \rho_s \cdot \exp \left[ -\frac{2}{\alpha} \left( \left( \frac{r}{r_s} \right)^\alpha - 1 \right) \right]$$



# cuspy: NFW, Moore



# mild: Einasto

## smooth: isothermal

# Indirect Detection

Boost Factor: local clumps in the DM halo enhance the density, boost the flux from annihilations. Typically:  $B \simeq 1 \rightarrow 20$


For illustration:




Bertone, Branchini, Pieri 2007

Computing the theory  
predictions


# Spectra at production




# Spectra at production






# Spectra at production



# Spectra at production



# Spectra at production



So what are the particle physics parameters?


1. Dark Matter mass
2. primary channel(s)

# Comparing with data

# Data sets

Positrons from PAMELA:

**Payload for  
Anti-  
Matter  
Exploration and  
Light-nuclei  
Astrophysics**



calibrated on accelerator fluxes

magnetic spectrometer:  
charge and energy

calorimeter:  $e^\pm$  vs  $p/\bar{p}$   
(make showers)  
(swipe thru)

Big challenge: backgnd contamination  
from p ( $10^4$  more numerous at 100 GeV)

# Data sets


## Positrons from PAMELA:

- steep  $e^+$  excess above 10 GeV!
- very large flux!

$$\text{positron fraction: } \frac{e^+}{e^+ + e^-}$$

(9430  $e^+$  collected)


(errors statistical only,  
that's why larger at high energy)



# Data sets

## Positrons from PAMELA:

- steep  $e^+$  excess above 10 GeV!
- very large flux!




# Data sets

## Antiprotons from PAMELA:

- consistent with the background

(about 1000  $\bar{p}$  collected)



# Results

Which DM spectra can fit the data?


# Results

Which DM spectra can fit the data?

E.g. a DM with:

- mass  $M_{\text{DM}} = 150 \text{ GeV}$
- annihilation  $\text{DM DM} \rightarrow W^+W^-$
- (a possible SuperSymmetric candidate: wino)

Positrons:




# Results


Which DM spectra can fit the data?

E.g. a DM with:  
-mass  $M_{\text{DM}} = 150 \text{ GeV}$   
-annihilation  $\text{DM DM} \rightarrow W^+W^-$   
(a possible SuperSymmetric candidate: wino)

Positrons:



Anti-protons:



[insisting on Winos]

# Results

Which DM spectra can fit the data?

E.g. a DM with: -mass  $M_{\text{DM}} = 10 \text{ TeV}$


-annihilation  $\text{DM DM} \rightarrow W^+W^-$

# Results


Which DM spectra can fit the data?

E.g. a DM with: -mass  $M_{\text{DM}} = 10 \text{ TeV}$   
-annihilation  $\text{DM DM} \rightarrow W^+W^-$

Positrons:



Anti-protons:




# Results

Which DM spectra can fit the data?


E.g. a DM with:  
-mass  $M_{\text{DM}} = 10 \text{ TeV}$   
-annihilation  $\text{DM DM} \rightarrow W^+ W^-$   
but...: -cross sec  $\sigma_{\text{ann}} v = 6 \cdot 10^{-22} \text{ cm}^3/\text{sec}$

Mmm...

Positrons:



Anti-protons:




# Results

Which DM spectra can fit the data?

Model-independent results:

fit to PAMELA positrons only




# Results

Which DM spectra can fit the data?

Model-independent results:

fit to PAMELA positrons + anti-protons




# Results

Which DM spectra can fit the data?

Model-independent results:

fit to PAMELA positrons + anti-protons




(1) annihilate into leptons (e.g.  $\mu^+ \mu^-$ )

# Results

Which DM spectra can fit the data?

Model-independent results:

fit to PAMELA positrons + anti-protons




- (1) annihilate into leptons (e.g.  $\mu^+ \mu^-$ ) or
- (2) annihilate into  $W^+ W^-$  with mass  $\gtrsim 10$  TeV

# Results

Which DM spectra can fit the data?

Model-independent results:

Cross section required by PAMELA



# Data sets

Electrons + positrons from ATIC, PPB-BETS:



PPB-BETS  
(Japan)

Polar  
Patrol  
Balloon  
of the  
Balloon-borne  
Electron  
Telescope with  
Scintillating  
fibers



Advanced  
Thin  
Ionization  
Calorimeter


- bigger/denser: higher energy
- calorimeter only, no magnet:  
no charge discrimination

# Data sets

Electrons + positrons from ATIC, PPB-BETS:

- an  $e^+ + e^-$  excess at  $\sim 700$  GeV??

(ATIC: 1724  $e^+ + e^-$  collected  
at  $>100$  GeV;  $4\sigma$  above bkgnd)



# Results

Which DM spectra can fit the data?

A DM with:


- mass  $M_{\text{DM}} = 1 \text{ TeV}$
- annihilation  $\text{DM DM} \rightarrow \mu^+ \mu^-$

# Results


Which DM spectra can fit the data?

A DM with: -mass  $M_{\text{DM}} = 1 \text{ TeV}$   
-annihilation  $\text{DM DM} \rightarrow \mu^+ \mu^-$


Positrons:



Anti-protons:



Electrons + Positrons:




# Results


Which DM spectra can fit the data?

A DM with: -mass  $M_{\text{DM}} = 1 \text{ TeV}$   
-annihilation  $\text{DM DM} \rightarrow \mu^+ \mu^-$


Positrons:



Anti-protons:



Electrons + Positrons:



Have we identified the DM  
for the first time???

Arkani-Hamed, Weiner et al. 0810: Yes!  
+ a ton of others

[pulsar]

# Results

## Which DM can fit the data?

M.Pospelov and A.Ritz, 0810.1502: Secluded DM - A.Nelson and C.Spitzer, 0810.5167: Slightly Non-Minimal DM - Y.Nomura and J.Thaler, 0810.5397: DM through the Axion Portal - R.Harnik and G.Kribs, 0810.5557: Dirac DM - D.Feldman, Z.Liu, P.Nath, 0810.5762: Hidden Sector - T.Hambye, 0811.0172: Hidden Vector - Yin, Yuan, Liu, Zhang, Bi, Zhu, 0811.0176: Leptonically decaying DM - K.Ishiwata, S.Matsumoto, T.Moroi, 0811.0250: Superparticle DM - Y.Bai and Z.Han, 0811.0387: sUED DM - P.Fox, E.Poppitz, 0811.0399: Leptophilic DM - C.Chen, F.Takahashi, T.T.Yanagida, 0811.0477: Hidden-Gauge-Boson DM - K.Hamaguchi, E.Nakamura, S.Shirai, T.T.Yanagida, 0811.0737: Decaying DM in Composite Messenger - E.Ponton, L.Randall, 0811.1029: Singlet DM - A.Ibarra, D.Tran, 0811.1555: Decaying DM - S.Baek, P.Ko, 0811.1646: U(1) Lmu-Ltau DM - C.Chen, F.Takahashi, T.T.Yanagida, 0811.3357: Decaying Hidden-Gauge-Boson DM - I.Cholis, G.Dobler, D.Finkbeiner, L.Goodenough, N.Weiner, 0811.3641: 700+ GeV WIMP - E.Nardi, F.Sannino, A.Strumia, 0811.4153: Decaying DM in TechniColor - K.Zurek, 0811.4429: Multicomponent DM - M.Ibe, H.Murayama, T.T.Yanagida, 0812.0072: Breit-Wigner enhancement of DM annihilation - E.Chun, J.-C.Park, 0812.0308: sub-GeV hidden U(1) in GMSB - M.Lattanzi, J.Silk, 0812.0360: Sommerfeld enhancement in cold substructures - M.Pospelov, M.Trott, 0812.0432: super-WIMPs decays DM - Zhang, Bi, Liu, Liu, Yin, Yuan, Zhu, 0812.0522: Discrimination with SR and IC - Liu, Yin, Zhu, 0812.0964: DMnu from GC - M.Pohl, 0812.1174: electrons from DM - J.Hisano, M.Kawasaki, K.Kohri, K.Nakayama, 0812.0219: DMnu from GC - A.Arvanitaki, S.Dimopoulos, S.Dubovsky, P.Graham, R.Harnik, S.Rajendran, 0812.2075: Decaying DM in GUTs - R.Allahverdi, B.Dutta, K.Richardson-McDaniel, Y.Santoso, 0812.2196: SuSy B-L DM- S.Hamaguchi, K.Shirai, T.T.Yanagida, 0812.2374: Hidden-Fermion DM decays - D.Hooper, A.Stebbins, K.Zurek, 0812.3202: Nearby DM clump - C.Delaunay, P.Fox, G.Perez, 0812.3331: DMnu from Earth - Park, Shu, 0901.0720: Split-UED DM - Gogoladze, R.Khalid, Q.Shafi, H.Yuksel, 0901.0923: cMSSM DM with additions - Q.H.Cao, E.Ma, G.Shaughnessy, 0901.1334: Dark Matter: the leptonic connection - E.Nezri, M.Tytgat, G.Vertongen, 0901.2556: Inert Doublet DM - C.-H.Chen, C.-Q.Geng, D.Zhuridov, 0901.2681: Fermionic decaying DM - J.Mardon, Y.Nomura, D.Stolarski, J.Thaler, 0901.2926: Cascade annihilations (light non-abelian new bosons) - P.Meade, M.Papucci, T.Volansky, 0901.2925: DM sees the light - D.Phalen, A.Pierce, N.Weiner, 0901.3165: New Heavy Lepton - T.Banks, J.-F.Fortin, 0901.3578: Pyrma baryons - Goh, Hall, Kumar, 0902.0814: Leptonic Higgs - K.Bae, J.-H. Huh, J.Kim, B.Kyae, R.Viollier, 0812.3511: electrophilic axion from flipped-SU(5) with extra spontaneously broken symmetries and a two component DM with  $Z_2$  parity - ...

# Results

## Which DM can fit the data?


M.Pospelov and A.Ritz, 0810.1502: Secluded DM - A.Nelson and C.Spitzer, 0810.5167: Slightly Non-Minimal DM - Y.Nomura and J.Thaler, 0810.5397: DM through the Axion Portal - R.Harnik and G.Kribs, 0810.5557: Dirac DM - D.Feldman, Z.Liu, P.Nath, 0810.5762: Hidden Sector - T.Hambye, 0811.0172: Hidden Vector - Yin, Yuan, Liu, Zhang, Bi, Zhu, 0811.0176: **Leptonically decaying** DM - K.Ishiwata, S.Matsumoto, T.Moroi, 0811.0250: Superparticle DM - Y.Bai and Z.Han, 0811.0387: sUED DM - P.Fox, E.Poppitz, 0811.0399: **Leptophilic DM** - C.Chen, F.Takahashi, T.T.Yanagida, 0811.0477: Hidden-Gauge-Boson DM - K.Hamaguchi, E.Nakamura, S.Shirai, T.T.Yanagida, 0811.0737: Decaying DM in Composite Messenger - E.Ponton, L.Randall, 0811.1029: Singlet DM - A.Ibarra, D.Tran, 0811.1555: Decaying DM - S.Baek, P.Ko, 0811.1646: U(1) Lmu-Ltau DM - C.Chen, F.Takahashi, T.T.Yanagida, 0811.3357: Decaying Hidden-Gauge-Boson DM - I.Cholis, G.Dobler, D.Finkbeiner, L.Goodenough, N.Weiner, 0811.3641: **700+ GeV WIMP** - E.Nardi, F.Sannino, A.Strumia, 0811.4153: Decaying DM in TechniColor - K.Zurek, 0811.4429: Multicomponent DM - M.Ibe, H.Murayama, T.T.Yanagida, 0812.0072: Breit-Wigner enhancement of DM annihilation - E.Chun, J.-C.Park, 0812.0308: sub-GeV hidden U(1) in GMSB - M.Lattanzi, J.Silk, 0812.0360: Sommerfeld enhancement in cold substructures - M.Pospelov, M.Trott, 0812.0432: super-WIMPs decays DM - Zhang, Bi, Liu, Liu, Yin, Yuan, Zhu, 0812.0522: Discrimination with SR and IC - Liu, Yin, Zhu, 0812.0964: DMnu from GC - M.Pohl, 0812.1174: electrons from DM - J.Hisano, M.Kawasaki, K.Kohri, K.Nakayama, 0812.0219: DMnu from GC - A.Arvanitaki, S.Dimopoulos, S.Dubovsky, P.Graham, R.Harnik, S.Rajendran, 0812.2075: Decaying DM in GUTs - R.Allahverdi, B.Dutta, K.Richardson-McDaniel, Y.Santoso, 0812.2196: SuSy B-L DM- S.Hamaguchi, K.Shirai, T.T.Yanagida, 0812.2374: Hidden-Fermion DM decays - D.Hooper, A.Stebbins, K.Zurek, 0812.3202: Nearby DM clump - C.Delaunay, P.Fox, G.Perez, 0812.3331: DMnu from Earth - Park, Shu, 0901.0720: Split-UED DM - Gogoladze, R.Khalid, Q.Shafi, H.Yuksel, 0901.0923: cMSSM DM with additions - Q.H.Cao, E.Ma, G.Shaughnessy, 0901.1334: Dark Matter: the **leptonic connection** - E.Nezri, M.Tytgat, G.Vertongen, 0901.2556: Inert Doublet DM - C.-H.Chen, C.-Q.Geng, D.Zhuridov, 0901.2681: Fermionic decaying DM - J.Mardon, Y.Nomura, D.Stolarski, J.Thaler, 0901.2926: Cascade annihilations (light non-abelian new bosons) - P.Meade, M.Papucci, T.Volansky, 0901.2925: DM sees the light - D.Phalen, A.Pierce, N.Weiner, 0901.3165: New Heavy Lepton - T.Banks, J.-F.Fortin, 0901.3578: Pyrma baryons - Goh, Hall, Kumar, 0902.0814: Leptonic Higgs - K.Bae, J.-H. Huh, J.Kim, B.Kyae, R.Viollier, 0812.3511: electrophilic axion from flipped-SU(5) with extra spontaneously broken symmetries and a two component DM with  $Z_2$  parity - ...

# Results

Which DM spectra can fit the data?

Model-independent results:

fit to PAMELA positrons\* + balloon experiments




\* adding anti-protons does not change much, non-leptonic channels give too smooth spectrum for balloons

# Results

Which DM spectra can fit the data?

Model-independent results:

fit to PAMELA positrons\* + balloon experiments



(1) annihilate into leptons (e.g.  $\mu^+ \mu^-$ ), mass  $\sim 1$  TeV


# Data sets

Electrons + positrons from FERMI and HESS:



**FERMI-LAT**

(USA + France + Italy + Germany + Japan + Sweden)



**HESS**

(Europe + Africa)


“Designed as a high-sensitivity gamma-ray observatory, the FERMI Large Area Telescope is also an electron detector with a large acceptance”

“The very large collection area of ground-based gamma-ray telescopes gives them a substantial advantage over balloon/satellite based instruments in the detection of high-energy cosmic-ray electrons.”

# Data sets

Electrons + positrons adding FERMI and HESS:

- no  $e^+ + e^-$  excess
- spectrum  $\sim E^{-3.04}$
- a (smooth) cutoff?




# Results

Which DM spectra can fit the data?

# Results

Which DM spectra can fit the data?


$\mu^+ \mu^-$ ,  $M_{\text{DM}} \simeq 1 \text{ TeV}$




# Results

Which DM spectra can fit the data?

$\mu^+ \mu^-$ ,  $M_{\text{DM}} \simeq 1 \text{ TeV}$




$\tau^+ \tau^-$ ,  $M_{\text{DM}} \simeq 2 \text{ TeV}$




# Results

Which DM spectra can fit the data?


$\mu^+ \mu^-$ ,  $M_{\text{DM}} \simeq 1 \text{ TeV}$



$\tau^+ \tau^-$ ,  $M_{\text{DM}} \simeq 2 \text{ TeV}$




$W^+ W^-$ ,  $M_{\text{DM}} \simeq 10 \text{ TeV}$



# Results

Which DM spectra can fit the data?


$\mu^+ \mu^-$ ,  $M_{\text{DM}} \simeq 1 \text{ TeV}$




Notice:

- same spectra still fit PAMELA positron and anti-protons!

$\tau^+ \tau^-$ ,  $M_{\text{DM}} \simeq 2 \text{ TeV}$




$W^+ W^-$ ,  $M_{\text{DM}} \simeq 10 \text{ TeV}$



# Results

Which DM spectra can fit the data?


$\mu^+ \mu^-$ ,  $M_{\text{DM}} \simeq 1 \text{ TeV}$




Notice:

- same spectra still fit PAMELA positron and anti-protons!

$\tau^+ \tau^-$ ,  $M_{\text{DM}} \simeq 2 \text{ TeV}$



$W^+ W^-$ ,  $M_{\text{DM}} \simeq 10 \text{ TeV}$




- no features in FERMI =>  $M_{\text{DM}} > 1 \text{ TeV}$
- a ‘cutoff’ in HESS =>  $M_{\text{DM}} \lesssim 3 \text{ TeV}$
- **smooth** lepton spectrum

# Results

Which DM spectra can fit the data?

Model-independent results:


fit to PAMELA + FERMI + HESS (no balloon):




(1) annihilate into leptons (e.g.  $\tau^+ \tau^-$ ), mass  $\sim 3$  TeV

# Two important remarks

A. Maybe it's just a pulsar,  
or other astrophysics



B. Associated *gamma ray* and  
radio constraints from  
the GC, Gal Halo and  
dwarf galaxies are severe



Bertone, Cirelli, Strumia, Taoso 0811.3744


[jump to conclusions]

# Astrophysical explanation?

# Astrophysical explanation?

[others?]

Or perhaps it's just a **young, nearby pulsar...**




'Mechanism': the spinning  $\vec{B}$  of the pulsar strips  $e^-$  that emit  $\gamma$  that make production of  $e^\pm$  pairs that are trapped in the cloud, further accelerated and later released at  $\tau \sim 0 \rightarrow 10^5$  yr (typical total energy output:  $10^{46}$  erg).

Must be young ( $T < 10^5$  yr) and nearby (< 1 kpc); if not: too much diffusion, low energy, too low flux.


Predicted flux:  $\Phi_{e^\pm} \approx E^{-p} \exp(E/E_c)$  with  $p \approx 2$  and  $E_c \sim \text{many TeV}$

( $1.4 < p < 2.4$ , Profumo 2008)

Not a  
new  
idea:




A.Boulares, APJ 342 (1989)



Atoyan, Aharonian, Volk (1995)

# Astrophysical explanation?

Or perhaps it's just a **young, nearby pulsar...**



Geminga pulsar

(funny that it means:  
“it is not there” in milanese)

‘Mechanism’: the spinning  $\vec{B}$  of the pulsar strips  $e^-$  that emit  $\gamma$  that make production of  $e^\pm$  pairs that are trapped in the cloud, further accelerated and later released at  $\tau \sim 0 \rightarrow 10^5$  yr.


Must be young ( $T < 10^5$  yr) and nearby (< 1 kpc);  
if not: too much diffusion, low energy, too low flux.

Predicted flux:  $\Phi_{e^\pm} \approx E^{-p} \exp(E/E_c)$  with  $p \approx 2$  and  $E_c \sim \text{many TeV}$

Try the fit with known nearby pulsars:


TABLE 1  
LIST OF NEARBY SNRs

SNR	Distance (kpc)	Age (yr)	$E_{\max}^a$ (TeV)
SN 185 .....	0.95	$1.8 \times 10^3$	$1.7 \times 10^2$
S147 .....	0.80	$4.6 \times 10^3$	63
HB 21 .....	0.80	$1.9 \times 10^4$	14
G65.3+5.7 .....	0.80	$2.0 \times 10^4$	13
Cygnus Loop.....	0.44	$2.0 \times 10^4$	13
Vela .....	0.30	$1.1 \times 10^4$	25
Monogem .....	0.30	$8.6 \times 10^4$	2.8
Loop1 .....	0.17	$2.0 \times 10^5$	1.2
Geminga .....	0.4	$3.4 \times 10^5$	0.67



# Astrophysical explanation?

Or perhaps it's just a **young, nearby pulsar...**





Geminga pulsar

'Mechanism': the spinning  $\vec{B}$  of the pulsar strips  $e^-$  that emit  $\gamma$  that make production of  $e^\pm$  pairs that are trapped in the cloud, further accelerated and later released at  $\tau \sim 0 \rightarrow 10^5$  yr.

Must be young ( $T < 10^5$  yr) and nearby ( $< 1$  kpc); if not: too much diffusion, low energy, too low flux.

Predicted flux:  $\Phi_{e^\pm} \approx E^{-p} \exp(E/E_c)$  with  $p \approx 2$  and  $E_c \sim \text{many TeV}$


Try the fit with known nearby pulsars and **diffuse mature pulsars**:



Hooper, Blasi, Serpico 2008

# Astrophysical explanation?

Or perhaps it's just a **young, nearby pulsar...**





Geminga pulsar

'Mechanism': the spinning  $\vec{B}$  of the pulsar strips  $e^-$  that emit  $\gamma$  that make production of  $e^\pm$  pairs that are trapped in the cloud, further accelerated and later released at  $\tau \sim 0 \rightarrow 10^5$  yr.

Must be young ( $T < 10^5$  yr) and nearby (< 1 kpc); if not: too much diffusion, low energy, too low flux.


Predicted flux:  $\Phi_{e^\pm} \approx E^{-p} \exp(E/E_c)$  with  $p \approx 2$  and  $E_c \sim \text{many TeV}$

ATIC needs a different (and very powerful) source:



# Astrophysical explanation?

Or perhaps it's just a **young, nearby pulsar...**





Geminga pulsar

'Mechanism': the spinning  $\vec{B}$  of the pulsar strips  $e^-$  that emit  $\gamma$  that make production of  $e^\pm$  pairs that are trapped in the cloud, further accelerated and later released at  $\tau \sim 0 \rightarrow 10^5$  yr.

Must be young ( $T < 10^5$  yr) and nearby (< 1 kpc); if not: too much diffusion, low energy, too low flux.

Predicted flux:  $\Phi_{e^\pm} \approx E^{-p} \exp(E/E_c)$  with  $p \approx 2$  and  $E_c \sim$  many TeV


PAMELA + FERMI + HESS can be well fitted by pulsars:



D.Grasso et al.  
(sub-FERMI collab.)  
0905.0636

# Astrophysical explanation?

Or perhaps it's just a **young, nearby pulsar...**



Geminga pulsar

'Mechanism': the spinning  $\vec{B}$  of the pulsar strips  $e^-$  that emit  $\gamma$  that make production of  $e^\pm$  pairs that are trapped in the cloud, further accelerated and later released at  $\tau \sim 0 \rightarrow 10^5$  yr.

Must be young ( $T < 10^5$  yr) and nearby ( $< 1$  kpc);  
if not: too much diffusion, low energy, too low flux.

Predicted flux:  $\Phi_{e^\pm} \approx E^{-p} \exp(E/E_c)$  with  $p \approx 2$  and  
 $E_c \sim$  many TeV

**Open issue.**

(look for anisotropies,  
(both for single source and collection in disk)

**antiprotons, gammas...**  
(Fermi is discovering a pulsar a week)


or shape of the spectrum...)

e.g. Yuksel, Kistler, Stanev 0810.2784  
Hall, Hooper 0811.3362


[back]

# Two important remarks

A. Maybe it's just a pulsar,  
or other astrophysics



B. Associated **gamma ray** and  
radio constraints from  
the GC, Gal Halo and  
dwarf galaxies are severe



Bertone, Cirelli, Strumia, Taoso 0811.3744

[jump to conclusions]

# DM detection

direct detection

production at colliders

$\gamma$  from annihil in galactic center  
and from synchrotron emission

HESS, radio telescopes

indirect

$e^+$  from annihil in galactic halo or center

PAMELA, ATIC, Fermi


$\bar{p}$  from annihil in galactic halo or center

$\bar{D}$  from annihil in galactic halo or center

$\nu, \bar{\nu}$  from annihil in massive bodies

# Indirect Detection

## $\gamma$ from DM annihilations in galactic center




$DM \rightarrow W^-, Z, b, \tau^-, t, h \dots \rightsquigarrow e^\mp, \overset{(-)}{p}, \overset{(-)}{D} \dots$  and  $\gamma$

$DM \rightarrow W^+, Z, \bar{b}, \tau^+, \bar{t}, h \dots \rightsquigarrow e^\pm, \overset{(-)}{p}, \overset{(-)}{D} \dots$  and  $\gamma$

# Indirect Detection

## $\gamma$ from DM annihilations in galactic center




$DM \rightarrow W^-, Z, b, \tau^-, t, h \dots \rightsquigarrow e^\mp, \overset{(-)}{p}, \overset{(-)}{D} \dots$  and  $\gamma$

$DM \rightarrow W^+, Z, \bar{b}, \tau^+, \bar{t}, h \dots \rightsquigarrow e^\pm, \overset{(-)}{p}, \overset{(-)}{D} \dots$  and  $\gamma$


# Indirect Detection

$\gamma$  from DM annihilations in galactic center



# Indirect Detection

## $\gamma$ from DM annihilations in Sagittarius Dwarf




$DM \rightarrow W^-, Z, b, \tau^-, t, h \dots \rightsquigarrow e^\mp, \overset{(-)}{p}, \overset{(-)}{D} \dots \text{ and } \gamma$

$DM \rightarrow W^+, Z, \bar{b}, \tau^+, \bar{t}, h \dots \rightsquigarrow e^\pm, \overset{(-)}{p}, \overset{(-)}{D} \dots \text{ and } \gamma$


# Indirect Detection

radio-waves from synchrotron radiation of  $e^\pm$  in GC




# Indirect Detection

radio-waves from synchrotron radiation of  $e^\pm$  in GC




- compute the population of  $e^\pm$  from DM annihilations in the GC
- compute the synchrotron emitted power for different configurations of galactic  $\vec{B}$   
(assuming ‘scrambled’  $B$ ; in principle, directionality could focus emission, lift bounds by  $O(\text{some})$ )



# Indirect Detection

$\gamma$  from Inverse Compton on  $e^\pm$  in halo




- upscatter of CMB, infrared and starlight photons on energetic  $e^\pm$
- probes regions outside of Galactic Center

# Comparing with data

# Gamma constraints


HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.



HESS coll.

# Gamma constraints


HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.



HESS coll.

# Gamma constraints


HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.




HESS coll.

# Gamma constraints

HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.




HESS coll.




# Gamma constraints

HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.




HESS coll.

a)  $M = 10 \text{ TeV}$  into  $W^+W^-$ , Galactic Center  
 $\sigma v_{\text{ann}} = 10^{-23} \text{ cm}^3/\text{sec}$



Data: HESS coll., astro-ph/0610509

b)  $M = 1 \text{ TeV}$  into  $\mu^-\mu^+$ , Galactic Ridge  
 $\sigma v_{\text{ann}} = 10^{-23} \text{ cm}^3/\text{sec}$




Data: HESS coll., astro-ph/0605021

# Gamma constraints


HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.

Moreover: no detection from Sgr dSph => upper bound.




HESS coll.

a)  $M = 10$  TeV into  $W^+W^-$ , Galactic Center  
 $\sigma v_{\text{ann}} = 10^{-23} \text{ cm}^3/\text{sec}$




b)  $M = 1$  TeV into  $\mu^-\mu^+$ , Galactic Ridge  
 $\sigma v_{\text{ann}} = 10^{-23} \text{ cm}^3/\text{sec}$



Data: HESS coll., astro-ph/0605021

# Gamma constraints

Several observations detected radio to IR emission from the Gal Center. The DM signal must not exceed that.




# Gamma constraints

Several observations detected radio to IR emission from the Gal Center. The DM signal must not exceed that.

Davies 1978 upper bound at 408 MHz.

Davies'78




# Gamma constraints

Several observations detected radio to IR emission from the Gal Center. The DM signal must not exceed that.

Davies 1978 upper bound at 408 MHz.


VLT 2003 emission at  $10^{14}$  Hz.

integrate emission over a small angle corresponding to angular resolution of instrument




# Gamma constraints


EGRET and FERMI have measured diffuse  $\gamma$ -ray emission. The DM signal must not exceed that.



FERMI coll.




Data: EGRET coll., Strong et al. astro-ph/0406254



Data: FERMI coll., several talks in 2009

# Gamma constraints


DM DM  $\rightarrow \mu^+ \mu^-$ , NFW profile



The PAMELA  
and ATIC regions  
are in **conflict**  
with gamma  
constraints,  
unless...

Bertone, Cirelli, Strumia, Taoso 0811.3744


# Gamma constraints




Bertone, Cirelli, Strumia, Taoso 0811.3744

# Gamma constraints


DM DM  $\rightarrow e^+ e^-$ , isothermal profile




DM DM  $\rightarrow \mu^+ \mu^-$ , isothermal profile




DM DM  $\rightarrow \tau^+ \tau^-$ , isothermal profile




DM DM  $\rightarrow W^+ W^-$ , isothermal profile



DM DM  $\rightarrow b\bar{b}$ , isothermal profile




DM DM  $\rightarrow t\bar{t}$ , isothermal profile



...not-too-steep profile needed.

# Gamma constraints




Bertone, Cirelli, Strumia, Taoso 0811.3744

...not-too-steep profile needed.

Or: take different boosts here (at Earth, for  $e^+$ ) than there (at GC for gammas).


Or: take ad hoc DM profiles (truncated at 100 pc, with central void..., after all we don't know).

# Inverse Compton $\gamma$ constraints



The PAMELA and ATIC regions are in **conflict** with these gamma constraints, and here...

# Inverse Compton $\gamma$ constraints




Cirelli, Panci 0904.3830

see also: Regis, Ullio 0904.4645

# Cosmology: bounds from reionization

DM particles that fit  
PAMELA+FERMI+HESS  
produce **too many**  
**free electrons:**  
bounds on optical depth  
of the Universe violated  
 $\tau = 0.084 \pm 0.016$  (WMAP-5yr)



see also:  
Huetsi, Hektor, Raidal 0906.4550  
Kanzaki et al., 0907.3985

Cirelli, Iocco, Panci, 0907.0719

# Answers

1. Are we seeing Dark Matter  
in cosmic rays?
  
2. Why  $\gtrsim 300$  new DM models have  
been proposed in one year?

# Answers

1. Are we seeing Dark Matter in cosmic rays?

I don't know, I fear it's unlikely, but maybe...  
Maybe it's a pulsar.

2. Why  $\gtrsim 300$  new DM models have been proposed in one year?

# Answers

1. Are we seeing Dark Matter  
in cosmic rays?

I don't know, I fear it's unlikely, but maybe...  
Maybe it's a pulsar.

2. Why  $\gtrsim 300$  new DM models have  
been proposed in one year?

Because the signals point to a "weird" DM so  
theorists try to reinvent the field:

- DM is heavy
- annihilates into leptons and not anti-protons
- huge cross section (boost? Sommerfeld?)
- must not produce too many gammas

# Answers

1. Are we seeing Dark Matter  
in cosmic rays?

I don't know, I fear it's unlikely, but maybe...  
Maybe it's a pulsar.

2. Why  $\gtrsim 300$  new DM models have  
been proposed in one year?

Because the signals point to a "weird" DM so  
theorists try to reinvent the field:

- DM is heavy
- annihilates into leptons and not anti-protons
- huge cross section (boost? Sommerfeld?)
- must not produce too many gammas

**Upcoming data: Fermi, ATIC-4, Pamela, HESS, AMS-02...**

# Answers

1. Are we seeing Dark Matter  
in cosmic rays?

I don't know, I fear it's unlikely, but maybe...  
Maybe it's a pulsar.

2. Why  $\gtrsim 300$  new DM models have  
been proposed in one year?

Because the signals point to a "weird" DM so  
theorists try to reinvent the field:


- DM is heavy
- annihilates into leptons and not anti-protons
- huge cross section (boost? Sommerfeld?)
- must not produce too many gammas

~~Upcoming data: Fermi, ATIC-4, Pamela, HESS, AMS-02...~~


# DM annihilations: the game

Dark Matter  
annihilations


# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game




# DM annihilations: the game



# DM annihilations: the game



# DM annihilations: the game



# Conclusions

Indirect DM searches are powerful and promising.

# Conclusions

Indirect DM searches are powerful and promising.

The recent **PAMELA** results might be a breakthrough:  
excess in positrons, nothing in anti-protons.

# Conclusions

Indirect DM searches are powerful and promising.

The recent **PAMELA** results might be a breakthrough:  
**excess** in positrons, nothing in anti-protons.

Would anything go with PAMELA? Not at all!

DM must

- annihilate into leptons (e.g.  $\mu^+ \mu^-$ ) or
- annihilate into  $W^+ W^-$  with mass  $\gtrsim 10$  TeV

and you need a huge flux.

# Conclusions

Indirect DM searches are powerful and promising.

The recent **PAMELA** results might be a breakthrough:  
excess in positrons, nothing in anti-protons.

Would anything go with PAMELA? Not at all!

DM must

- annihilate into leptons (e.g.  $\mu^+ \mu^-$ ) or
- annihilate into  $W^+ W^-$  with mass  $\gtrsim 10$  TeV

and you need a huge flux.

**Not your garden variety vanilla DM...**

# Conclusions

Indirect DM searches are powerful and promising.

The recent **PAMELA** results might be a breakthrough:  
excess in positrons, nothing in anti-protons.

Would anything go with PAMELA? Not at all!

DM must

- annihilate into leptons (e.g.  $\mu^+ \mu^-$ ) or
- annihilate into  $W^+ W^-$  with mass  $\gtrsim 10$  TeV

and you need a huge flux.

**Not your garden variety vanilla DM...**

Adding **balloon** data (ATIC, PPB-BETS):

DM must annihilate into  $\mu^+ \mu^-$  and have  $M_{\text{DM}} \simeq 1$  TeV

# Conclusions

Indirect DM searches are powerful and promising.

The recent **PAMELA** results might be a breakthrough:  
excess in positrons, nothing in anti-protons.

Would anything go with PAMELA? Not at all!

DM must

- annihilate into leptons (e.g.  $\mu^+ \mu^-$ ) or
- annihilate into  $W^+ W^-$  with mass  $\gtrsim 10$  TeV

and you need a huge flux.

**Not your garden variety vanilla DM...**

Adding **balloon** data (ATIC, PPB-BETS):

DM must annihilate into  $\mu^+ \mu^-$  and have  $M_{\text{DM}} \simeq 1$  TeV

Adding **FERMI & HESS** data:

DM must annihilate into  $\tau^+ \tau^-$  (?) and have  $M_{\text{DM}} \simeq 2 \div 3$  TeV

# Conclusions

Indirect DM searches are powerful and promising.

The recent **PAMELA** results might be a breakthrough:  
excess in positrons, nothing in anti-protons.

Would anything go with PAMELA? Not at all!

DM must

- annihilate into leptons (e.g.  $\mu^+ \mu^-$ ) or
- annihilate into  $W^+ W^-$  with mass  $\gtrsim 10$  TeV

and you need a huge flux.

**Not your garden variety vanilla DM...**

Adding **balloon** data (ATIC, PPB-BETS):

DM must annihilate into  $\mu^+ \mu^-$  and have  $M_{\text{DM}} \simeq 1$  TeV

Adding **FERMI & HESS** data:

DM must annihilate into  $\tau^+ \tau^-$  (?) and have  $M_{\text{DM}} \simeq 2 \div 3$  TeV

But: **gamma, synchrotron** and **ICS** constraints are severe!

Need a not-too-steep DM profile.

# Conclusions

Indirect DM searches are powerful and promising.

The recent **PAMELA** results might be a breakthrough:  
excess in positrons, nothing in anti-protons.

Would anything go with PAMELA? Not at all!

DM must

- annihilate into leptons (e.g.  $\mu^+ \mu^-$ ) or
- annihilate into  $W^+ W^-$  with mass  $\gtrsim 10$  TeV

and you need a huge flux.

**Not your garden variety vanilla DM...**

Adding **balloon** data (ATIC, PPB-BETS):

DM must annihilate into  $\mu^+ \mu^-$  and have  $M_{\text{DM}} \simeq 1$  TeV

Adding **FERMI & HESS** data:

DM must annihilate into  $\tau^+ \tau^-$  (?) and have  $M_{\text{DM}} \simeq 2 \div 3$  TeV

But: **gamma, synchrotron** and **ICS** constraints are severe!


Need a not-too-steep DM profile.

Future data (PAMELA, FERMI, AMS02...) will be crucial.  
Will it be just some young, nearby **pulsar**?

# **Back up slides**

# The cosmic inventory

Most of the Universe is Dark.



*FAvgQ: what's the difference  
between DM and DE?*

DM behaves like matter

- overall it **dilutes** as volume expands
- **clusters** gravitationally on small scales
- $w = P/\rho = 0$  (NR matter)  
(radiation has  $w = -1/3$ )


DE behaves like a constant

- it does not dilute
- does not cluster, it is prob homogeneous
- $w = P/\rho \simeq -1$
- pulls the acceleration, FRW eq.  $\frac{\ddot{a}}{a} = -\frac{4\pi G_N}{3}(1 - 3w)\rho$

[back]

# DM N-body simulations

$2 \times 10^6$  CDM particles, 43 Mpc cubic box




Andrey Kravtsov, [cosmicweb.uchicago.edu](http://cosmicweb.uchicago.edu)

[back]

# DM N-body simulations

2dF:  $2.2 \cdot 10^5$  galaxies  
SDSS:  $10^6$  galaxies,  
2 billion lyr



Springel, Frenk, White, Nature 440 (2006)


Millennium:  
 $10^{10}$  particles,  
 $500 h^{-1} \text{ Mpc}$

[back]


# The Evidence for DM

How would the power spectra be without DM?  
(and no other extra ingredient)

CMB



LSS



(in particular: no DM  $\Rightarrow$  no 3<sup>rd</sup> peak!)

(you need DM to gravitationally  
“catalyse” structure formation)

[back]


# Indirect Detection

**Boost Factor:** local clumps in the DM halo enhance the density, boost the flux from annihilations. Typically:  $B \simeq 1 \rightarrow 20$  ( $10^4$ )

In principle,  $B$  is different for  $e^+$ , anti-p and gammas,  
energy dependent,


dependent on many astro assumptions (inner density profile of clump, tidal disruptions and smoothing...),  
with an energy dependent variance, at high energy for  $e^+$ , at low energy for anti-p.

positrons



Lavalle et al. 2006

antiprotons



Lavalle et al. 2007

# Indirect Detection

Propagation for positrons:

$$\frac{\partial f}{\partial t} - K(E) \cdot \nabla^2 f - \frac{\partial}{\partial E} (b(E)f) = Q$$

diffusion

(in turbulent  $\bar{B} \approx \mu\text{G}$ ,  
assumed space indep.)

$$K(E) = K_0(E/\text{GeV})^\delta$$

energy loss

$$b(E) = (E/\text{GeV})^2/\tau_E$$

$$\tau_E = 10^{16} \text{ s}$$

$$Q = \frac{1}{2} \left( \frac{\rho}{M_{\text{DM}}} \right)^2 f_{\text{inj}},$$


$$f_{\text{inj}} = \sum_k \langle \sigma v \rangle_k \frac{dN_{e^+}^k}{dE}$$

Model	$\delta$	$K_0$ in $\text{kpc}^2/\text{Myr}$	$L$ in kpc
min (M2)	0.55	0.00595	1
med	0.70	0.0112	4
max (M1)	0.46	0.0765	15

Solution:

$$\Phi_{e^+}(E, \vec{r}_\odot) = B \frac{v_{e^+} \tau_E}{4\pi} \int_E^{M_{\text{DM}}} dE' Q(E') \cdot I(\lambda_D(E, E'))$$

$$\lambda_D^2 = 4K_0 \tau_E \left[ \frac{(E/\text{GeV})^{\delta-1} - (E'/\text{GeV})^{\delta-1}}{\delta - 1} \right]$$



[back]

# Indirect Detection


Where do positrons come from?

Mostly locally, within 1 kpc  
(more so at higher energy).

Typical lifetime (due to syn rad & IC):

$$\tau \approx 5 \cdot 10^5 \text{ yr} \frac{\text{TeV}}{E} \frac{1}{\left(\frac{B}{5\mu\text{G}}\right)^2 + 1.6 \frac{w}{\text{eV/cm}^3}}$$

*(w = density of IS photons)*



T.Delahaye et al., 2008


[back]

# 3. Indirect Detection

Results for positrons:

Astro uncertainties:

- propagation model
- DM halo profile
- boost factor B



[back]


# 3. Indirect Detection

Results for positrons:

Astro uncertainties:

- propagation model
- DM halo profile
- boost factor B

Distinctive signal,  
quite robust vs astro.



[back]

# 3. Indirect Detection

Propagation for antiprotons:

$$\frac{\partial f}{\partial t} - K(T) \cdot \nabla^2 f + \frac{\partial}{\partial z} (\text{sign}(z) f V_{\text{conv}}) = Q - 2h \delta(z) \Gamma_{\text{ann}} f$$

diffusion

$$K(T) = K_0 \beta (p/\text{GeV})^\delta$$

$T$  kinetic energy


convective wind

spallations

Model	$\delta$	$K_0$ in $\text{kpc}^2/\text{Myr}$	$L$ in kpc	$V_{\text{conv}}$ in km/s
min	0.85	0.0016	1	13.5
med	0.70	0.0112	4	12
max	0.46	0.0765	15	5

Solution:

$$\Phi_{\bar{p}}(T, \vec{r}_\odot) = B \frac{v_{\bar{p}}}{4\pi} \left( \frac{\rho_\odot}{M_{\text{DM}}} \right)^2 R(T) \sum_k \frac{1}{2} \langle \sigma v \rangle_k \frac{dN_{\bar{p}}^k}{dT}$$



[back]

# Indirect Detection

Solar wind Modulation of cosmic rays:


$$\frac{d\Phi_{\bar{p}\oplus}}{dT_{\oplus}} = \frac{p_{\oplus}^2}{p^2} \frac{d\Phi_{\bar{p}}}{dT},$$

spectrum  
at Earth

spectrum  
far from Earth

$$T = T_{\oplus} + |Ze|\phi_F$$


Fisk  
potential  $\phi_F \simeq 500$  MV



# Indirect Detection

Solar polarity Modulation of cosmic rays:

solar magnetic polarity reverses at (the max of) each cycle;  
during ‘- polarity’ state, positive particles are more deflected away



+ = rotation parallel  
to magnetic field;  
- = antiparallel

# Indirect Detection

Background computations for positrons:


$$\Phi_{e^+}^{\text{bkg}} = \frac{4.5 E^{0.7}}{1 + 650 E^{2.3} + 1500 E^{4.2}}$$

main source: CR nuclei  
spallating on IS gas

$$\Phi_{e^-}^{\text{bkg}} = \Phi_{e^-}^{\text{bkg, prim}} + \Phi_{e^-}^{\text{bkg, sec}} = \frac{0.16 E^{-1.1}}{1 + 11 E^{0.9} + 3.2 E^{2.15}} + \frac{0.70 E^{0.7}}{1 + 110 E^{1.5} + 580 E^{4.2}}$$

Baltz, Edsjo 1999

On the basis of CR simulations of  
Moskalenko, Strong 1998



More recently:  
Delahaye et al., 0809.5268  
P.Salati, Cargese 2007


We marginalize w.r.t. the slope  
 $E^p$ ,  $p = \pm 0.05$   
and let normalization free.

[back]

# Indirect Detection

Background estimation for positrons:

using new  
measurements of  
electron fluxes  
Casadei, Bindi 2004



T.Delahaye et al., 0809.5268


[back]

# Indirect Detection

## Background estimation for positrons:

relaxing the assumption of isotropy\* in propagation model ( $\Lambda$ CDM = anisotropic convection driven transport model), allows to fit PAMELA with pure background

\* (ROSAT X-ray satellite has seen fast, strong SN winds coming out from galaxy plane: not isotropic)



Gebauer 0811.2767


# Indirect Detection

Background estimation for positrons:

SNRs in the spiral arm as sources of electrons (not positrons), whose flux drops at 10 GeV for energy loss

= PAMELA

additional more local SNRs inject  
further electrons at 100 GeV = ATIC



# Indirect Detection


Background estimation for positrons:

SNRs in the spiral arm as sources of electrons (not positrons), whose flux drops at 10 GeV for energy loss

= PAMELA

additional more local SNRs inject further electrons at 100 GeV = ATIC



**But:** preliminary PAMELA data on absolute  $e^-$  flux show harder spectrum ( $E^{-3.33}$ ) than this prediction...;  
do nearby sources agree with B/C...?



# Indirect Detection

Background computations for antiprotons:

$$\log_{10} \Phi_{\bar{p}}^{\text{bkg}} = -1.64 + 0.07 \tau - \tau^2 - 0.02 \tau^3 + 0.028 \tau^4 \quad \tau = \log_{10} T/\text{GeV}$$



We marginalize w.r.t. the slope  
 $E^p$ ,  $p = \pm 0.05$   
and let normalization free.


[back]

# Indirect Detection

Results for anti-protons:

Astro uncertainties:

- propagation model
- DM halo profile
- boost factor B




# Indirect Detection

Results for anti-protons:

Astro uncertainties:

- propagation model
- DM halo profile
- boost factor B



# Challenges for the 'conventional' DM candidates

Needs:

	SuSy DM	KK DM
- TeV or multi-TeV masses	difficult	ok
- no hadronic channels	difficult	difficult
- no helicity suppression	no	ok

 for any Majorana DM,  
s-wave annihilation cross section

$$\sigma_{\text{ann}}(\text{DM } \bar{\text{DM}} \rightarrow f\bar{f}) \propto \left( \frac{m_f}{M_{\text{DM}}} \right)^2$$

# Results

Which DM spectra can fit the data?


Ok, let's *insist* on Wino with: -mass  $M_{\text{DM}} = 200 \text{ GeV}$   
-annihilation  $\text{DM DM} \rightarrow W^+W^-$

If one:


- assumes non-thermal production of DM
- takes positron energy loss 5 times larger than usual
- takes "min" propagation only
- gives up ATIC
- neglects conflict with EGRET bound (4 times too many gammas)

then:

Positrons:



Anti-protons:



G.Kane, A.Pierce, P.Grajek, D.Phalen, S.Watson 0812.4555

# Results

Which DM spectra can fit the data?

Ok, let's *insist* on KK DM with:

-mass  $M_{\text{DM}} = 600 - 800 \text{ GeV}$

-annihilation  $\text{DM DM} \rightarrow l^+l^- (BR = 60\%)$   
 $\text{DM DM} \rightarrow q\bar{q} (BR = 35\%)$


Good fit with: - boost  $B = 1800$

- propagation model


$$B: K(E_e) = 1.4 \times 10^{28} (E_e/4 \text{ GeV})^{0.43} \text{ cm}^2/\text{s}, L=1 \text{ kpc}$$

very large energy loss with very small L


Positrons:



Electrons + Positrons:




Anti-protons:



# Data sets

Electrons + positrons from Fermi-LAT:


Fermi detects gammas by pair production: it's inherently an  $e^+e^-$  detector



[back]

# Results

Which DM spectra can fit the data?



# Astrophysical explanation?

see S.Profumo, 0812.4457

the electron spectrum has a steep deepening!

**T.Delahaye et al., 09.2008**

Casadei, Bindi 2004

**Tsvi Piran et al., 0902.0376**

- difficult to get PAMELA slope?
- does it explain ATIC or HESS?

CR proton collisions on giant molecular clouds produce  $e^+e^-$ !

Dogiel, Sharov 1990

- does not work at  $E > 30$  GeV

Coutu et al (HEAT), 1990

Gamma Ray Bursts produce  $e^+e^-$ !

Ioka 0812.4851

- maybe, constrained by gammas

$\beta^+$  decays of  $^{56}\text{Co}$  in SN produce  $e^+$ !

ICRC 1990

- low energy and low flux

...

[back]

# Model building

- Minimal extensions of the SM:  
heavy WIMPS (Minimal DM, Inert Doublet)

Cirelli, Strumia et al. 2005-2009

Tytgat et al. 0901.2556

- More drastic extensions:  
New models with a rich Dark sector

M.Pospelov and A.Ritz, 0810.1502: Secluded DM - A.Nelson and G.Spitzer, 0810.5167: Slightly Non-Minimal DM - Y.Nomura and J.Thaler, 0810.5397: DM through the Axion Portal - R.Harnik and G.Kribs, 0810.5557: Dirac DM - D.Feldman, Z.Liu, P.Nath, 0810.5762: Hidden Sector - T.Hambye, 0811.0172: Hidden Vector - K.Ishiwata, S.Matsumoto, T.Moroi, 0811.0250: Superparticle DM - Y.Bai and Z.Han, 0811.0587: sUED DM - P.Fox, E.Poppitz, 0811.0399: Leptophilic DM - C.Chen, F.Takahashi, T.T.Yanagida, 0811.0477: Hidden-Gauge-Boson DM - E.Ponton, L.Randall, 0811.1029: Singlet DM - S.Baek, P.Ko, 0811.1646: U(1) Lmu-Ltau DM - I.Cholis, G.Dobler, D.Finkbeiner, L.Goodenough, N.Weiner, 0811.3641: 700+ GeV WIMP - K.Zurek, 0811.4429: Multicomponent DM - M.Ibe, H.Murayama, T.T.Yanagida, 0812.0072: Breit-Wigner enhancement of DM annihilation - E.Chun, J.-C.Park, 0812.0308: sub-GeV hidden U(1) in GMSB - M.Lattanzi, J.Silk, 0812.0360: Sommerfeld enhancement in cold substructures - M.Pospelov, M.Trott, 0812.0432: super-WIMPs decays DM - Zhang, Bi, Liu, Liu, Yin, Yuan, Zhu, 0812.0522: Discrimination with SR and IC - Liu, Yin, Zhu, 0812.0964: DMnu from GC - M.Pohl, 0812.1174: electrons from DM - J.Hisano, M.Kawasaki, K.Kohri, K.Nakayama, 0812.0219: DMnu from GC - R.Allahverdi, B.Dutta, K.Richardson-McDaniel, Y.Santoso, 0812.2196: SuSy B-L DM - S.Hamaguchi, K.Shirai, T.T.Yanagida, 0812.2374: Hidden-Fermion DM decays - D.Hooper, A.Stebbins, K.Zurek, 0812.3202: Nearby DM clump - C.Delaunay, P.Fox, G.Perez, 0812.3331: DMnu from Earth - Park, Shu, 0901.0720: Split-UED DM - Gogoladze, R.Khalid, Q.Shafi, H.Yuksel, 0901.0923: cmSSM DM with additions - Q.H.Cao, E.Ma, G.Shaughnessy, 0901.1334: Dark Matter: the leptonic connection - E.Nezri, M.Tytgat, G.Vertongen, 0901.2556: Inert Doublet DM - J.Mardon, Y.Nomura, D.Stolarski, J.Thaler, 0901.2926: Cascade annihilations (light non-abelian new bosons) - P.Meade, M.Papucci, T.Volansky, 0901.2925: DM sees the light - D.Phalen, A.Pierce, N.Weiner, 0901.3165: New Heavy Lepton - T.Banks, J.-F.Fortin, 0901.3578: Pyrma baryons - K.Bae, J.-H. Huh, J.Kim, B.Kyae, R.Viollier, 0812.3511: electrophilic axion from flipped-SU(5) with extra spontaneously broken symmetries and a two component DM with  $Z_2$  parity - ...

- Decaying DM

Ibarra et al., 2007-2009

Nardi, Sannino, Strumia 0811.4153

A.Arvanitaki, S.Dimopoulos, S.Dubovsky, P.Graham, R.Harnik, S.Rajendran, 0812.2075

# Model building

- Minimal extensions of the SM:  
heavy WIMPS (Minimal DM, Inert Doublet)

Cirelli, Strumia et al. 2005-2009

Tytgat et al. 0901.2556

- More drastic extensions:  
New models with a rich Dark sector

M.Pospelov and A.Ritz, 0810.1502: Secluded DM - A.Nelson and C.Spitzer, 0810.5167: Slightly Non-Minimal DM - Y.Nomura and J.Thaler, 0810.5397: DM through the Axion Portal - H.Cai, K.G.Kriba, 0810.5557: Muon DM - M.D.Feldman, Z.Liu, P.Nath, 0810.5762: Hidden Sector - T.Hambye, 0811.0172: Hidden Vector - K.Ishiwata, S.Matsumoto, T.Miyata, 0811.0256: Inert Doublet DM - and Z.Han, 0811.0387: SUED DM - P.Fox, E.Poppitz, 0811.0599: Leptophilic DM - C.Chen, F.Takahashi, T.T.Yanagida, 0811.0477: Hidden-Gauge-Boson DM - E.Farouz, L.Randall, 0811.1029: Singlet DM - S.Bae, P.Ko, 0811.1646: U(1) Lmu-Ltau DM - I.Cholis, G.Dobler, D.Finkbeiner, L.Goodenough, N.Weiner, 0811.3641: 700+ GeV WIMP - K.Zurek, 0811.4429: Multicomponent DM - M.Ibe, H.Murayama, T.T.Yanagida, 0812.0072: Breit-Wigner enhancement - J.A.Costa, 0812.0425: Self-Interacting DM - J.Silk, 0812.0360: Sommerfeld enhancement in cold substructures - K.Pospelov, M.Pohl, 0812.0425: Self-Interacting DM - J.Silk, 0812.0522: Discrimination with SR and IC - Liu, Yin, Zhu, 0812.0964: DMnu from GC - M.Pohl, 0812.1174: electrons from DM - J.Hisano, M.Kawasaki, K.Kohri, K.Nakayama, 0812.0219: DMnu from GC - R.Allahverdi, B.Dutta, K.Richardson, M.Tyutgat, 0812.3126: G.S., R.J. Ellis, C.Hanaguchi, K.Saito, T.T.Yanagida, 0812.2574: Hidden-Flavor DM decays - D.Hooper, A.Stebbins, K.Zurek, 0812.3407: New DM in TC - C.D.Garcia, C.S., S. Saito, 0812.3407: New DM in TC - S. Saito, 0901.2201: D.J.~Gildenhez, R.Khalid, Q.Shafi, H.Yuksel, 0901.3923: cmSSM DM with additions - Q.H.Cao, E.Ma, G.Shaughnessy, 0901.1834: Dark Matter: the leptonic connection - E.Nezri, M.Tytgat, G.Vertongen, 0901.2656: Inert Doublet DM - J.Mardon, Y.Nomura, D.Stolarski, J.Thaler, 0901.2929: Cascade annihilations (light non-abelian new bosons) - P.Meade, M.Papucci, T.Volansky, 0901.2925: DM sees the light - D.Phalen, A.Pierce, N.Weiner, 0901.5136: w/ Lee, 0901.2925: DM sees the light - J.Farina, 0901.5774: Probing a baryons - K.Bae, J.-H. Huh, J.Kim, B.Kyae, R.Viollier, 0812.3511: electrophilic axion from flipped-SO(6) with extra spontaneously broken symmetries and a two component DM with  $Z_2$  parity - ...

- Decaying DM

Ibarra et al., 2007-2009

Nardi, Sannino, Strumia 0811.4153

A.Arvanitaki, S.Dimopoulos, S.Dubovsky, P.Graham, R.Harnik, S.Rajendran, 0812.2075

# The “Theory of DM”


Arkani-Hamed, Weiner, Finkbeiner et al. 0810.0713

0811.3641

Basic ingredients:

- $\chi$  Dark Matter particle, decoupled from SM, mass  $M \sim 700+$  GeV
- $\phi$  new gauge boson (“Dark photon”),  
couples only to DM, with typical gauge strength,  $m_\phi \sim$  few GeV  
- mediates Sommerfeld enhancement of  $\chi\bar{\chi}$  annihilation:  
 $\alpha M/m_V \gtrsim 1$  fulfilled

- decays only into  $e^+e^-$  or  $\mu^+\mu^-$   
for kinematical limit



# The “Theory of DM”


Arkani-Hamed, Weiner, Finkbeiner et al. 0810.0713

0811.3641

Basic ingredients:

- $\chi$  Dark Matter particle, decoupled from SM, mass  $M \sim 700+$  GeV
- $\phi$  new gauge boson (“Dark photon”),
  - couples only to DM, with typical gauge strength,  $m_\phi \sim$  few GeV
  - mediates Sommerfeld enhancement of  $\chi\bar{\chi}$  annihilation:  
 $\alpha M/m_V \gtrsim 1$  fulfilled

- decays only into  $e^+e^-$  or  $\mu^+\mu^-$  for kinematical limit




Extras:


- $\chi$  is a multiplet of states and  $\phi$  is non-abelian gauge boson:  
splitting  $\delta M \sim 200$  KeV (via loops of non-abelian bosons)
- inelastic scattering explains DAMA
- excited state decay  $\chi\chi \rightarrow \chi\chi^* \rightarrow e^+e^-$  explains INTEGRAL

# The “Theory of DM”

Phenomenology:



Meade, Papucci, Volanski  
0901.2925




Mardon, Nomura, Stolarski,  
Thaler 0901.2926

# Variations

(selected)


- ★ pioneering: Secluded DM, U(1) Stückelberg extension of SM

Pospelov, Ritz et al 0711.4866 P.Nath et al 0810.5762



- ★ Axion Portal:  $\phi$  is pseudoscalar axion-like

Nomura, Thaler 0810.5397



- ★ singlet-extended UED:  $\chi$  is KK RNnu,  $\phi$  is an extra bulk singlet

Bai, Han 0811.0387

- ★ split UED:  $\chi$  annihilates only to leptons because quarks are on another brane

Park, Shu 0901.0720

- ★ DM carrying lepton number:  $\chi$  charged under  $U(1)_{L_\mu - L_\tau}$ ,  $\phi$  gauge boson


Cirelli, Kadastik, Raidal, Strumia 0809.2409 Fox, Poppitz 0811.0399 ( $m_\phi \sim$  tens GeV)

- ★ New Heavy Lepton:  $\chi$  annihilates into  $\Xi$  that carries lepton number and

decays weakly ( $\sim$  TeV)

( $\sim$  100s GeV)


Phalen, Pierce, Weiner 0901.3165



.....

“PAMELA did not do in-flight checks of the  $\beta$  rejection rate”

What a *little* dash of protons can do!



PAMELA claims  $\beta$  rejection of  $10^{-5}$ . CAUTION! This is not verified using independent technique in flight.

M.Schubnell, ENTApP workshop CERN, 02.2009

# “PAMELA did do in-flight checks of the $p$ rejection rate”

Method: in the calorimeter, leptons leave all their energy and on the top; protons leave little energy and in the bottom.

## Proton background evaluation (pre-semplifier method)

Rigidity: 20-28 GV







Fraction of charge released along the calorimeter track (left, hit, right)

+

Constraints on:

Energy-momentum match

Shower starting-point




Step 1: use the upper portion of the calorimeter to select electrons only ( $\bar{p}$  negligible)

**Step 2:** shower in lower portion selects **protons only**

Step 3: full analysis  
(see that peak is statistically consistent with  $e^-$  peak of step 1)

# Gamma constraints


HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.



HESS coll.

# Gamma constraints


HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.



HESS coll.

# Gamma constraints


HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.




HESS coll.

# Gamma constraints

HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.




HESS coll.




# Gamma constraints

HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.




HESS coll.

a)  $M = 10 \text{ TeV}$  into  $W^+W^-$ , Galactic Center  
 $\sigma v_{\text{ann}} = 10^{-23} \text{ cm}^3/\text{sec}$




b)  $M = 1 \text{ TeV}$  into  $\mu^-\mu^+$ , Galactic Ridge  
 $\sigma v_{\text{ann}} = 10^{-23} \text{ cm}^3/\text{sec}$



# Gamma constraints


HESS has detected  $\gamma$ -ray emission from Gal Center and Gal Ridge. The DM signal must not exceed that.

Moreover: no detection from Sgr dSph => upper bound.




HESS coll.

a)  $M = 10 \text{ TeV}$  into  $W^+W^-$ , Galactic Center  
 $\sigma v_{\text{ann}} = 10^{-23} \text{ cm}^3/\text{sec}$



Data: HESS coll., astro-ph/0610509


b)  $M = 1 \text{ TeV}$  into  $\mu^-\mu^+$ , Galactic Ridge  
 $\sigma v_{\text{ann}} = 10^{-23} \text{ cm}^3/\text{sec}$



Data: HESS coll., astro-ph/0605021

# Gamma constraints

Several observations detected radio to IR emission from the Gal Center. The DM signal must not exceed that.




# Gamma constraints

Several observations detected radio to IR emission from the Gal Center. The DM signal must not exceed that.

Davies 1978 upper bound at 408 MHz.

Davies'78




# Gamma constraints

Several observations detected radio to IR emission from the Gal Center. The DM signal must not exceed that.

Davies 1978 upper bound at 408 MHz.


VLT 2003 emission at  $10^{14}$  Hz.

integrate emission over a small angle corresponding to angular resolution of instrument




# Gamma constraints


EGRET and FERMI have measured diffuse  $\gamma$ -ray emission. The DM signal must not exceed that.



FERMI coll.



Data: EGRET coll., Strong et al. astro-ph/0406254



Data: FERMI coll., several talks in 2009