

Talk 9.6

Effects of the dose-rate on the radiolysis of water and small protein biomolecules

**A. DANVIN¹, A. NASREDDINE², A. ARNONE¹, C. GALINDO¹, P. PEUPARDIN¹, N. LUDWIG², F. KUNTZ², L. BARTOLUCCI³, D. JARNET³,
N. ARBOR¹, R. BARILLON¹, M. DELNERO¹, Q. RAFFY¹**

¹IPHC, Université de Strasbourg, 23 rue du Lœss, 67037 Strasbourg Cedex, France

²Aerial-CRT, 250 Rue Laurent Fries, 67400 Illkirch-Graffenstaden

³ICANS, 17 Rue Albert Calmette, 67200 Strasbourg

Email: antoine.danvin@iphc.cnrs.fr

Ionizing radiations are known to have important effects on living organisms, as their energy deposition causes a lot of damages. Studying fundamental chemical mechanisms of the effects of ionizing radiations on biomolecules is crucial to have a better understanding of their radiobiological effects. These ionizing radiations can come from radioactive isotopes, or be used in a therapeutic context in radiotherapy to treat cancers by damaging tumoral tissues. Particle therapy, using accelerated ions, is very interesting for its better targeting of tumors compared to classical X-Rays. FLASH radiotherapy, which has been attracting a lot of interest recently, could have a preserving effect towards healthy tissues, using very high dose rates (>40Gy/s) [1]. At this moment, the molecular mechanisms of the FLASH effect are still far from being completely understood.

Our team is developing a systematic study of the chemical effects of water radiolysis species on protein biomolecules, especially under ion irradiation. For this purpose, we compare yields of water radiolysis species (esp. HO[•], e⁻_{aq}, H₂O₂) to that of amino acids, peptides and proteins, in order to study the mechanisms of degradation of the biomolecules. Experiments are performed with ions and lowLET ionizing radiations (X-rays, γ, electrons) for comparison, to identify eventual ion specific mechanisms [2]. The effect of dose-rate on the radiolysis mechanisms are also scrutinized.

Hydroxyle radical (HO[•]) is the most potent chemical species towards biomolecules degradation. It has been quantified using scavenging probes at various dose-rates, from 0.1 to about 2000 Gy/s, under irradiation by 1MeV electrons, showing a significant dose-rate effect. Radiolysis of amino acids and of a small peptide, aspartame, was studied in the very same conditions and correlated to the dose-rate effect on HO[•], as shown in *Figure 1* with the example of the 2,5-dopa, one of the radiolysis products of phenylalanine.




Figure 1. Evolution of G(HO[•]) and G(2,5-dopa) ponderated by G(HO[•]) as a function of the dose-rate under irradiation by 1MeV electrons.

References

- [1] de Kruijff, R.M., FLASH radiotherapy: ultra-high dose rates to spare healthy tissue. *International Journal of Radiation Biology*, 2019, 1–5.
- [2] Ludwig, N et al, Radiolysis of Phenylalanine in Solution with Bragg-Peak Energy Protons. *Radiation Measurements* 2018, 116, 55–59.