



ID de Contribution: 133

Type: ORAL

Effects of the dose-rate on the radiolysis of water and small protein biomolecules

jeudi 30 juin 2022 10:15 (15 minutes)

Ionizing radiations are known to have important effects on living organisms, as their energy deposition causes a lot of damages. Studying fundamental chemical mechanisms of the effects of ionizing radiations on biomolecules is crucial to have a better understanding of their radiobiological effects. These ionizing radiations can come from radioactive isotopes, or be used in a therapeutic context in radiotherapy to treat cancers by damaging tumoral tissues. Particle therapy, using accelerated ions, is very interesting for its better targetting of tumors compared to classical X-Rays. FLASH radiotherapy, which has been attracting a lot of interest recently, could have a preserving effect towards healthy tissues, using very high dose rates ($>40\text{Gy/s}$) [1]. At this moment, the molecular mechanisms of the FLASH effect are still far from being completely understood.

Our team is developing a systematic study of the chemical effects of water radiolysis species on protein biomolecules, especially under ion irradiation. For this purpose, we compare yields of water radiolysis species (esp. $\text{HO}\cdot$, e-aq , H_2O_2) to that of amino acids, peptides and proteins, in order to study the mechanisms of degradation of the biomolecules. Experiments are performed with ions and low-LET ionizing radiations (X-rays, γ , electrons) for comparison, to identify eventual ion specific mechanisms [2]. The effect of dose-rate on the radiolysis mechanisms are also scrutinized.

Hydroxyle radical ($\text{HO}\cdot$) is the most potent chemical species towards biomolecules degradation. It has been quantified using scavenging probes at various dose-rates, from 0.1 to about 2000Gy/s , under irradiation by 1MeV electrons, showing a significant dose-rate effect. Radiolysis of amino acids and of a small peptide, aspartame, was studied in the very same conditions and correlated to the dose-rate effect on $\text{HO}\cdot$, as shown in Figure 1 with the example of the 2,5-dopa, one of the radiolysis products of phenylalanine.

[1] de Kruijff, R.M., FLASH radiotherapy: ultra-high dose rates to spare healthy tissue. *International Journal of Radiation Biology*, 2019, 1–5.

[2] Ludwig, N et al, Radiolysis of Phenylalanine in Solution with Bragg-Peak Energy Protons. *Radiation Measurements* 2018, 116, 55–59.

Auteur principal: DANVIN, Antoine (IPHC / Radiochimie)

Co-auteurs: Dr NASREDDINE, Abbas (Aerial-CRT); ARNONE, Aurélia (IPHC DRS Groupe Radiochimie); GALINDO, Catherine (IPHC UMR7178); PEAPARDIN, Philippe (IPHC); LUDWIG, Nicolas (IPHC); Dr KUNTZ, Florent (Aerial-CRT); BARTOLUCCI, Laurent (ICANS); JARNET, Delphine (ICANS); ARBOR, Nicolas (IPHC-Strasbourg); BAILLON, Rémi (IPHC); DEL NERO, Mirella (CNRS-IN2P3 IPHC); RAFFY, Quentin (IPHC / Radiochimie)

Orateur: DANVIN, Antoine (IPHC / Radiochimie)

Classification de Session: Radiobiological Effects, Epidemiology & Health Impacts