



ID de Contribution: 144

Type: ORAL

Natural Nuclear Reactors: prediction, search, discovery, operation and implications

mardi 28 juin 2022 16:30 (20 minutes)

Natural nuclear reactors operated in Oklo (Gabon) about 2.0 billion years ago [1]. This phenomenon was predicted and led to a systematic search for uranium deposits that went critical. Fifty years ago, just a few years after this search was abandoned, the Oklo phenomenon was discovered by chance. Many elements extracted from the Oklo reactor material still carry clear isotopic signatures of ^{235}U fission, ^{239}Pu production and neutron capture reactions. Isotopic compositions of these elements provided reconstruction of neutron fluence, amount of consumed ^{235}U , and an effective duration of nuclear fission chain reaction that was estimated to last for hundreds of thousands of years [2]. It was not clear, however, whether the reactor was operating continuously or in pulses. One proposed mechanism was based on burning up highly neutron absorbing impurities (RRE and/or boron) [3]. As the strong absorbers were burned up at one edge of the reactor zone and uranium at the other one, the active zone could have shifted along the U-vein making different parts of the natural reactor been active at different times [3]. Another potential self-regulation mechanism could have involved water acted as a neutron moderator. As the temperature of the reactor increases, all unbounded water was converted into steam, reducing neutron thermalisation and shutting down the chain reaction. Only when the reactor cooled down and the water concentration increases again, could the chain reaction resume. A tiny sample from reactor zone RZ-13 kindly provided by Maurice Pagel, Philippe Holliger and François Gauthier Lafaye carried the answer to this and several other questions. Novel analytical techniques LENGAs (Laser Extraction Noble Gas Analyses) and NAUTILUS (NAval Ultra-Trace Isotope Laboratory's Universal Spectrometer, the US Naval Research Laboratory) used for analyses of this 4×3 mm slab revealed:

- * the highest concentration ($\sim 8 \times 10^{17}$ atom/g) of fission Xe ever observed in natural materials [4]
- * cycling operation of RZ-13 of Oklo and self-regulating mechanism [5],
- * lowest $^{235}\text{U}/^{238}\text{U} = 0.3655$ and capture of fission Cs and Ba 5 yr. after the shutdown [6],
- * Al-phosphates and metallic aggregates preserve certain fission products over a geologic time [4, 7].

[1] R. Bodu, H. Bouzigues, N. Morin, and J. Pfiffelmann. C. R. Acad. Sci. Paris, 1972, 275 D, 1731.

[2] J. R. De Laeter and H. Hidaka. Mass Spectrometry Review. 2007, 26, 683–712.

[3] R. Naudet. IAEA-SM-204/41, 1975, 589–601. [4] A. Meshik, K. Kehm and C. Hohenberg. Geochim. Cosmochim. Acta, 2000, 64 1651–1661. [5] A. Meshik, C. Hohenberg, and O. Pravdivtseva. Physical Review Letters, 2004, 93, 18, 182302-1–4. [6] E. Groopman, D. Willingham, A. Meshik and O. Pravdivtseva. PNAS, 2018, 115, 8676–8681. [7] E. Groopman, L. Nittler, D. Willingham, A. Meshik and O. Pravdivtseva. Applied Geochemistry, 2021, 131, 105047.

Auteurs principaux: Prof. MESHNIK, Alex (Washington University in St. Louis); Dr GROOPMAN, Evan (National Institute of Standards and Technology); Dr PRAVDIVTSEVA, Olga (Washington University in St. Louis)

Orateur: Prof. MESHIK, Alex (Washington University in St. Louis)

Classification de Session: From Oklo samples to natural core simulations