

# Optimization of gravitational wave optical followup galaxy targeting

**Ducoin Jean-Grégoire**

In collaboration with :

**Corre David**

**Leroy Nicolas**

TS2020 2019




- 1 Galaxies targeting - Standard approach
- 2 Galaxies targeting - Adding galaxies properties
- 3 Conclusion

# EM follow-up of GW events

## Starting point : GW skymap

Example S190425z :



### search of optical counterpart

- Poor 2D localization (up to some 1000 deg<sup>2</sup>)
- Imprecise distance information (~20% error)

# Galaxies targeting - Standard approach

## Galaxies targeting

Hypothesis : the source is located within a galaxy

- Choice of the catalog, what we need :
    - ▶ all sky
    - ▶ provide distance
    - ▶ completeness compatible with LIGO-Virgo range
- ⇒ GLADE (<http://aquarius.elte.hu/glade/>)  
Constructed (combined and matched) from four existing galaxy catalogs : GWGC, 2MPZ, 2MASS XSC and HyperLEDA. GLADE contains 3,262,883 objects.
- Selection in the catalogue of compatible galaxies for a certain 3D volume : RA, Dec, distance

# Galaxies targeting - Standard approach

How do we use the galaxies ?

We need to define a grade (weight) to put on each galaxy

Standard definition of the grade


We use the 3D probability :

$$P_{pos} = P_{dV} = \frac{P_{pixel}}{Pixel\ area} N_{pixel} e^{-\frac{1}{2} \left( \frac{D_{galaxy} - \mu_{pixel}}{\sigma_{pixel}} \right)^2}$$

Where  $\mu_{pixel}$  ,  $\sigma_{pixel}$  and  $N_{pixel}$  are respectively the mean distance, the standard deviation and the normalization factor of the Gaussian distribution at the given pixel.  $D_{galaxy}$  is the galaxy distance fetch from the catalog.

# Results

Example of tiles obtained, GW170817, FOV =  $20' \times 20'$



- 1 Galaxies targeting - Standard approach
- 2 Galaxies targeting - Adding galaxies properties
  - Previous works
  - Our works
  - Results
- 3 Conclusion

## Upgrading the grade

More complete definition of the grade

Only information available on GLADE

- B,J,H,K Luminosity (not for all galaxies)  
⇒ sufficient to deduce interesting properties from it?

LCOGT grade ⇒ stellar mass

Use the B luminosity as an "indicator of mass"

We then calculate the  $B$ -band luminosity of the galaxy,  $L_B$  (based on the  $B$ -band magnitude and distance provided in the GLADE catalog), and assign it a score

$$S_{\text{lum}} = \frac{L_B}{\sum L_B}, \quad (3)$$

(Arcavi et al. 2017)





The B band is highly sensitive to the galaxy dust attenuation

⇒ We should use near infrared band

⇒ K band is provided by GLADE but :

- K band is still a bit affected by the dust extinction
- only ~67% of the galaxies in the catalog (up to 400Mpc) have K band information

⇒ Utilization of the WISE1 band ( $3.4 \mu\text{m}$ )

## Our works

Cross-match AllWISE and GLADE (400Mpc) :

After all treatment we have **~90%** of the galaxies with WISE1 band

Determination of the stellar mass

From WISE1 band we can determine the stellar mass using a **constant mass to light ratio** : (Kettley et al. 2017)

$$\Upsilon_*^{3.4\mu m} \sim 0.60 M_\odot / L_{\odot, 3.4\mu m}$$


Adding a factor to the grade

As done previously we can now change the grade adding :

$$P_{mass} = \frac{M_{*,galaxy}}{\sum M_{*,galaxy}} \Rightarrow P_{tot} = P_{pos} \times P_{mass}$$

# Results

## Comparison with LCOGT method :



A NGC4993 like galaxy ( $\sim 7.2 \times 10^9 L_{\odot}$  B band luminosity) can have a stellar mass which can span from  $\sim 3.8 \times 10^7 M_{\odot}$  to  $\sim 1.0 \times 10^{12} M_{\odot}$

⇒ Our grade is going to behave very differently from one using B band luminosity  
 ⇒ B band luminosity is a very poor indicator of the stellar mass (assuming our determination of stellar mass is ok)

# Reformulation

$$P_{tot} = P_{pos} \times P_{mass}$$

Huge drawback of the product expression :

Can't define  $P_{mass}$  when you don't have the stellar mass info (= the W1 mag)

⇒ forced to throw away **~10%** of the catalog

We chose to reformulate the grade :

$$P_{tot} = P_{pos} (1 + \alpha P_{mass})$$

# Reformulation

$$P_{tot} = P_{pos}(1 + \alpha P_{mass})$$

whit  $\alpha$  that ensure the two factor in the addition are, in mean, contributing as much :

$$\frac{\sum P_{pos}}{N} = \frac{\sum P_{pos} \alpha P_{mass}}{N}$$

$$\Rightarrow \alpha = \frac{\sum P_{pos}}{\sum P_{pos} P_{mass}}$$


$\Rightarrow$  Put  $P_{mass} = 0$  to fall back on  $P_{pos}$

# Results




## GW170817

Only counterpart for a GW found at the moment

Lightcurve from *Fermi*/GBM (50 – 300 keV)




Gravitational-wave time-frequency map



⇒ Mandatory to test our grade on it

# Results

## GW170817



- 90% skymap  $\sim 30 \text{deg}^2$
- distance  $40 \pm 8 \text{ Mpc}$
- 65 galaxies compatibles

## Results

With the standard 3D localization  $\Rightarrow$  NGC 4993 ranked 5



Rank	RA	Dec	$P_{loc}$	Distance	Galaxy name
1	196.27	-22.38	0.0604	33.71	ESO575-053
2	196.88	-23.17	0.0588	38.04	PGC803966
3	194.26	-17.39	0.0588	25.94	WINGSJ125701.38-172325.2
4	197.13	-23.35	0.047	43.15	ESO508-014
5	197.45	-23.38	0.0465	39.35	NGC4993
6	197.18	-23.78	0.0464	36.55	PGC797164
7	196.72	-22.84	0.0452	30.51	ESO508-004
8	197.02	-23.8	0.0403	41.06	IC4197
9	197.47	-24.24	0.0397	39.47	ESO508-019
10	197.69	-23.87	0.0385	38.73	2MASS 13104593-2351566
11	196.89	-23.82	0.0369	41.12	796755
12	196.77	-23.68	0.0364	40.49	NGC4968
13	197.32	-24.38	0.0338	35.5	6dFJ1309178-242256
14	196.91	-23.58	0.0334	45.19	ESO508-010
15	196.06	-22.88	0.0314	36.56	PGC169663
16	196.74	-23.92	0.027	40.67	IC4180
17	193.71	-16.05	0.024	47.39	PGC043966
18	196.35	-23.52	0.0208	37.17	PGC799951
19	194.26	-17.39	0.0206	20.21	WINGSJ125701.40-172325.3
20	197.33	-24.38	0.0196	28.99	ESO508-015
21	197.69	-23.87	0.0191	29.06	ESO508-024
22	194.0	-19.27	0.019	39.95	ESO575-029
23	196.35	-23.5	0.0172	42.35	PGC169670
24	198.19	-25.99	0.017	39.19	PGC772879
25	196.89	-24.01	0.0169	47.07	NGC4970
26	194.26	-17.39	0.015	18.87	WINGSJ125701.40-172325.3
27	194.37	-19.69	0.0108	48.87	NGC4830
28	193.11	-15.52	0.0105	50.9	PGC043664
29	197.06	-21.0	0.0104	31.45	ESO575-061
30	193.84	-17.1	0.0092	53.88	PGC044023



## Results

With the B luminosity  $\Rightarrow$  NGC 4993 ranked 2



Rank	RA	Dec	$P_{tot}$	Distance	Galaxy name
1	197.02	-23.8	0.1371	41.06	IC4197
2	197.45	-23.38	0.1112	39.35	NGC4993
3	196.89	-24.01	0.0807	47.07	NGC4970
4	196.77	-23.68	0.0717	40.49	NGC4968
5	197.47	-24.24	0.0638	39.47	ESO508-019
6	196.74	-23.92	0.0634	40.67	IC4180
7	194.37	-19.69	0.0481	48.87	NGC4830
8	196.27	-22.38	0.0432	33.71	ESO575-053
9	194.0	-19.27	0.0423	39.95	ESO575-029
10	197.69	-23.87	0.0403	29.06	ESO508-024
11	196.91	-23.58	0.0398	45.19	ESO508-010
12	192.25	-14.4	0.0283	51.89	IC3799
13	197.69	-23.87	0.0269	38.73	2MASS 13104593-2351566
14	196.72	-22.84	0.0162	30.51	ESO508-004
15	193.71	-16.05	0.016	47.39	PGC043966
16	193.11	-15.52	0.0154	50.9	PGC043664
17	193.36	-17.01	0.0152	56.57	NGC4763
18	197.18	-23.78	0.0144	36.55	PGC797164
19	197.33	-24.38	0.0135	28.99	ESO508-015
20	197.13	-23.35	0.0129	43.15	ESO508-014
21	193.22	-15.41	0.0088	57.42	NGC4756
22	192.52	-14.73	0.0082	55.63	PGC043424
23	196.6	-24.16	0.0074	53.47	ESO508-003
24	199.1	-26.56	0.007	47.93	ESO508-033
25	194.69	-17.54	0.0066	53.76	PGC044500
26	193.83	-14.95	0.0064	43.14	PGC044021
27	194.25	-17.32	0.005	57.9	PGC044234
28	192.83	-14.57	0.0042	55.44	IC3831
29	194.64	-16.8	0.004	52.46	PGC044478
30	193.62	-16.35	0.0037	57.36	PGC043908

# Results

With the stellar mass addition  $\Rightarrow$  NGC 4993 ranked 1



Rank	RA	Dec	$P_{Tot}$	Distance	Galaxy name	Stellar mass
1	197.45	-23.38	0.119	39.35	NGC4993	10.56
2	197.02	-23.8	0.1055	41.06	IC4197	10.57
3	196.77	-23.68	0.0811	40.49	NGC4968	10.48
4	196.89	-24.01	0.0694	47.07	NGC4970	10.8
5	196.74	-23.92	0.0592	40.67	IC4180	10.47
6	194.37	-19.69	0.0535	48.87	NGC4830	10.89
7	196.27	-22.38	0.0467	33.71	ESO575-053	9.68
8	196.91	-23.58	0.0326	45.19	ESO508-010	9.92
9	196.88	-23.17	0.0296	38.04	PGC803966	7.84
10	194.26	-17.39	0.0294	25.94	WINGSJ125701.38-172325.2	-
11	197.18	-23.78	0.0251	36.55	PGC797164	8.87
12	197.13	-23.35	0.0246	43.15	ESO508-014	8.61
13	196.72	-22.84	0.0233	30.51	ESO508-004	8.42
14	197.47	-24.24	0.0222	39.47	ESO508-019	9.01
15	197.69	-23.87	0.0192	38.73	2MASS 13104593-2351566	-
16	196.89	-23.82	0.0187	41.12	796755.0	7.97
17	197.32	-24.38	0.0171	35.5	6dFJ1309178-242256	8.08
18	196.06	-22.88	0.0159	36.56	PGC169663	8.18
19	194.0	-19.27	0.0129	39.95	ESO575-029	9.5
20	192.52	-14.73	0.0126	55.63	PGC043424	11.06
21	193.71	-16.05	0.0125	47.39	PGC043966	8.58
22	196.35	-23.52	0.0106	37.17	PGC799951	8.26
23	193.36	-17.01	0.0105	56.57	NGC4763	10.74
24	194.26	-17.39	0.0103	20.21	WINGSJ125701.40-172325.3	-
25	197.33	-24.38	0.0099	28.99	ESO508-015	7.78
26	193.11	-15.52	0.0098	50.9	PGC043664	9.89
27	197.69	-23.87	0.0095	29.06	ESO508-024	-
28	196.35	-23.5	0.0088	42.35	PGC169670	8.44
29	198.19	-25.99	0.0086	39.19	PGC772879	7.87
30	199.1	-26.56	0.0083	47.93	ESO508-033	10.16

# Results

With the stellar mass addition  $\Rightarrow$  NGC 4993 ranked 1

Rank	RA	Dec	$P_{Tot}$	Distance	Galaxy name	Stellar mass
1	197.45	-23.38	0.119	39.35	NGC4993	10.56
2	197.02	-23.8	0.1055	41.06	IC4197	10.57
3	196.77	-23.68	0.0811	40.49	NGC4968	10.48
4	196.89	-24.01	0.0694	47.07	NGC4970	10.8
5	196.74	-23.92	0.0592	49.67	IC4189	10.47
6						
7						
8						
9						
10						
11						
12						
13						
14						
15						
16						
17						
18						
19						
20						
21						
22						
23						
24						
25						
26	193.11	-15.52	0.0098	50.9	PGC043664	9.89
27	197.69	-23.87	0.0095	29.06	ESO508-024	-
28	196.35	-23.5	0.0088	42.35	PGC169670	8.44
29	198.19	-25.99	0.0086	39.19	PGC772879	7.87
30	199.1	-26.56	0.0083	47.93	ESO508-033	10.16



- 1 Galaxies targeting - Standard approach
- 2 Galaxies targeting - Adding galaxies properties
- 3 Conclusion**

## Conclusion

# A brand new galaxy selection !



- This method is already used by Kilonova-Catcher and will be then by GRANDMA and SVOM
- Paper about this development is on writing process
- All the tools will be publicly available via dedicated web-site

MERCI!

# Backup slide

$$P_{tot} = P_{pos} (1 + \alpha \beta P_{mass})$$

- ⇒  $\beta$  which will determine at which point the mass factor will count in the grade
- ⇒  $\beta$  is skymap independent
- ⇒  $\beta$  should be to fit with a statistically significant sample of gravitational wave host galaxies, but as we don't have such sample yet ⇒  $\beta = 1$