



ID de Contribution: 430

Type: oral contribution

Towards in-gas-jet studies of isomeric $^{229}\text{Th}^+$

mardi 6 juin 2023 16:00 (15 minutes)

Short half-lives, low production rates and the need to produce them by fusion-evaporation reactions all complicate laser spectroscopy studies of (trans)actinides. The In-Gas Laser Ionization and Spectroscopy (IGLIS) technique has been successfully employed in studies on short-lived actinides (see for instance [1,2]). The addition of a convergent-divergent (de Laval) nozzle to create a cold hypersonic gas jet combines efficiency with sub-GHz spectral resolution. The new generation of nozzles with a Mach number of 8 enables laser spectroscopy studies of actinides with spectral resolutions around 200 MHz [3].

The light actinide ^{229}Th and its nuclear clock isomer have attracted significant attention in the last years. A remarkable feature is the suggested short half-life (< 10 ms) of the isomer in its, not-yet observed, singly charged state [4]. We report on the design of a fast-extraction gas cell (evacuation time of ~ 1 ms) and tailor-made recoil ion sources of ^{233}U prepared by TU Vienna and JGU Mainz which are installed inside the gas cell to provide the isomeric thorium ions. A new set of de Laval nozzles was designed and characterized to operate under the required low-stagnation-pressure conditions of the recoil sources as well as for spectroscopy studies of (trans)actinides in the JetRIS experiment at GSI [5]. A level search above the second ionization potential of thorium revealed several auto-ionizing states which are used to improve laser ionization efficiency for future in-gas-jet laser spectroscopy studies of $^{229\text{m}}\text{Th}^+$.

- [1] C. Granados et al. Phys. Rev. C, 96:054331, 2017.
- [2] S. Raeder et al. Phys. Rev. Lett., 120:232503, 2018.
- [3] R. Ferrer et al. Physical Review Research, 3:043041, 2021.
- [4] L. von der Wense et al. Nature, 533:47–51, 2016.
- [5] S. Raeder et al. Nucl. Instrum. Methods Phys. Res. B, 463:272–276, 2020.

Author: CLAESSENS, Arno (KU Leuven)

Co-auteurs: M. DRAGON, Andreas (Johannes Gutenberg University Mainz, 55099 Mainz, Germany; Helmholtz-Institut Mainz, 55099 Mainz, Germany); DE ROUBIN, Antoine (KU Leuven, Instituut voor Kern- en Stralingsfysica, 3001 Leuven, Belgium); DÜLLMANN, Christoph E. (Johannes Gutenberg University Mainz, 55099 Mainz, Germany, Helmholtz Institute Mainz, 55099 Mainz, Germany, GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany); RENISCH, Dennis (Department of Chemistry – TRIGA Site, JGU Mainz, 55099 Mainz, Germany; Helmholtz-Institute Mainz, 55099 Mainz, Germany); IVANDIKOV, Fedor (IKS, KU Leuven, Belgium); ROMANS, Jekabs (KU Leuven); M. VAN DEN BERGH, Paul (KU Leuven, Instituut voor Kern- en Stralingsfysica, Celestijnenlaan 200D, 3001 Leuven, Belgium); Prof. VAN DUPPEN, Piet (KU Leuven, Instituut voor Kern- en Stralingsfysica, Celestijnenlaan 200D, 3001 Leuven, Belgium); CHHETRI, Premaditya (Instituut voor Kern- en Stralingsfysica, KU Leuven); FERRER, Rafael (KU Leuven - IKS); M. KRAEMER, Sandro (Instituut voor Kern- en Stralingsfysica, KU Leuven); BARA, Silvia (IKS, KU Leuven, Belgium); SCHUMM, Thorsten (Atominstytut, TU Wien); Dr ROSECKER, Veronika (Atominstytut, TU Wien, 1020 Vienna, Austria); M. ELSKENS, Yens (KU Leuven, Instituut voor Kern- en Stralingsfysica, 3001 Leuven, Belgium); KUDRYAVTSEV, Yuri (Instituut voor Kern- en Stralingsfysica, KU Leuven)

Orateur: CLAESSENS, Arno (KU Leuven)

Classification de Session: parallel session

Classification de thématique: facilities/instruments