Advances in Radioactive Isotope Science



ID de Contribution: 313

Type: invited presentation

High-resolution in-beam γ -ray spectroscopy at RIBF

mercredi 7 juin 2023 14:55 (25 minutes)

In-beam gamma-ray spectroscopy experiments have been actively performed at Radioactive Isotope Beam Factory (RIBF) of the RIKEN Nishina Center owing to the high secondary beam intensities from the BigRIPS fragment separator [1]. These experiments mostly employed the DALI2 NaI array with the very high γ -ray detection efficiency [2], and the Zero Degree Spectrometer [1] and the SAMURAI spectrometer [3] for reaction product identifications. Among the abundant scientific achievements from in-beam γ -ray spectroscopy experiments, the first spectroscopy of ⁵⁴Ca [4], ⁷⁸Ni [5], and ⁷⁰Kr [6] are examples of the capability of the RIBF facility.

Despite the notable accomplishments, experiments have been mostly limited to even-even nuclei in the vicinity of shell closures due to the moderate energy resolution of the DALI2 array. For the new capability in spectroscopy, a germanium-based high-resolution γ -ray detector array was constructed in 2019 under the High-resolution Cluster Array at RIBF (HiCARI) project [7]. The HiCARI array was comprised of several different types of high-purity germanium detectors, six segmented triple clusters from the Miniball collaboration, four segmented Clover detectors from the IMP, a quad-type tracking detector from the RCNP, and a triple-cluster tracking detector P3 from the LBNL. Through the improved position and energy resolution, the spectroscopy capabilities could be extended to further regions of interest such as odd-mass and deformed nuclei. Moreover, the HiCARI array was capable to measure level lifetimes based on the line-shape method [8].

In 2020 and 2021, 8 experiments were successfully carried out during 31.5 day of beam times with ²³⁸U and ⁷⁰Zn primary beams. The campaign included a wide range of exotic neutron-rich nuclei covering various physics motivations, such as shell and shape evolutions. An overview of the HiCARI project and first preliminary results from the rich physics program will be presented.

- [1] T. Kubo et al., Prog. Theor. Exp. Phys. 2012, 03C003 (2012).
- [2] S. Takeuchi et al., Nucl. Instrum. And Methods Phys. Res. A 763, 596 (2014).
- [3] T. Kobayashi et al., Nucl. Instrum. And Methods Phys. Res. B 317, 294 (2013).
- [4] D. Steppenbeck et al., Nature 502, 207 (2013).
- [5] R. Taniuchi et al., Nature 569, 52 (2019).
- [6] K. Wimmer et al., Phys. Lett. B 785, 441 (2018).
- [7] K. Wimmer et al., RIKEN Accel. Prog. Rep. 54, S27 (2021).
- [8] P. Doornenbal et al., Nucl. Instrum. Methods Phys. Res. A 613, 218 (2010).

Author: MOON, Byul (Center for Exotic Nuclear Studies, Institute for Basic Science)

Co-auteurs: Dr DOORNENBAL, Pieter (RIKEN Nishina Center); Dr WIMMER, Kathrin (GSI Helmholtzzentrum für Schwerionenforschung GmbH)

Orateur: MOON, Byul (Center for Exotic Nuclear Studies, Institute for Basic Science)

Classification de Session: plenary 11

Classification de thématique: spectroscopy