ID de Contribution: 114

Type: Non spécifié

## Two-quasiparticle states in some odd-odd heavy deformed nuclei within a microscopic approach

mercredi 1 février 2023 11:20 (20 minutes)

We study low-lying states of two-quasiparticle character in some well-deformed odd-odd nuclei around <sup>178</sup>Hf within the framework of the Skyrme energy-density functional (SEDF) approach, including BCS pairing correlations with selfconsistent blocking. We use the SIII SEDF parametrization with time-odd terms and seniority pairing residual interaction as in a previous study of two-quasiparticle *K*-isomeric states in actinide and heavier nuclei [to be published in Phys. Rev. C]. The strength of the seniority interaction is determined through an overall fit on the first  $2^+$  excitation energies [Phys. Rev. C \textbf{99}, 064306 (2019)]. Axial and parity symmetries are assumed throughout the Hartree–Fock–BCS calculations but time-reversal symmetry is broken.

After checking the relevance of the SIII single-particle spectrum by comparison with experimental bandhead states of low-lying one-quasiparticle states in odd-mass neighbors of <sup>178</sup>Hf, we calculate two-quasiparticle states in doubly-odd neighboring nuclei for the relevant neutron and proton configurations. A special attention is drawn on the Gallagher–Moskowski splitting and a comparison with experiment and results obtained by Robledo, Bernard and Bertsch [Phys. Rev. C **89**, 021303(R) (2014)] with the Gogny EDF.

**Co-auteurs:** KONTOWICZ, N. (Université de Bordeaux); KOH, Meng Hock (Department of Physics, Faculty of Science, Universiti Teknologi Malaysia); MINKOV, Nikolay (Institute of Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences,); BARTEL, J. (IPHC, UMR 7178, Université de Strasbourg, CNRS); MOLIQUE, H. (IPHC, UMR 7178, Université de Strasbourg, CNRS)

**Orateur:** LUDOVIC, Bonneau (University of Bordeaux, France)

Classification de Session: Session 10: Shapes and shape coexistence