

# CMS: Status and First Results


Martijn Mulders (CERN)  
for the CMS collaboration

La Thuile, March 7, 2010

XLV<sup>th</sup> Rencontres de Moriond

ElectroWeak Interactions and Unified Theories

# Outline




- CMS and status at start-up
- Commissioning with LHC proton-proton collisions
- First CMS publication on LHC collision data

Total weight 12500 t  
Overall diameter 15 m  
Overall length 21.6 m

# The CMS detector

3.8T Superconducting  
Solenoid

Iron Return  
Yoke



All Silicon tracker  
(pixels and micro-strips)

Lead Tungstate Crystal  
EM Calorimeter (ECAL)

Hermetic ( $|\eta| < 5.2$ )  
Hadron Calorimeter (HCAL)


Muon System with high  
redundancy (RPCs,  
Drift Tubes,  
Cathode Strip Chambers)

Pixel  
Tracker  
ECAL  
HCAL  
Muons  
Solenoid coil




# Commissioning with Muons


> 1 billion cosmics



> 1 million beam halo



> 1000 beam splash (\*)



*CMS invested maximum effort to understand detector performance before LHC start-up*

- **Cosmic Runs at Four Tesla (CRAFT)** in Fall 2008 and Summer 2009: two month-long cosmic data taking campaigns → **2x 300M** events with full detector and B field on
- **Beam halo** (mainly in September 2008) → alignment of End Caps
- **Beam splash** (17 in 2008, 1105 in 2009, 51 in 2010) → synchronization of detector, uniformity of response

(\*) LHC sector test dumping beam on collimator 150m away from CMS →  $O(100k)$  muons















# Status in November 2009:

- Experience with **sustained operation** of CMS as an integrated experiment
- Good **alignment** already at start-up
- Improved understanding **magnetic field**
- **Muon reconstruction** studied up to 1 TeV
- This, and more, **documented in 23 papers** submitted to (and now accepted by) JINST

**And:** detector simulation with realistic conditions (mis-alignment, calibrations) ready for LHC start-up → **used without further tuning** in all following results



Dec09 LHC2 - CMS



# LHC re-start

November 21, 2009





# First LHC p-p collisions

First collisions                      23 November  
First stable beams                  6 December  
First 2.36 TeV collisions        14 December

Recorded **85% of delivered luminosity**

Number of collected events:

$3.9 \times 10^5 \approx 10 \mu\text{b}^{-1} @ 900 \text{ GeV}$


$2.0 \times 10^4 \approx 0.4 \mu\text{b}^{-1} @ 2360 \text{ GeV}$

Tracker on, beam background rejected

Fully 'open' trigger



Minimum Bias trigger rate 0.5-15 Hz

*Quick analysis delivered preliminary results within hours/days*







# $\pi^0$ and $\eta$ in ECAL:



$\pi^0 \rightarrow \gamma\gamma$

Only ECAL barrel ( $|\eta| < 1.479$ )  
 $p_T(\gamma) > 300$  MeV  
 $p_T(\pi^0) > 900$  MeV  
shower shape

*No corrections for shower containment, thresholds, energy loss upstream of ECAL  $\rightarrow$  mass is a bit low*



$\eta \rightarrow \gamma\gamma$

Photon pairs in barrel  
 $ET(\gamma) > 400$  MeV;  
 $ET(\eta) > 2.0$  GeV;  
shower shape

*Good agreement data and MC: peak position and S/B*


$\rightarrow$  energy scale in data and MC agree within 2% (even at these low energies!) 8 / 35





# More $\pi^0$ 's

data




→ mass within 2% of known  $\pi^0$  mass (PDG: 135 MeV)

$$\pi^0 \rightarrow \gamma\gamma$$

(more accurate)


Photon pairs in the  
ECAL barrel ( $|\eta| < 1$ )  
 $E(\gamma) > 400$  MeV  
 $E(\pi^0) > 1.5$  GeV

*Monte-Carlo based  
correction of photon  
cluster energy is applied*



$$\pi^0 \rightarrow \gamma\gamma \rightarrow \gamma e^+ e^-$$

(more challenging)



One photon in the ECAL barrel ( $|\eta| < 1.479$ )  
 $ET(\gamma) > 300$  MeV  
Second photon reconstructed as  $e^+e^-$  pair,  
using tracker only  
 $p_T(\pi^0) > 1.5$  GeV





# $V^0$ decays in the Tracker

- Like **photon conversions**, look more generally for neutral particles that decay far away ( $> 1$  cm or so) from primary vertex, to a pair of oppositely charged tracks
- Useful to find weak decays of  **$K^0$  (and  $\Lambda^0$ )** to  $\pi^+\pi^-$  (or  $p\pi^-$ )

**Track requirements:**  $\geq 6$  hits and  $\chi^2/\text{dof} < 5$   
 $d_0/\sigma(d_0) > 0.5$ .

**Vertex requirements:**  $\chi^2/\text{dof} < 7$ ,  $> 15\sigma$  separation from beam spot in radial direction. No daughter track hits  $> 4\sigma$  inside of vertex






# Strange particles in the Tracker

*First  $K$  and  $\Lambda$  peaks  
presented within hours  
after first 900 GeV run  
with magnet on!*


Peak shape and S/B  
agree beautifully  
between data and MC

Momentum scale  
correct to better than  
0.1% (PDG/data and  
data/MC)


→ confirms excellent  
knowledge of B field





# Cascade baryon and $K^*(892)$



**data** *Fit : Gaussian and Breit-Wigner*



*Again: excellent agreement peak position with PDG mass*




[Analysis details in backup slides]





# Towards b-tagging:



*Getting closer to the primary vertex*

Secondary vertex with 4 tracks

Vertex  $\chi^2/\text{ndf} = 1.67 / 5$

Vertex mass: 1.64 GeV/c<sup>2</sup>


Transverse decay length  
significance:  $L_{xy}/\sigma = 0.12 / 0.019 \text{ [cm]} = 6.6$

3D decay length significance:  
 $L_{3D}/\sigma = 0.26 / 0.037 \text{ [cm]} = 7.0$




# B-tagging variables

*basic variables relevant for b-tagging  
are well described by simulation*




Signed 3D impact parameter for tracks, with  $\geq 7$  hits, associated to a jet. Impact parameter with respect to primary vertex.

Secondary vertices with above tracks, after K rejection:  $L_{xy} < 2.5\text{cm}$ ,  $|M_{\text{vtx}} - M_K| > 0.015\text{ GeV}$





Vertex 3D decay length significance




Number of tracks in Vertex



# dE/dx in the Tracker



dE/dx estimated from  
charge deposited in  
silicon tracker hits  
(analog readout)  
used for particle ID at low  
momentum





# Intermediate Summary

- Good understanding **Electro-Magnetic** calorimeter: energy scale for low-pT **photons** correct to **2% level**
- Beautiful performance of the tracker → **Weak** decays confirm momentum scale for low-pT **tracks** (B field) to **0.1% level**:

Mass bias	$K_S$	$\Lambda$	$\Xi^-$	$K^{*+}$	Tuesday
$(\text{mass}_{\text{data}} - \text{mass}_{\text{PDG}}) - (\text{mass}_{\text{MC}} - \text{mass}_{\text{Gen}})$	$-0.37 \pm 0.07$ MeV	$0.04 \pm 0.06$ MeV	$0.0 \pm 0.9$ MeV	$-4.0 \pm 3.1$ MeV	$-0.22 \pm 0.26$ MeV

*Ready for Unification...*

(Rencontres the Moriond **ElectroWeak** and Unified Theories)






# Unification of calorimetry and tracking

(to further improve the capability of CMS to measure ElectroWeak processes and detect potential new signatures of Unified Theories)

- **Particle Flow** approach: link tracks to calorimeter clusters to reconstruct individual **photons**, charged and neutral **hadrons** → to optimize energy resolution and particle ID
- **CMS** is ideally suited:
  - Powerful B field+ tracker
  - EM calorimeter with fine granularity






# Eta-phi view



CMS, December 2009, 2.36 TeV  
Run 124120 / Event 6613074  
Particle Flow Reconstruction






$(\eta, \phi)$  view of a particle-flow reconstructed event. Reconstructed particles are represented as circles with a radius proportional to their  $p_T$ . The direction of the MET computed from all particles is drawn as a solid horizontal straight line. Particle-based jets with  $p_T > 20 \text{ GeV/c}$  are shown as thinner circles representing the extension of the jet in the  $(\eta, \phi)$  coordinates.




# Linking tracks to Calo-clusters


*The goal: improved jet (and MET) resolution, especially at low  $p_T$*



$\Delta R$  between  
tracks  
( $p_T > 1\text{GeV}$ )  
and closest  
linked  
calorimeter  
cluster



# Particle Flow and HCAL calibration



- Compare **calorimeter cluster energy** to **track momentum** (integrated over full tracker acceptance  $|\eta| < 2.4$ )
- Calibration in simulation and data agree to  $1.5 \pm 4\%$
- This implies that **HCAL calibration scale** agrees **within  $\sim 5\%$**



Using the anti-kT ( $R=0.5$ ) jet algorithm

Three kinds of inputs:

- Calorimeter Jets


- Inputs: Calorimeter Towers
- $E_T$  tower thresholds

- Jets-Plus-Tracks (JPT) Jets

- Inputs: Calorimeter Jets, corrected with tracks
- Single-pion calorimeter response map

- Particle-Flow (PF) Jets

- PF candidate particles
- Photons, charged & neutral hadrons



run 124009: evt 10872958


→ Use superior resolution of tracker (at low  $p_T$ ) to improve jet resolution

→ Combine tracking and calorimetry for *all* particles in the event




# Jet $p_T$ and composition





## Calorimeter



## Jets-plus-tracks





## Particle Flow





# Missing $E_T$


- Raw calorimeter missing  $E_T$  is already rather **stable vs time**
  - Investigation of outliers → identification and **cleaning** of 3 types of **noise**: HF (particle hits PMT window), correlated HCAL noise (specific pattern) and occasional ECAL single hot channel:






# Missing $E_T$


Calorimeter



Track-corrected MET



Particle Flow





*Even in this early stage, without final detector calibration, the missing  $E_T$  is well described in simulation, and tails are small !*





# Missing $E_T$ significance

In these events, no real MET (from neutrino's or other invisible particles) is expected, so any observed MET is a measure of the resolution:



$\text{Sum} E_T > 3 \text{ GeV}$

Particle-flow based MET relative resolution is about twice as good as for Calorimeter-only MET

$$\sigma(E_{x,y}^{\text{miss}}) = a \oplus b \sqrt{\Sigma E_T}$$



Particle-flow based MET:

$a = 0.55 \text{ GeV}$

$b = 45\%$



# A barrel Muon






# Di-muon event in the EndCaps




## CMS Experiment at the LHC, CERN

Data recorded: 2009-Dec-14 03:46:50.815379 GMT  
Run: 124120  
Event: 5686693  
Lumi section: 19  
Orbit: 19245141  
Crossing: 51



1 event observed in mass window 2-4 GeV  
S/B between 3.0 and 3.2 GeV  $\sim 16/1$



$p_T(\mu_1) = 3.6$  GeV,  $p_T(\mu_2) = 2.6$  GeV,  $m(\mu\mu) = 3.03$  GeV



# The First CMS physics paper

JHEP02 (2010) 041

FOR DISCUSSION BY SPRINGER

RECEIVED: February 4, 2010

ACCEPTED: February 7, 2010

PUBLISHED: February 10, 2010



BY SPRINGER

RECEIVED: February 4, 2010

ACCEPTED: February 7, 2010

PUBLISHED: February 10, 2010

## Transverse-momentum and pseudorapidity distributions of charged hadrons in pp collisions at $\sqrt{s} = 0.9$ and 2.36 TeV

CMS Collaboration

**ABSTRACT:** Measurements of inclusive charged-hadron transverse-momentum and pseudorapidity distributions are presented for proton-proton collisions at  $\sqrt{s} = 0.9$  and 2.36 TeV. The data were collected with the CMS detector during the LHC commissioning in December 2009. For non-single-diffractive interactions, the average charged-hadron transverse momentum is measured to be  $0.46 \pm 0.01$  (stat.)  $\pm 0.01$  (syst.) GeV/c at 0.9 TeV and  $0.50 \pm 0.01$  (stat.)  $\pm 0.01$  (syst.) GeV/c at 2.36 TeV, for pseudorapidities between  $-2.4$  and  $+2.4$ . At these energies, the measured pseudorapidity densities in the central region,  $dN_{ch}/d\eta|_{|\eta|<0.5}$ , are  $3.48 \pm 0.02$  (stat.)  $\pm 0.13$  (syst.) and  $4.47 \pm 0.04$  (stat.)  $\pm 0.16$  (syst.), respectively. The results at 0.9 TeV are in agreement with previous measurements and confirm the expectation of near equal hadron production in pp and pp collisions. The results at 2.36 TeV represent the highest-energy measurements at a particle collider to date.

**KEYWORDS:** Hadron-Hadron Scattering


**ARXIV EPRINT:** [1002.0621](https://arxiv.org/abs/1002.0621)

<http://www.springerlink.com/content/t35h6211438476k0/>

JHEP02(2010)041

# Charged hadron $dN/d\eta$ and $dN/dp_T$

- Hadron production in soft pp collisions cannot be calculated perturbatively and has to be **measured in data** and **modeled phenomenologically**
- Important for **high-luminosity** LHC runs with **pile-up** and relevant as reference for **heavy ion physics**
- Various processes involved: elastic, single-diffractive, and **non-single-diffractive (NSD)** = double diffractive + non-diffractive → aim to measure the **NSD component**




Very low  $p_T$  = a big challenge for tracking: 0.1 GeV/c in a B field of 3.8T corresponds to a bending radius of  $\sim 8$  cm




# $dN/dp_T$ results

$dN/dp_T$  in bins of eta:



Integral for  $|\eta| < 2.4$




Fitted with the [empirical Tsallis function](#) (exponential at low  $p_T$ , power law at high  $p_T$ ). Integral used for  $dN/d\eta$  particle count (5% correction at low  $p_T$ )

$\langle p_T \rangle = 0.46 \pm 0.01(\text{stat}) \pm 0.01(\text{syst})$  @0.9TeV

$\langle p_T \rangle = 0.50 \pm 0.01(\text{stat}) \pm 0.01(\text{syst})$  @2.36TeV

# Three methods for $dN/d\eta$

*Pixel detector:* 53.3cm long,  
3 layers with radii: 4.4, 7.3, 10.2 cm



$p_T > 30 \text{ MeV}/c$


Clusters per layer  
 $|\eta| < 2$

3 measurements of  $dN/d\eta$

Immune to mis-alignment

Simplest method

Requires noise-free detector




$p_T > 75 \text{ MeV}/c$

2 of 3 pixel layers

$|\eta| < 2$

3 measurements of  $dN/d\eta$

Sensitive to mis-alignment



Over 50% Efficient for  $p_T > 0.1, 0.2, 0.3 \text{ GeV}/c$  for  $\pi, K, p$


Full tracks (pixel and strips)  
 $|\eta| < 2.4$

$dN/d\eta$  and  $dN/dp_T$


Sensitive to mis-alignment

Most complex

# dN/d $\eta$ Results




3 methods give consistent results. Error bars show systematic errors (ranging from 4.4% to 2.4%), excluding common contributions




The 3 CMS methods are averaged. Shaded band indicates systematic error, of which largest part is due to uncertainty in SD/DD contamination (2%). UA5 and CMS results are symmetrized in  $\eta$ . UA5 and ALICE errors are statistical only

# Results: scaling with Energy



Variation of average transverse momentum with center-of-mass energy




Variation of  $dN/d\eta$  with center-of-mass energy.

$dN/d\eta(@2.36\text{TeV})/dN/d\eta(@0.9\text{TeV}) = (28.4 \pm 1.4 \pm 2.6)\%$   
 significantly larger than prediction from PYTHIA & PHOJET tunes used in the analysis 18.4% & 14.5%



# Summary

- Many results from a small sample ( $\sim 10\mu\text{b}^{-1}$ ) of data
- Equivalent to  $< 1/1000$  of a second of LHC data at design luminosity!
- Started commissioning key ingredients for physics analysis, with excellent results so far
- But: still many orders of magnitude away from normal physics operation
- Expect a million times more data ( $\sim 10\text{pb}^{-1}$ ) very soon!







# CONCLUSION

- CMS arrived prepared to first collision data and was ready to quickly analyze the data and to produce physics results
  - We understand our detector: amazing agreement with simulation without further need of tuning, thanks to many years of preparation with test beams and cosmic runs
  - First paper on collision data is published, 5 other papers are in preparation
  - Excellent detector performance shown with high data quality
- Looking forward to (lots of) 7 TeV data
  - Ready to explore the Standard Model in a new energy domain
  - Prepare for searches

*Beam splash Event February 28, 2010*





# More Information:

CMS overview of published and preliminary physics results:  
[http://cms-physics.web.cern.ch/cms-physics/CMS\\_Physics\\_Results.htm](http://cms-physics.web.cern.ch/cms-physics/CMS_Physics_Results.htm)




## 23 CRAFT performance papers submitted to JINST:

09-001	Commissioning and Performance of the CMS Pixel Tracker with Cosmic Rays	<a href="http://arxiv.org/abs/0911.5434">http://arxiv.org/abs/0911.5434</a>
09-002	Commissioning and Performance of the CMS Silicon Strip Tracker with Cosmic Ray Muons	<a href="http://arxiv.org/abs/0911.4996">http://arxiv.org/abs/0911.4996</a>
09-003	Alignment of the CMS Silicon Tracker During Commissioning with Cosmic Ray Particles	<a href="http://arxiv.org/abs/0910.2505">http://arxiv.org/abs/0910.2505</a>
09-004	Performance and Operation of the CMS Electromagnetic Calorimeter	<a href="http://arxiv.org/abs/0910.3423">http://arxiv.org/abs/0910.3423</a>
09-005	Measurement of the muon stopping power of Lead Tungstate	<a href="http://arxiv.org/abs/0911.5397">http://arxiv.org/abs/0911.5397</a>
09-006	Time Reconstruction and Performance of the CMS Electromagnetic Calorimeter	<a href="http://arxiv.org/abs/0911.4044">http://arxiv.org/abs/0911.4044</a>
09-007	CMS Data Processing Workflows During an Extended Cosmic Ray Run	<a href="http://arxiv.org/abs/0911.4842">http://arxiv.org/abs/0911.4842</a>
09-008	Commissioning of the CMS Experiment and the Cosmic Run at Four Tesla	<a href="http://arxiv.org/abs/0911.4845">http://arxiv.org/abs/0911.4845</a>
09-009	Performance of the CMS Hadron Calorimeter with Cosmic Rays and Accelerator Produced Muons	<a href="http://arxiv.org/abs/0911.4991">http://arxiv.org/abs/0911.4991</a>
09-010	Performance study of Barrel CMS Resistive Plate Chambers with Cosmic Rays	<a href="http://arxiv.org/abs/0911.4045">http://arxiv.org/abs/0911.4045</a>
09-011	Performance of the CMS Cathode Strip Chambers with Cosmic Rays	<a href="http://arxiv.org/abs/0911.4992">http://arxiv.org/abs/0911.4992</a>
09-012	Performance of the CMS Drift Tube Chambers with Cosmic Rays	<a href="http://arxiv.org/abs/0911.4855">http://arxiv.org/abs/0911.4855</a>
09-013	Performance of the CMS Level-1 Trigger during Commissioning with Cosmic Rays	<a href="http://arxiv.org/abs/0911.5422">http://arxiv.org/abs/0911.5422</a>
09-014	Performance of CMS Muon Reconstruction in Cosmic-Ray Events	<a href="http://arxiv.org/abs/0911.4994">http://arxiv.org/abs/0911.4994</a>
09-015	Precise Mapping of the Magnetic Field in the CMS Barrel Yoke using Cosmic Rays	<a href="http://arxiv.org/abs/0910.5530">http://arxiv.org/abs/0910.5530</a>
09-016	Alignment of the CMS Muon System with Cosmic-Ray and Beam-Halo Muons	<a href="http://arxiv.org/abs/0911.4022">http://arxiv.org/abs/0911.4022</a>
09-017	Aligning the CMS Muon Chambers with the Muon Alignment System during an Extended Cosmic Ray Run	<a href="http://arxiv.org/abs/0911.4770">http://arxiv.org/abs/0911.4770</a>
09-018	Performance of CMS Hadron Calorimeter Timing and Synchronization using Cosmic Ray and LHC Beam Data	<a href="http://arxiv.org/abs/0911.4877">http://arxiv.org/abs/0911.4877</a>
09-019	Identification and Filtering of Uncharacteristic Noise in the CMS Hadron Calorimeter	<a href="http://arxiv.org/abs/0911.4881">http://arxiv.org/abs/0911.4881</a>
09-020	Commissioning of the CMS High-Level Trigger with Cosmic Rays	<a href="http://arxiv.org/abs/0911.4889">http://arxiv.org/abs/0911.4889</a>
09-022	Performance of the CMS Drift-Tube Chamber Local Trigger with Cosmic Rays	<a href="http://arxiv.org/abs/0911.4893">http://arxiv.org/abs/0911.4893</a>
09-023	Calibration of the CMS Drift Tube Chambers and Measurement of the Drift Velocity with Cosmic Rays	<a href="http://arxiv.org/abs/0911.4895">http://arxiv.org/abs/0911.4895</a>
09-025	Fine Synchronization of the CMS Muon Drift-Tube Local Trigger using Cosmic Rays	<a href="http://arxiv.org/abs/0911.4904">http://arxiv.org/abs/0911.4904</a>





# Primary Vertexing





Primary vertex  
distribution for  
a single run:  
clean Gaussian  
distributions



Resolution estimated  
by splitting vertices in 2  
and comparing fits:



# LHC beam spot



Commissioned method for determination of LHC beam spot, important for:

- Initial guess of interaction point, before primary vertex fit
- As vertex constraint in High Level Trigger

Mean of primary vertex distribution and beam spot positions are consistent






# Lifetime Measurements

**Monte Carlo is simulated with the same conditions as in data.**

- Data and MC are split into bins of  $c\tau$  and a fit for the yield is performed in each bin.
- Divide MC yields by true (exponential) distribution to obtain correction factor.
- Correct data and fit for lifetime.


**PDG:  $89.53 \pm 0.05$  ps**

**CMS:  $89.80 \pm 2.10$  ps**



**PDG:  $263.1 \pm 2.0$  ps**


**CMS:  $271.0 \pm 20$  ps**



**→ accurate tracking and vertex simulation, even outside the beam region**

# Cascade Baryon signal

- ♦ All 3 tracks must have  $\geq 6$  hits and miss primary by  $3\sigma$  (in 3D).
- ♦  $\Lambda^0$  vertex must be separated by  $10\sigma$  radially from beam spot, have  $\chi^2 < 7$ , and track hits no more than  $4\sigma$  inside.
- ♦  $\Lambda^0$  candidates must be within 8 MeV of PDG mass.
- ♦ Constrain  $\Lambda^0$  mass in vertex fit. Fit probability  $> 1\%$ .
- ♦ Data mass  $1322.8 \pm 0.8$  MeV is consistent with PDG value ( $1321.71 \pm 0.07$  MeV).
- ♦ Data width  $4.0 \pm 0.8$  MeV similar to MC ( $3.6 \pm 0.1$  MeV).





# $K^*(892)$ signal

Basic idea: combine  $K_S$  candidates with charged tracks from the primary vertex.

$K_S$  requirements:


- Tracks have  $\geq 6$  hits, normalized  $\chi^2 < 5$ ,  $d_o/\sigma(d_o) > 2$ .
- Vertex is  $> 15\sigma$  from beam spot (radially), does not have track hits  $> 4\sigma$  inside of position, has  $\chi^2 < 7$ .
- $K_S$  3D momentum vector passes  $< 2$  mm of primary.
- Invariant mass within  $20 \text{ MeV}/c^2$  of PDG value.

Pion requirements:


- Normalized  $\chi^2 < 2$  with  $\geq 7$  hits and  $\geq 2$  pixel hits.
- $p_T > 0.5 \text{ GeV}/c$ ,  $|\eta| < 2$ ,  $d_{xy} < 2 \text{ mm}$ ,  $|d_z| < 3 \text{ mm}$ .

# The $K^*(892)$ resonance

data



MC




- Relativistic Breit-Wigner for signal with the width fixed to PDG value.
- Background function:

$$\frac{1}{\left(m^2 - M^2\right)^2 + \Gamma^2 M^2}$$

$$A \left( 1 - \exp \left( \frac{m_K + m_\pi - m}{B} \right) \right)$$


# Photon Conversions



18-fold structure is from cooling pipes  
Smeared by radial resolution  $\sim 0.5\text{cm}$



# Nuclear Interactions




Resolution of the vertex  $\sim 500 \mu\text{m}$

Good agreement between data and MC  
means a good understanding of the material budget



# Electrons




2.5 GeV electron candidate with bremsstrahlung


Low statistics for signal in these data

Comparison with MC performed mainly for background (only 1/3 of electron candidates are electrons, mostly from conversions)

Commissioning will continue in the next run  
Agreement with MC is promising



low  $p_T$



Good agreement  
Track and  
Calorimeter cluster

Reconstructed electrons candidates combining two seeding algorithms


- “ecal driven” optimized for W/Z electrons, starting from clusters of energy > 4 GeV
- “tracker driven” more suitable for low  $p_T$  electron and electrons in jets

# Jet Corrections


- Derived from Pythia QCD simulation @ 900 GeV and 2360 GeV
- Derived for and applied to calorimeter jets & particle-flow jets

Jet Energy correction factor is function of jet  $p_T$  and  $\eta$ :

Calorimeter jets









Particle Flow jets






# Inclusive Jet $p_T$ and $\eta$











# Di-jet events: Jet composition









# Missing ET and Sum ET





# Tracking Quality dN/d $\eta$





*Good understanding of tracker performance was crucial to quickly produce final results*

Hits on track

Cluster charge

Vertex distribution  
with no tails – beam  
spot in simulation  
Matched to data





# Event Selection $dN/d\eta$

- Aimed at selecting NonSingleDiffractive events with high efficiency (rejecting a large fraction of SingleDiffractive).

Efficiencies:

- NSD:  $\approx 86\%$

- SD:  $\approx 19\%$

- DD:  $\approx 34\%$

NSD are chosen to minimize effect of model dependence of the corrections and allow comparison with previous experiments

- $\approx 10$  Hz collision rate (pile-up probability  $< 2 \times 10^{-4}$ )

- Event selection common to the 3 methods requiring:


- Trigger level: at least 1 hit in Beam scintillation counters AND coincidence with beam pickups (BPTX)

- $> 3$  GeV total energy on both sides Forward calorimeter (HF)

- Beam halo rejection

- Beam background rejection

- A collision vertex



# dN/d $\eta$ : systematic uncertainties

Table 3: Summary of systematic uncertainties. While the various sources of uncertainties are largely independent, most of the uncertainties are correlated between data points and between the analysis methods. The event selection and acceptance uncertainty is common to the three methods and affects them in the same way. The values in parentheses apply to the  $\langle p_T \rangle$  measurement.

Source	Pixel Counting [%]	Tracklet [%]	Tracking [%]
Correction on event selection	3.0	3.0	3.0 (1.0)
Acceptance uncertainty	1.0	1.0	1.0
Pixel hit efficiency	0.5	1.0	0.3
Pixel cluster splitting	1.0	0.4	0.2
Tracklet and cluster selection	3.0	0.5	-
Efficiency of the reconstruction	-	3.0	2.0
Correction of looper hits	2.0	1.0	-
Correction of secondary particles	2.0	1.0	1.0
Misalignment, different scenarios	-	1.0	0.1
Random hits from beam halo	1.0	0.2	0.1
Multiple track counting	-	-	0.1
Fake track rate	-	-	0.5
$p_T$ extrapolation	0.2	0.3	0.5
Total, excl. common uncertainties	4.4	3.7	2.4
Total, incl. common uncert. of 3.2%	5.4	4.9	4.0 (2.8)



# $dN/d\eta$ : DD/SD/NSD fractions

Table 2: Expected fractions of SD, DD, ND and NSD processes ("Frac.") obtained from the PYTHIA and PHOJET event generators before any selection and the corresponding selection efficiencies ("Sel. Eff.") determined from the MC simulation.

Energy	PYTHIA				PHOJET			
	0.9 TeV		2.36 TeV		0.9 TeV		2.36 TeV	
	Frac.	Sel. Eff.	Frac.	Sel. Eff.	Frac.	Sel. Eff.	Frac.	Sel. Eff.
SD	22.5%	16.1%	21.0%	21.8%	18.9%	20.1%	16.2%	25.1%
DD	12.3%	35.0%	12.8%	33.8%	8.4%	53.8%	7.3%	50.0%
ND	65.2%	95.2%	66.2%	96.4%	72.7%	94.7%	76.5%	96.5%
NSD	77.5%	85.6%	79.0%	86.2%	81.1%	90.5%	83.8%	92.4%




# $dN/d\eta$ : Tsallis Function

$$E \frac{d^3 N_{\text{ch}}}{dp^3} = \frac{1}{2\pi p_T} \frac{E}{p} \frac{d^2 N_{\text{ch}}}{d\eta dp_T} = C(n, T, m) \frac{dN_{\text{ch}}}{dy} \left( 1 + \frac{E_T}{nT} \right)^{-n}$$

Limits:


- exponential at low  $p_T$
- power-law at high  $p_T$

# $dN/d\eta$ : Model Dependence




Corrections based either on PYTHIA or on PHOJET event generators yield the same final result

# Tracking Method




Differential yield of charged hadrons in different  $\eta$  bins (vertically shifted by 4 units). Points fitted with the empirical Tsallis function (exponential at low  $p_T$ , power law at high  $p_T$ ) —————> Integral gives hadron count (a 5% correction)





Use all pixel & strip layers  
 Acceptance ( $|\eta| < 2.4$ ,  $> 50\%$  for  $p_T \approx 0.1, 0.2, 0.3$  for  $\pi, K, p$ )  
 Compatibility with beam spot and primary vertex is required  
 Low fake rate ( $< 1\%$ ) achieved with additional cleaning on cluster shapes  
 Immune to beam background  
 More sensitive to beam spot & alignment

# Tracking Method: efficiency



# Pixel Cluster Counting



Counting clusters of pixel hits in pixel barrel layers (acceptance  $p_T > 30 \text{ MeV}/c$   $|\eta| < 2$ )

Applying a cut on cluster length  $\approx |\sinh(\eta)|$  to eliminate loopers and secondaries (shorter clusters)



Corrections for loopers, weak decays, secondaries

Independent results for the 3 layers agree

Insensitive to detector misalignment, sensitive to beam background



# Tracklets Method



**Tracklets:** pairs of clusters in 2 different pixel barrel layers (acceptance  $p_T > 75 \text{ MeV}/c$   $|\eta| < 2$ )

$|\Delta\eta|$  and  $|\Delta\phi|$  between clusters are used to **select signal from primaries**

**Combinatorial background** is subtracted using  **$\Delta\phi$  sidebands**


**Corrections** are applied for efficiency, secondaries, weak decays

**Less sensitive** to beam background




# $\pi^0$ and $\eta$ in ECAL:

data



MC




$\pi^0 \rightarrow \gamma\gamma$


Photon pairs in the ECAL barrel ( $|\eta| < 1$ )  
 $E(\gamma) > 400$  MeV  
 $E(\pi^0) > 1.5$  GeV

*Monte-Carlo based correction of photon cluster energy is applied*

data



MC







$\eta \rightarrow \gamma\gamma$

Photon pairs in barrel  
 $ET(\gamma) > 400$  MeV;  
 $ET(\eta) > 2.0$  GeV;  
 shower shape

*No corrections applied*

*Good agreement data and MC: peak position and S/B*

# Missing ET







# CRAFT: Calibration

## *An excellent understanding of muon response in calorimeters*

- Hadronic calorimeter: good agreement data and simulation over large momentum range
- Crystal calorimeter: first measurement of muon critical energy in Lead Tungstate:




*For a typical energy deposit of 250 MeV!*





# CRAFT: Muons

<http://arxiv.org/abs/0911.4994>



Efficiencies

Charge mis-ID



<0.01% below 20 GeV/c

~1% at 0.5 TeV

Relative  $p_T$  resolution



<1% below 20 GeV/c

~8% at 0.5 TeV/c



# CRAFT: Alignment


<http://arxiv.org/abs/0910.2505>



Tracking performance evaluated by comparing top and bottom half of cosmic muon, reconstructed independently

- Alignment achieved with CRAFT data gives tracking performance close to MC with perfect alignment
- 16027/16588 (97%) of silicon detector modules aligned
  - 3-4  $\mu\text{m}$  in barrel
  - 3-14  $\mu\text{m}$  in endcap
- Internal alignment barrel muon chambers  $\sim 80 \mu\text{m}$  and positions relative to tracker: 200-700  $\mu\text{m}$

<http://arxiv.org/abs/0911.4022>









# CRAFT: Magnetic Field

<http://arxiv.org/abs/0910.5530>

- **Field in Tracker Volume** mapped by probes in 2006 to excellent precision of  $0.5 \cdot 10^{-4}$
- **Up to inner muon station:** NMR probes and cosmic tracks confirm  $\int B \cdot dl$  to better than 0.1% in the barrel
- **Yoke:** cosmic tracks show field in yoke over-estimated by 20%
  - Too tight boundaries used in finite element model (r,z)
  - New map provided with 3-8% accuracy in barrel yoke (more than sufficient for physics)





# CRAFT: Muons

<http://arxiv.org/abs/0911.4994>

*Muons studied in great detail*

Large momentum range

Relative pT resolution

<1% below 20 GeV/c

~8% at 0.5 TeV/c

In Calorimeters

muon system + tracker combined

