



ID de Contribution: 81

Type: Non spécifié

## Primary scintillation in Xe for electrons and alpha-particles

Xenon scintillation has been widely used in rare event detection experiments, such as neutrinoless double beta decay and dark matter [1,2]. Yet, experimental values on the primary scintillation yield in the absence of recombination remain scarce and dispersed. The mean energy required to produce a vacuum ultraviolet scintillation photon ( $W_{sc}$ ) in gaseous xenon has been measured in the range of 30-120 eV [3]. Lower  $W_{sc}$ -values are often found for alpha particles when compared to electrons produced by gamma or x-rays, being this difference not understood. We carried out a systematic study of the absolute primary scintillation yield in Xe at the atmospheric pressure, using a Gas Proportional Scintillation Counter. A simulation model of the detector's geometric efficiency was developed and benchmarked using waveform-shape analysis of primary and secondary scintillation signals.  $W_{sc}$ -values in the range of 30-50 eV were obtained for gamma and x-rays with energies from 5.9 up to 60 keV, and for ~2-MeV alpha particles. No significant differences were found between alpha particles and electrons.

**Auteurs principaux:** HENRIQUES, Carlos; BERNARDES MONTEIRO, Cristina M (University of Coimbra); GONZÁLEZ-DÍAZ, Diego; TEIXEIRA, J.M.R. (University of Coimbra); SILVA, P.A.O.C. (University of Coimbra); MANO, R.D.P. (University of Coimbra)

**Orateur:** HENRIQUES, Carlos

**Classification de Session:** Fundamental measurements session, chair Fernando Amaro