

## APPEL A PROJETS DIM ACAV<sup>+</sup> – ANNÉE 2020

### ALLOCATIONS DOCTORALES – DOSSIER DE CANDIDATURE

Les dossiers de candidature « Allocation doctorale » devront être envoyés à l'adresse [dim.acav@obspm.fr](mailto:dim.acav@obspm.fr) avant le dimanche 09 février 2020 minuit.



#### Sommaire :

1. Synthèse du projet
2. Données administratives
3. Dossier scientifique :
  - Sujet de l'allocation
  - Plan de recherche et calendrier
  - CV du porteur de projet
  - Liste de publications
4. Avis de la direction du laboratoire
5. Suggestion d'experts extérieurs

#### Préambule :

Les proposants sont invités à préparer la partie administrative de leur dossier avec l'école doctorale, l'université d'inscription et l'établissement gestionnaire pour :

- 1) valider le nom du directeur de thèse
- 2) définir l'université d'inscription
- 3) définir l'établissement gestionnaire du contrat de thèse.

## 1. Synthèse du projet

**Nom - Prénom du porteur :** POPOV Boris

**Laboratoire :** LPNHE

**Axe thématique :** Astrophysique des ondes gravitationnelles et des particules

**Acronyme :** HK-SIDEREAL

**Titre Français du projet:** Hyper-Kamiokande, un observatoire unique d'événements rares dans l'Univers - Préparation de l'expérience

**Titre Anglais du projet:** Preparation of the Hyper-Kamiokande experiment – a unique observatory for rare events in the Universe

### Résumé en Français (10 lignes maximum) :

L'expérience Hyper-Kamiokande (HK) sera construite au Japon par une collaboration internationale. Le détecteur sera constitué de 260000 tonnes d'eau ultra pure, soit cinq fois plus que l'expérience existante Super-Kamiokande (SK). La taille gigantesque de HK lui permettra de détecter un nombre de neutrinos provenant de différentes sources sans précédent, notamment provenant de rayons cosmiques, du Soleil, de supernovae et de faisceaux produit artificiellement par un accélérateur de particules existant. En plus de détecter des neutrinos, la possible désintégration spontanée de protons contenus dans l'eau du détecteur pourrait être mesurée, ce qui serait une découverte révolutionnaire. Les physiciens du LPNHE à Paris vont contribuer à la fois à la construction du détecteur, notamment au système de distribution et synchronisation de l'horloge, ainsi qu'à la calibration des données, la reconstruction des événements et aux analyses de physique.

### Résumé en Anglais (10 lignes maximum) :

The Hyper-Kamiokande (HK) experiment will be built in Japan by an international collaboration. The detector will hold 260000 tonnes of ultra-pure water – more than five times larger compared to the existing Super-Kamiokande (SK). The enormous size of the HK will enable it to detect unprecedented numbers of neutrinos produced by various sources — including cosmic rays, the Sun, supernovae and beams artificially produced by an existing particle accelerator. In addition to catching neutrinos, it will



monitor the water for the possible spontaneous decay of protons in atomic nuclei, which, if observed, would be a revolutionary discovery. Physicists from LPNHE-Paris plan to contribute to both the detector construction via a precise time distribution and synchronization system and to the physics analysis via data calibration and reconstruction.



## 2. Données administratives

## **Porteur du projet :**

## Laboratoire :

#### **Etablissement signataire de la convention:**

**Cet établissement devra s'engager à accueillir un stagiaire pendant une période de deux mois minimum.**

Nom : Sorbonne Université  
Adresse : 4 place Jussieu, 75005 Paris

École Doctorale francilienne de rattachement :

Etablissement d'enseignement supérieur de rattachement de l'École Doctorale:



**Co-financement :**

**Collaborations laboratoires franciliens (obligatoire) :**

SYRTE – Observatoire de Paris  
APC (Memphyno setup)

**Collaborations nationales :**

Laboratoire Leprince-Ringuet – UMR 7638  
CEA-Saclay

**Collaborations internationales :**

Collaboration international Hyper-Kamiokande (15 pays, 75 labos)

**Partenariats industriels (entreprises, PME, start-up...) :**

MICRONIQUE : <https://www.micronique.fr>  
RESA : <http://www.resa-spatiales.com>

### 3. Dossier scientifique à rédiger en Anglais

#### Sujet de thèse (2 pages)

During the last decade the LPNHE-neutrino group has been involved in the T2K experiment which uses the Super-Kamiokande (SK) detector to register (anti)neutrino interactions from neutrino and antineutrino beams produced by the J-PARC accelerator. Recently, with the final approval of the Hyper-Kamiokande (HK) project by the Japanese government and endorsement obtained from the LPNHE Scientific Council, the group has also become involved in the HK experiment. The goal is to enlarge and to complement our current activities with studies of neutrinos from astrophysical sources using data from HK experiment. Indeed, full participation in the HK experiment will open a possibility for our group to study solar, atmospheric, supernovae neutrinos and by performing combined analyses of accelerator neutrino and antineutrino data with measurements of (anti)neutrinos from natural sources.

Contrary to T2K, which had access only to a subset of SK data in a small time window around the neutrino beam spill, the Hyper-Kamiokande experiment has a broad physics program covering many areas of particle and astroparticle physics. Based on the proven technology of (Super-)Kamiokande, its much larger detector volume and additional improvements in key areas like photosensors and near/intermediate detectors make HK a straightforward yet powerful extension of the very successful Japan-based neutrino program.

Right from the start of their history, the large Water Cherenkov detectors have been particularly successful in detecting neutrinos from astrophysical sources. Back in 1987, Kamiokande detected few neutrinos emitted by the famous 1987A supernova (SN) explosion, while in 1998 SK observed, for the first time, flavour oscillations of neutrinos produced in the atmosphere and in the Sun. The former observation opened a new window on neutrino astronomy and other exotic searches such as axion by constraining models describing the SN explosion mechanism, while the later proved the existence of neutrino oscillations predicted about 40 years earlier by Bruno Pontecorvo. More recently, new results from the T2K experiment using the muon neutrino beam from the J-PARC accelerator directed towards the SK detector conclusively showed that muon neutrinos transform to electron neutrinos, discovering appearance of new neutrino type in neutrino oscillations.

As the next-generation Water Cherenkov detector, Hyper-Kamiokande with a fiducial volume 8 times larger than Super-Kamiokande will start data taking by 2027. A new Cherenkov detector of 260 kton of water located 295 km down the muon neutrino beam generated by the J-PARC facility will be equipped with 20,000 20-inches photomultipliers (PMTs) with a large quantum efficiency. In addition, it is planned to install about 5,000 multi photomultipliers (mPMTs) that will enhance the detector capabilities at low energy. Indeed, the main advantage of the mPMTs is their improved timing resolution (from 2.6 ns for the 20-inches PMT to 1.6 ns for the 3-inches ones) allowing a reduction of the dark noise rate and an improvement of the spatial reconstruction efficiency.

HK, thanks to its gigantic mass, will detect thousands of electron antineutrinos (via inverse beta-decay) and electron neutrinos (via elastic scattering) from SN bursts in the galactic center. Using the elastic scattering events, it will be possible to reconstruct the direction towards a SN at a distance of 10 kpc with an accuracy of about 1 degree. The events observed in HK will allow to provide detailed information about the time profile and the energy spectrum to further inspect SN explosion mechanism. In addition, it will be possible to detect neutrinos also from extra-galactic SN explosions. Even for distances of 4 Mpc, we will observe few tenths of neutrinos in HK and, at such distances, one SN is expected every three years. HK will also be able to detect the SN relic neutrinos (SRN) that are neutrinos produced by all SN explosions since the beginning of the universe. Such neutrinos fill the present universe and have a flux of few tens/cm<sup>2</sup>/sec. The observation of SRN would allow understanding how heavy elements have been synthesized in stellar formation.

On top of that, HK will collect a large sample of atmospheric high-energy neutrinos. Such measurements will complement the long-baseline program and joint analyses between beam and atmospheric neutrinos are planned in order to improve the sensitivity to neutrino mass ordering and CP violation in neutrino oscillations. Indeed, the matter effects are rather small for the 295 km baseline of Tokai to HK making the sensitivity to the mass ordering limited, while atmospheric neutrinos emitted on the other side of the Earth and measured at the HK detector have crossed the Earth's core and experienced strong matter effects. Therefore, the HK detector as part of the Japanese long-baseline neutrino program will be extremely useful to further constrain neutrino oscillation parameters.

The success of this experiment relies on the excellent reconstruction of incoming neutrino energies and directions using the PMTs. The detection of SN events strongly depends on the reconstruction of the associated low-energy events and on the synchronisation of HK PMTs and other experiments around the world. Moreover, reconstruction of neutrino interaction vertices in the detector requires an accurate timing determination of the event occurrence. It is therefore essential that an excellent clock distribution system is built for distributing and synchronising time among all the detectors. Preliminary studies indicate that the timing precision should be lower than 1 ns with a maximum jitter of 100 ps rms along with the capability of sending data using this link thanks to a sufficient bandwidth. Several solutions are under consideration at present, but two of them seem to be the most promising. One is based on the CERN White Rabbit (WR) protocol and the other on a custom solution.

World-wide, several detectors currently running or nearing completion are sensitive to a core-collapse supernova neutrino signal in the Milky Way using the so-called SNEWS (SuperNova Early Warning System) network. The neutrino burst signal emerges promptly from a supernova's core, whereas it may take hours for the first photons to be visible. Therefore, the detection of the neutrino burst from the next Galactic supernova can provide an early warning for astronomers. Requiring a coincident signal from several detectors will provide the astronomical community with a very high confidence early warning of the supernova's occurrence.

This thesis will be dedicated to the time synchronisation of the photodetectors within the HK far detector. The impact of synchronisation precision on the low energy physics including SN will be studied and will provide motivated requirements for HK. From this, the implementation and characterization of a suitable solution will be done on several experimental setups in view of later integration. New algorithms based on Machine Learning techniques will be developed and tested for reconstructing low energy Cherenkov rings induced by SN neutrinos. Finally, the HK detector will be integrated into a worldwide network for SN detection.

**Plan de recherche et calendrier (2 pages) incluant une description claire des collaborations**

This PhD thesis will be co-directed by Mathieu Guigue (MdC Sorbonne Université), member of the LPNHE-neutrino group, who has recently obtained a Sorbonne Université “Emergence” grant to work on technical developments towards the Hyper-Kamiokande experiment. The work of the student will also be coordinated with the development on time synchronisation performed by experienced electronics engineer Stefano Russo (PhD Ingénieur de recherche, CNRS) working at LPNHE.

A detailed research plan is as follows.

**First year: studies and development of a time synchronization system for Hyper Kamiokande**

One of the contributions foreseen by the LPNHE group to the Hyper-Kamiokande experiment is related to the communication block and, in particular, to the time synchronization and clock distribution for both the large PMTs and mPMTs since it represents a critical part of the experiment and we do have, in our electronic group, experts on this kind of technology.

The experiment's TDCs or FADCs timing synchronization is crucial for precise measurements of photon arrival and the actual requirement is set to have timing resolution of the photo-sensors at the level of 1 ns with a maximum jitter of 100 ps RMS. The distribution of a clock signal, synchronous with the GPS and a local atomic clock, to all Front-End (FE) nodes plus some signals to align FE's local counters call for a system structured as a data exchange protocol to transfer physics related information or slow control. Several solutions are under consideration at present. One is based on the CERN White Rabbit (WR) protocol and the other on a custom solution.

The **White Rabbit** protocol developed by CERN implements a fully deterministic gigabit ethernet-based network for general-purpose data transfer and sub-nanosecond accuracy time transfer. The link, from the timing distribution point of view, is structured as single-master-many-slaves using 18 ports' network switches and has sufficient bandwidth to allow time synchronization and data transfer including slow control. A White Rabbit node can be also connected inside a standard Ethernet network. A possible architecture for the HK experiment could be constituted by a series of switches, the one defined as master receiving the GPS's signal and distributing the clock to all the others via a daisy chain connection. Each switch of the chain can be connected to a maximum of 8 FEs and the other ports are used as a link to the data acquisition system. From this description it is clear that the White Rabbit represents a very attractive solution for the experiment also considering that the protocol is developed and maintained by CERN and the firmware and the hardware designs are distributed freely.

Another architecture is based on a **custom** design that takes advantage from a very specific configuration mode present in almost all the high speed serializer-deserializer embedded in modern FPGAs. Japanese collaborators are actively developing this mode that allows the realization of a high-speed point-to-point connection between two nodes using SK technology.

Recently, we have established close collaboration with physicists from the SYRTE laboratory at the Paris Observatory. This is very important for the project described above as SYRTE colleagues have already accumulated a significant experience in precise time determination and clock synchronization between different locations. The HK experiment could largely profit from this know-how. This is crucial for the efficient inclusion of HK detector into the SNEWS network. In addition, a neutrino burst alert may be able to serve as a trigger for detectors that are not able to trigger on a supernova signal by themselves, allowing extra data to be recorded. Precise timing information will be used later for physics analysis.

The PhD student will work on the characterization of the time synchronisation in the context of HK in collaboration with SYRTE and technical teams (including Stefano Russo) at LPNHE. The newly designed equipment will then be produced by one of enterprises located in Ile-de-France (e.g. MICRONIQUE, RESA). The PhD student will also perform physics analysis to study the impact of

precise timing on the quality of reconstructed events in the HK detector especially on low-energy SN events. This will require a development of new reconstruction algorithms based on artificial intelligence. Performance of those algorithms can be checked using already accumulated SK data. Moreover, data transfer, calibration and reconstruction scheme will be worked out using the very performant IN2P3 Computing Centre at Lyon. There is a possibility to contribute on the computing aspects by providing storage elements and computing resources to the T2K and HK collaborations, in addition to expertise on the distributed computing using DIRAC, a python-based software framework for distributed computing.

#### **2nd year: Characterization and impact of mPMT on HK low-energy physics**

Another possible contribution to the HK project is related to the **mPMT development and tests**. R&D on an alternative photosensor option based on mPMTs modules is actively being carried out by several countries (mostly Canada, Italy, UK) with the goal of providing up to a half of the photocathode coverage for HK. Their lower dark rate compared to the large 20-inch PMTs are making them great candidates for the detection of low-energy events, in particular increasing the sensitivity of HK to the up-turn of the  ${}^8\text{B}$  spectrum of solar neutrinos, probing the MSW-LMA prediction in the transition region between vacuum and matter-dominated neutrino oscillations. Their design also allows better vertex reconstruction especially close to the water tank walls, increasing the detector fiducial volume. Participation in the mPMT development and tests is particularly appealing for our group as we could profit from the existing KM3NeT expertise, from the developments and tests of the small PMTs for the JUNO experiment and from the presence of the Memphyno water tank at the APC laboratory in Paris to perform underwater tests of the mPMT modules.

The Memphyno setup is equipped with two layers of scintillators allowing the identification of cosmic muons crossing the water tank where the tested prototype sits. Initial tests have already started using a first mPMT prototype from Italy currently installed and taking data in water in the Memphyno setup. More mPMTs prototypes are expected to be produced over the next couple of years, and this facility will be of great use for a first in-situ characterisation of their performances, while using the SU "Emergence" grant recently awarded to develop more detailed characterisation capabilities.

The second year of the PhD program will thus be dedicated to the underwater tests of the mPMT prototypes currently under development using the Memphyno setup at APC. The results obtained with these tests could allow the student to further study the impact of the mPMT on the low-energy physics accessible for the HK detector and further develop appropriate reconstruction algorithms using Machine Learning techniques

#### **3rd year: Combined oscillation analysis using HK atmospheric and beam data**

The last year of the PhD will be dedicated to the extraction of new physics results from a joint analysis of T2K and SK data and to the calculation of the expected sensitivities for HK project from a combination of atmospheric and accelerator data. Indeed, the combination of atmospheric data with neutrino beam data allows to better constrain the matter effects on the neutrino oscillation, thus leading to a better constraint on the neutrino oscillation parameters. The results of this combined analysis and the projected sensitivity of the HK experiment will be presented at international conferences and in peer-reviewed papers.

Finally, starting from February 2023, the student will focus on the preparation of the PhD manuscript, with an expected defence in September 2023.

## CV du directeur de thèse (2 pages maximum)

Boris POPOV

Né en 1968 à Dubna (Russie), de nationalité russe

Directeur de Recherche au CNRS depuis 2013

### Études

1991 : diplômé du Moscow Institute of Physics and Technology.

1998 : Doctorat de l'Université Paris 7

(recherches d'oscillations de neutrinos dans l'expérience NOMAD au CERN)

### Fonctions

1991 : physicien titulaire au département de physique des particules élémentaires du Laboratoire des Problèmes Nucléaires au JINR à Dubna

1999-2006 : responsable du groupe neutrinos du Laboratoire des Problèmes Nucléaires au JINR à Dubna

2007-2012 : CDD haut niveau du CNRS (groupe neutrino du LPNHE-Paris)

2013-aujourd'hui : Directeur de Recherche au CNRS (groupe neutrino du LPNHE)

2019-aujourd'hui : coordinateur du groupe neutrino du LPNHE

### Travaux

**1993-2008** : expérience NOMAD au CERN

développement des programmes de reconstruction des chambres à dérive

coordinateur du groupe de travail sur l'identification des électrons

recherche des oscillations  $\nu_\mu \rightarrow \nu_l$  dans le canal  $\tau \rightarrow e\nu_e\nu_l$  (thèse de Doctorat)

mesure de la production de particules étranges et de la polarisation des  $\Lambda$

(direction de 2 thèses)

étude des interactions quasi-élastiques dans NOMAD (direction d'une thèse)

**1999-2008** : expérience HARP au CERN

analyse des données pour K2K et MiniBooNE et réduction d'un facteur 2 des incertitudes de flux

Membre du steering committee de l'expérience et de son Editorial Board

**2006-2020** : expérience NA61/SHINE au CERN

coordinateur du software et de l'analyse de l'expérience

premières mesures sur cible mince pour T2K (direction d'une thèse)

premières mesures sur la cible réplique de T2K (direction d'une thèse)

**2007-2020** : expérience T2K au Japon

conception, construction et installation du détecteur

coordinateur du groupe de travail sur le faisceau de T2K



referee interne pour la publication de découverte des oscillations  $\nu_\mu \rightarrow \nu_e$  dans T2K  
récipiendaire du prix La Recherche 2012 (Physique) au nom de T2K pour cette découverte  
*2016 Breakthrough Prize in Fundamental Physics* avec Daya Bay, Kamland, K2K & T2K, SNO,  
SK collègues

**2018- present:** expérience Hyper-Kamiokande au Japon  
conception du détecteur et développement d'un système de distribution et de synchronisation  
de temps  
études de sensibilité physique

## Liste de publications

1) [Constraint on the Matter-Antimatter Symmetry-Violating Phase in Neutrino Oscillations.](#)

By T2K Collaboration (K. Abe et al.), *to be published in Nature*,  
[arXiv:1910.03887 [hep-ex]].

2) [Hyper-Kamiokande Design Report.](#)

By Hyper-Kamiokande Collaboration (K. Abe et al.).  
[arXiv:1805.04163 [physics.ins-det]].

3) [Physics potentials with the second Hyper-Kamiokande detector in Korea.](#)

By Hyper-Kamiokande Collaboration (K. Abe et al.).  
[arXiv:1611.06118 [hep-ex]], PTEP 2018 (2018) no.6, 063C01 [10.1093/ptep/pty044](https://doi.org/10.1093/ptep/pty044).

4) [Measurement of neutrino and antineutrino neutral-current quasielasticlike interactions on oxygen by detecting nuclear deexcitation  \$\gamma\$  rays.](#)

By T2K Collaboration (K. Abe et al.).  
[arXiv:1910.09439 [hep-ex]], Phys.Rev. D100 (2019) no.11, 112009 [10.1103/PhysRevD.100.112009](https://doi.org/10.1103/PhysRevD.100.112009).

5) [Measurements of hadron production in  \$\pi^+ + C\$  and  \$\pi^+ + Be\$  interactions at 60 GeV/c\\$.](#)

By NA61/SHINE Collaboration (A. Aduszkiewicz et al.).  
[arXiv:1909.06294 [hep-ex]], Phys.Rev. D100 (2019) no.11, 112004 [10.1103/PhysRevD.100.112004](https://doi.org/10.1103/PhysRevD.100.112004).

6) [Measurement of the muon neutrino charged-current single  \$\pi^+\$  production on hydrocarbon using the T2K off-axis near detector ND280.](#)

By T2K Collaboration (K. Abe et al.).  
[arXiv:1909.03936 [hep-ex]], Phys.Rev. D101 (2020) no.1, 012007 [10.1103/PhysRevD.101.012007](https://doi.org/10.1103/PhysRevD.101.012007).

7) [Measurements of production and inelastic cross sections for  \$p + C\$ ,  \$p + Be\$ , and  \$p + Al\$  at 60 GeV/c\\$ and  \$p + C\$  and  \$p + Be\$  at 120 GeV/c\\$.](#)

By NA61/SHINE Collaboration (A. Aduszkiewicz et al.).  
[arXiv:1909.03351 [hep-ex]], Phys.Rev. D100 (2019) no.11, 112001 [10.1103/PhysRevD.100.112001](https://doi.org/10.1103/PhysRevD.100.112001).

8) [J-PARC Neutrino Beamline Upgrade Technical Design Report.](#)

By T2K Collaboration and J-PARC Neutrino Facility Group (K. Abe et al.).  
[arXiv:1908.05141 [physics.ins-det]].

9) [Measurement of  \$\phi\$  Meson Production in p + p Interactions at 40, 80 and 158 GeV/c with the NA61/SHINE Spectrometer at the CERN SPS.](#)

By NA61/SHINE Collaboration (A. Aduszkiewicz et al.).  
[arXiv:1908.04601 [nucl-ex]].

10) [Performances of a resistive MicroMegas module for the Time Projection Chambers of the T2K Near Detector upgrade.](#)

By D. Attié et al..  
[arXiv:1907.07060 [physics.ins-det]], Nucl.Instrum.Meth. A957 (2020) 163286 [10.1016/j.nima.2019.163286](https://doi.org/10.1016/j.nima.2019.163286).

11) Measurement of the  $\nu_\mu$  charged-current cross sections on water, hydrocarbon, iron, and their ratios with the T2K on-axis detectors.

By T2K Collaboration (K. Abe et al.).

[arXiv:1904.09611 [hep-ex]], Prog Theor Exp Phys (2019) [10.1093/ptep/ptz070](https://doi.org/10.1093/ptep/ptz070).

12) Search for heavy neutrinos with the T2K near detector ND280.

By T2K Collaboration (K. Abe et al.).

[arXiv:1902.07598 [hep-ex]], Phys.Rev. D100 (2019) no.5, 052006 [10.1103/PhysRevD.100.052006](https://doi.org/10.1103/PhysRevD.100.052006).

13) Search for light sterile neutrinos with the T2K far detector Super-Kamiokande at a baseline of 295 km.

By T2K Collaboration (K. Abe et al.).

[arXiv:1902.06529 [hep-ex]], Phys.Rev. D99 (2019) no.7, 071103 [10.1103/PhysRevD.99.071103](https://doi.org/10.1103/PhysRevD.99.071103).

14) Search for neutral-current induced single photon production at the ND280 near detector in T2K.

By T2K Collaboration (K. Abe et al.).

[arXiv:1902.03848 [hep-ex]], J.Phys. G46 (2019) no.8, 08LT01 [10.1088/1361-6471/ab227d](https://doi.org/10.1088/1361-6471/ab227d).

15) Research and Development for Near Detector Systems Towards Long Term Evolution of Ultra-precise Long-baseline Neutrino Experiments.

By Luis Alvarez Ruso et al..

[arXiv:1901.04346 [physics.ins-det]].

16) Future Opportunities in Accelerator-based Neutrino Physics.

By Andrea Dell'Acqua et al..

[arXiv:1812.06739 [hep-ex]].

17) Measurements of  $\pi^\pm$ ,  $K^\pm$  and proton double differential yields from the surface of the T2K replica target for incoming 31 GeV/c protons with the NA61/SHINE spectrometer at the CERN SPS.

By NA61/SHINE Collaboration (N. Abgrall et al.).

[arXiv:1808.04927 [hep-ex]], Eur.Phys.J. C79 (2019) no.2, 100 [10.1140/epjc/s10052-019-6583-0](https://doi.org/10.1140/epjc/s10052-019-6583-0).

18) Search for CP Violation in Neutrino and Antineutrino Oscillations by the T2K Experiment with  $2.2 \times 10^{21}$  Protons on Target.

By T2K Collaboration (K. Abe et al.).

[arXiv:1807.07891 [hep-ex]], Phys.Rev.Lett. 121 (2018) no.17, 171802 [10.1103/PhysRevLett.121.171802](https://doi.org/10.1103/PhysRevLett.121.171802).

19) Measurements of total production cross sections for  $\pi^+ + C$ ,  $\pi^+ + Al$ ,  $K^+ + C$ , and  $K^+ + Al$  at 60 GeV/c and  $\pi^+ + C$  and  $\pi^+ + Al$  at 31 GeV/c.

By NA61/SHINE Collaboration (A. Aduszkiewicz et al.).

[arXiv:1805.04546 [hep-ex]], Phys.Rev. D98 (2018) no.5, 052001 [10.1103/PhysRevD.98.052001](https://doi.org/10.1103/PhysRevD.98.052001).

20) Characterization of nuclear effects in muon-neutrino scattering on hydrocarbon with a measurement of final-state kinematics and correlations in charged-current pionless interactions at T2K.

By T2K Collaboration (K. Abe et al.).

[arXiv:1802.05078 [hep-ex]], Phys.Rev. D98 (2018) no.3, 032003 [10.1103/PhysRevD.98.032003](https://doi.org/10.1103/PhysRevD.98.032003).

21) Measurement of inclusive double-differential  $\nu_\mu$  charged-current cross section with improved acceptance in the T2K off-axis near detector.

By T2K Collaboration (K. Abe et al.).

[arXiv:1801.05148 [hep-ex]], Phys.Rev. D98 (2018) 012004 [10.1103/PhysRevD.98.012004](https://doi.org/10.1103/PhysRevD.98.012004).

- 22) [First measurement of the  \$\nu\_\mu / \mu\$  charged-current cross section on a water target without pions in the final state.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1708.06771 [hep-ex]], Phys.Rev. D97 (2018) no.1, 012001 [10.1103/PhysRevD.97.012001](https://doi.org/10.1103/PhysRevD.97.012001).

- 23) [Measurement of neutrino and antineutrino oscillations by the T2K experiment including a new additional sample of  \$\nu\_e\$  interactions at the far detector.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1707.01048 [hep-ex]], Phys.Rev. D96 (2017) no.9, 092006, Erratum: Phys.Rev. D98 (2018) no.1, 019902.

[10.1103/PhysRevD.98.019902](https://doi.org/10.1103/PhysRevD.98.019902), [10.1103/PhysRevD.96.092006](https://doi.org/10.1103/PhysRevD.96.092006).

- 24) [Measurement of  \$\bar{\nu}\_\mu / \mu\$  and  \$\nu\_\mu / \mu\$  charged current inclusive cross sections and their ratio with the T2K off-axis near detector.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1706.04257 [hep-ex]], Phys.Rev. D96 (2017) no.5, 052001 [10.1103/PhysRevD.96.052001](https://doi.org/10.1103/PhysRevD.96.052001).

- 25) [Measurement of meson resonance production in  \$\pi^+ + C\$  interactions at SPS energies.](#)

By NA61/SHINE Collaboration (A. Aduszkiewicz et al.).

[arXiv:1705.08206 [nucl-ex]], Eur.Phys.J. C77 (2017) no.9, 626 [10.1140/epjc/s10052-017-5184-z](https://doi.org/10.1140/epjc/s10052-017-5184-z).

- 26) [Measurements of  \$\pi^\pm\$ ,  \$K^\pm\$ ,  \$p\$  and  \$\bar{p}\$  spectra in proton-proton interactions at 20, 31, 40, 80 and 158 GeV/c with the NA61/SHINE spectrometer at the CERN SPS.](#)

By NA61/SHINE Collaboration (A. Aduszkiewicz et al.).

[arXiv:1705.02467 [nucl-ex]], Eur.Phys.J. C77 (2017) no.10, 671 [10.1140/epjc/s10052-017-5260-4](https://doi.org/10.1140/epjc/s10052-017-5260-4).

- 27) [Measurement of the single  \$\pi^0\$  production rate in neutral current neutrino interactions on water.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1704.07467 [hep-ex]], Phys.Rev. D97 (2018) no.3, 032002 [10.1103/PhysRevD.97.032002](https://doi.org/10.1103/PhysRevD.97.032002).

- 28) [Updated T2K measurements of muon neutrino and antineutrino disappearance using  \$1.5 \times 10^{21}\$  protons on target.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1704.06409 [hep-ex]], Phys.Rev. D96 (2017) no.1, 011102 [10.1103/PhysRevD.96.011102](https://doi.org/10.1103/PhysRevD.96.011102).

- 29) [Search for Lorentz and CPT violation using sidereal time dependence of neutrino flavor transitions over a short baseline.](#)

By T2K Collaboration (Ko Abe et al.).

[arXiv:1703.01361 [hep-ex]], Phys.Rev. D95 (2017) no.11, 111101 [10.1103/PhysRevD.95.111101](https://doi.org/10.1103/PhysRevD.95.111101).

- 30) [Combined Analysis of Neutrino and Antineutrino Oscillations at T2K.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1701.00432 [hep-ex]], Phys.Rev.Lett. 118 (2017) no.15, 151801 [10.1103/PhysRevLett.118.151801](https://doi.org/10.1103/PhysRevLett.118.151801).

- 31) [Two-particle correlations in azimuthal angle and pseudorapidity in inelastic p + p interactions at the CERN Super Proton Synchrotron.](#)

By NA61/SHINE Collaboration (A. Aduszkiewicz et al.).

[arXiv:1610.00482 [nucl-ex]], Eur.Phys.J. C77 (2017) no.2, 59 [10.1140/epjc/s10052-017-4599-x](https://doi.org/10.1140/epjc/s10052-017-4599-x).

32) [Proposal for an Extended Run of T2K to  \$20 \times 10^{21}\$  POT.](#)

By Ko Abe et al..

[arXiv:1609.04111 [hep-ex]].

33) [Sensitivity of the T2K accelerator-based neutrino experiment with an Extended run to  \$20 \times 10^{21}\$  POT.](#)

By T2K Collaboration (Ko Abe et al.).

[arXiv:1607.08004 [hep-ex]].

34) [First measurement of the muon neutrino charged current single pion production cross section on water with the T2K near detector.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1605.07964 [hep-ex]], Phys.Rev. D95 (2017) no.1, 012010 [10.1103/PhysRevD.95.012010](https://doi.org/10.1103/PhysRevD.95.012010).

35) [Measurement of Coherent  \$\pi^+\$  Production in Low Energy Neutrino-Carbon Scattering.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1604.04406 [hep-ex]], Phys.Rev.Lett. 117 (2016) no.19, 192501 [10.1103/PhysRevLett.117.192501](https://doi.org/10.1103/PhysRevLett.117.192501).

36) [Measurements of  \$\pi^\pm\$  differential yields from the surface of the T2K replica target for incoming 31 GeV/c protons with the NA61/SHINE spectrometer at the CERN SPS.](#)

By NA61/SHINE Collaboration (N. Abgrall et al.).

[arXiv:1603.06774 [hep-ex]], Eur.Phys.J. C76 (2016) no.11, 617 [10.1140/epjc/s10052-016-4440-y](https://doi.org/10.1140/epjc/s10052-016-4440-y).

37) [Measurement of double-differential muon neutrino charged-current interactions on C\\$ 8\\$H\\$ 8\\$ without pions in the final state using the T2K off-axis beam.](#)

By T2K Collaboration (Ko Abe et al.).

[arXiv:1602.03652 [hep-ex]], Phys.Rev. D93 (2016) no.11, 112012 [10.1103/PhysRevD.93.112012](https://doi.org/10.1103/PhysRevD.93.112012).

38) [Measurement of Muon Antineutrino Oscillations with an Accelerator-Produced Off-Axis Beam.](#)

By T2K Collaboration (Ko Abe et al.).

[arXiv:1512.02495 [hep-ex]], Phys.Rev.Lett. 116 (2016) no.18, 181801 [10.1103/PhysRevLett.116.181801](https://doi.org/10.1103/PhysRevLett.116.181801).

39) [Production of  \$\Lambda\$ -hyperons in inelastic p+p interactions at 158  \$\mathrm{GeV}/c\$ .](#)

By NA61/SHINE Collaboration (A. Aduszkiewicz et al.).

[arXiv:1510.03720 [hep-ex]], Eur.Phys.J. C76 (2016) no.4, 198 [10.1140/epjc/s10052-016-4003-2](https://doi.org/10.1140/epjc/s10052-016-4003-2).

40) [Measurements of  \$\pi^\pm\$ ,  \$K^\pm\$ ,  \$K^0\$  and  \$\bar{\Lambda}\$  and proton production in proton-carbon interactions at 31 GeV/c with the NA61/SHINE spectrometer at the CERN SPS.](#)

By NA61/SHINE Collaboration (N. Abgrall et al.).

[arXiv:1510.02703 [hep-ex]], Eur.Phys.J. C76 (2016) no.2, 84 [10.1140/epjc/s10052-016-3898-y](https://doi.org/10.1140/epjc/s10052-016-3898-y).

41) [Multiplicity and transverse momentum fluctuations in inelastic proton-proton interactions at the CERN Super Proton Synchrotron.](#)

By NA61/SHINE Collaboration (A. Aduszkiewicz et al.).

[arXiv:1510.00163 [hep-ex]], Eur.Phys.J. C76 (2016) no.11, 635 [10.1140/epjc/s10052-016-4450-9](https://doi.org/10.1140/epjc/s10052-016-4450-9).

42) [Measurement of the muon neutrino inclusive charged-current cross section in the energy range of 1–3 GeV with the T2K INGRID detector.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1509.06940 [hep-ex]], Phys.Rev. D93 (2016) no.7, 072002 [10.1103/PhysRevD.93.072002](https://doi.org/10.1103/PhysRevD.93.072002).

43) [Fixed-target hadron production experiments.](#)

By Boris A. Popov, EPJ Web Conf. 99 (2015) 02002 [10.1051/epjconf/20159902002](https://doi.org/10.1051/epjconf/20159902002).

44) [Measurement of the electron neutrino charged-current interaction rate on water with the T2K ND280  \$\pi^0\$  detector.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1503.08815 [hep-ex]], Phys.Rev. D91 (2015) 112010 [10.1103/PhysRevD.91.112010](https://doi.org/10.1103/PhysRevD.91.112010).

45) [Measurement of the  \$\nu\_\mu\$  charged current quasielastic cross section on carbon with the T2K on-axis neutrino beam.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1503.07452 [hep-ex]], Phys.Rev. D91 (2015) no.11, 112002 [10.1103/PhysRevD.91.112002](https://doi.org/10.1103/PhysRevD.91.112002).

46) [Upper bound on neutrino mass based on T2K neutrino timing measurements.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1502.06605 [hep-ex]], Phys.Rev. D93 (2016) no.1, 012006 [10.1103/PhysRevD.93.012006](https://doi.org/10.1103/PhysRevD.93.012006).

47) [Measurements of neutrino oscillation in appearance and disappearance channels by the T2K experiment with  \$6.6 \times 10^{20}\$  protons on target.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1502.01550 [hep-ex]], Phys.Rev. D91 (2015) no.7, 072010 [10.1103/PhysRevD.91.072010](https://doi.org/10.1103/PhysRevD.91.072010).

48) [Measurement of the  \$\nu\_e\$  charged-current quasielastic cross section on carbon with the ND280 detector at T2K.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1411.6264 [hep-ex]], Phys.Rev. D92 (2015) no.11, 112003 [10.1103/PhysRevD.92.112003](https://doi.org/10.1103/PhysRevD.92.112003).

49) [Search for short baseline  \$\nu\_e\$  disappearance with the T2K near detector.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1410.8811 [hep-ex]], Phys.Rev. D91 (2015) 05110 [10.1103/PhysRevD.91.051102](https://doi.org/10.1103/PhysRevD.91.051102).

50) [Neutrino oscillation physics potential of the T2K experiment.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1409.7469 [hep-ex]], PTEP 2015 (2015) no.4, 043C01 [10.1093/ptep/ptv031](https://doi.org/10.1093/ptep/ptv031).

51) [Measurement of the Inclusive Electron Neutrino Charged Current Cross Section on Carbon with the T2K Near Detector.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1407.7389 [hep-ex]], Phys.Rev.Lett. 113 (2014) no.24, 241803 [10.1103/PhysRevLett.113.241803](https://doi.org/10.1103/PhysRevLett.113.241803).

52) [Measurement of the inclusive  \$\nu\_\mu\$  charged current cross section on iron and hydrocarbon in the T2K on-axis neutrino beam.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1407.4256 [hep-ex]], Phys.Rev. D90 (2014) no.5, 052010 [10.1103/PhysRevD.90.052010](https://doi.org/10.1103/PhysRevD.90.052010).

53) [Measurement of the neutrino-oxygen neutral-current interaction cross section by observing nuclear deexcitation  \$\gamma\$  rays.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1403.3140 [hep-ex]], Phys.Rev. D90 (2014) no.7, 072012 [10.1103/PhysRevD.90.072012](https://doi.org/10.1103/PhysRevD.90.072012).

54) [Measurement of the intrinsic electron neutrino component in the T2K neutrino beam with the ND280 detector.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1403.2552 [hep-ex]], Phys.Rev. D89 (2014) 092003 [10.1103/PhysRevD.89.092003](https://doi.org/10.1103/PhysRevD.89.092003), Phys.Rev. D89 (2014) 099902 [10.1103/PhysRevD.89.099902](https://doi.org/10.1103/PhysRevD.89.099902).

55) [Precise Measurement of the Neutrino Mixing Parameter  \$\theta\_{23}\$  from Muon Neutrino Disappearance in an Off-Axis Beam.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1403.1532 [hep-ex]], Phys.Rev.Lett. 112 (2014) no.18, 181801 [10.1103/PhysRevLett.112.181801](https://doi.org/10.1103/PhysRevLett.112.181801).

56) [NA61/SHINE facility at the CERN SPS: beams and detector system.](#)

By NA61 Collaboration (N. Abgrall et al.).

[arXiv:1401.4699 [physics.ins-det]], JINST 9 (2014) P06005 [10.1088/1748-0221/9/06/P06005](https://doi.org/10.1088/1748-0221/9/06/P06005).

57) [Recent Results from the T2K Experiment.](#)

By K. Abe et al., Nucl.Phys.Proc.Suppl. 246-247 (2014) 23-28 [10.1016/j.nuclphysbps.2013.10.060](https://doi.org/10.1016/j.nuclphysbps.2013.10.060).

58) [The mass-hierarchy and CP-violation discovery reach of the LBNO long-baseline neutrino experiment.](#)

By LAGUNA-LBNO Collaboration (S.K. Agarwalla et al.).

[arXiv:1312.6520 [hep-ph]], JHEP 1405 (2014) 094 [10.1007/JHEP05\(2014\)094](https://doi.org/10.1007/JHEP05(2014)094).

59) [Observation of Electron Neutrino Appearance in a Muon Neutrino Beam.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1311.4750 [hep-ex]], Phys.Rev.Lett. 112 (2014) 061802 [10.1103/PhysRevLett.112.061802](https://doi.org/10.1103/PhysRevLett.112.061802).

60) [Measurement of negatively charged pion spectra in inelastic p+p interactions at  \$S\_{\text{lab}} = 20, 31, 40, 80\$  and 158 GeV/c.](#)

By NA61/SHINE Collaboration (N. Abgrall et al.).

[arXiv:1310.2417 [hep-ex]], Eur.Phys.J. C74 (2014) no.3, 2794 [10.1140/epjc/s10052-014-2794-6](https://doi.org/10.1140/epjc/s10052-014-2794-6).

61) [Measurements of production properties of  \$K^- \pi^0\$  mesons and  \$\Lambda\$  hyperons in proton-carbon interactions at 31 GeV/c.](#)

By NA61/SHINE Collaboration (N. Abgrall et al.).

[arXiv:1309.1997 [physics.acc-ph]], Phys.Rev. C89 (2014) no.2, 025205 [10.1103/PhysRevC.89.025205](https://doi.org/10.1103/PhysRevC.89.025205).

62) [A Precision Measurement of Charm Dimuon Production in Neutrino Interactions from the NOMAD Experiment.](#)

By NOMAD Collaboration (O. Samoylov et al.).

[arXiv:1308.4750 [hep-ex]], Nucl.Phys. B876 (2013) 339-375 [10.1016/j.nuclphysb.2013.08.021](https://doi.org/10.1016/j.nuclphysb.2013.08.021).

63) [Measurement of Neutrino Oscillation Parameters from Muon Neutrino Disappearance with an Off-axis Beam.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1308.0465 [hep-ex]], Phys.Rev.Lett. 111 (2013) no.21, 211803 [10.1103/PhysRevLett.111.211803](https://doi.org/10.1103/PhysRevLett.111.211803).

64) [Evidence of Electron Neutrino Appearance in a Muon Neutrino Beam.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1304.0841 [hep-ex]], Phys.Rev. D88 (2013) no.3, 032002. [10.1103/PhysRevD.88.032002](https://doi.org/10.1103/PhysRevD.88.032002).

- 65) [Measurement of the inclusive  \$\bar{\nu}\_e\$  charged current cross section on carbon in the near detector of the T2K experiment.](#)  
By T2K Collaboration (K. Abe et al.).  
[arXiv:1302.4908 [hep-ex]], Phys.Rev. D87 (2013) no.9, 092003 [10.1103/PhysRevD.87.092003](https://doi.org/10.1103/PhysRevD.87.092003).
- 66) [Energy dependence of identified hadron spectra and event-by-event fluctuations in p+p interactions from NA61/SHINE at the CERN SPS.](#)  
By NA61/SHINE Collaboration (Maciej Rybczynski et al.).  
[arXiv:1301.3360 [nucl-ex]], PoS ConfinementX (2012) 207 [10.22323/1.171.0207](https://doi.org/10.22323/1.171.0207).
- 67) [Hadron production experiments.](#)  
By Boris A. Popov.  
[arXiv:1212.1030 [hep-ex]], Nucl.Phys.Proc.Suppl. 235-236 (2013) 135-142 [10.1016/j.nuclphysbps.2013.04.003](https://doi.org/10.1016/j.nuclphysbps.2013.04.003).
- 68) [T2K neutrino flux prediction.](#)  
By T2K Collaboration (K. Abe et al.).  
[arXiv:1211.0469 [hep-ex]], Phys.Rev. D87 (2013) no.1, 012001, Addendum: Phys.Rev. D87 (2013) no.1, 019902 [10.1103/PhysRevD.87.019902](https://doi.org/10.1103/PhysRevD.87.019902), [10.1103/PhysRevD.87.012001](https://doi.org/10.1103/PhysRevD.87.012001).
- 69) [Expression of Interest for a very long baseline neutrino oscillation experiment \(LBNO\).](#)  
By A. Stahl et al..
- 70) [Pion emission from the T2K replica target: method, results and application.](#)  
By NA61/SHINE Collaboration (N. Abgrall et al.).  
[arXiv:1207.2114 [hep-ex]], Nucl.Instrum.Meth. A701 (2013) 99-114 [10.1016/j.nima.2012.10.079](https://doi.org/10.1016/j.nima.2012.10.079).
- 71) [First Muon-Neutrino Disappearance Study with an Off-Axis Beam.](#)  
By T2K Collaboration (K. Abe et al.).  
[arXiv:1201.1386 [hep-ex]], Phys.Rev. D85 (2012) 031103. [10.1103/PhysRevD.85.031103](https://doi.org/10.1103/PhysRevD.85.031103).
- 72) [Measurement of Production Properties of Positively Charged Kaons in Proton-Carbon Interactions at 31 GeV/c.](#)  
By NA61/SHINE Collaboration (N. Abgrall et al.).  
[arXiv:1112.0150 [hep-ex]], Phys.Rev. C85 (2012) 035210 [10.1103/PhysRevC.85.035210](https://doi.org/10.1103/PhysRevC.85.035210).
- 73) [A search for single photon events in neutrino interactions.](#)  
By NOMAD Collaboration (C.T. Kullenberg et al.).  
[arXiv:1111.3713 [hep-ex]], Phys.Lett. B706 (2012) 268-275 [10.1016/j.physletb.2011.11.049](https://doi.org/10.1016/j.physletb.2011.11.049).
- 74) [Measurements of the T2K neutrino beam properties using the INGRID on-axis near detector.](#)  
By K. Abe et al..  
[arXiv:1111.3119 [physics.ins-det]], Nucl.Instrum.Meth. A694 (2012) 211-223 [10.1016/j.nima.2012.03.023](https://doi.org/10.1016/j.nima.2012.03.023).
- 75) [Results from hadroproduction experiments.](#)  
By Boris A. Popov.  
[arXiv:1106.4380 [hep-ex]].
- 76) [Indication of Electron Neutrino Appearance from an Accelerator-produced Off-axis Muon Neutrino Beam.](#)  
By T2K Collaboration (K. Abe et al.).  
[arXiv:1106.2822 [hep-ex]], Phys.Rev.Lett. 107 (2011) 041801 [10.1103/PhysRevLett.107.041801](https://doi.org/10.1103/PhysRevLett.107.041801).



77) [The T2K Experiment.](#)

By T2K Collaboration (K. Abe et al.).

[arXiv:1106.1238 [physics.ins-det]], Nucl.Instrum.Meth. A659 (2011) 106-135 [10.1016/j.nima.2011.06.067](https://doi.org/10.1016/j.nima.2011.06.067).

78) [Measurements of Cross Sections and Charged Pion Spectra in Proton-Carbon Interactions at 31 GeV/c.](#)

By NA61/SHINE Collaboration (N Abgrall et al.).

[arXiv:1102.0983 [hep-ex]], Phys.Rev. C84 (2011) 034604 [10.1103/PhysRevC.84.034604](https://doi.org/10.1103/PhysRevC.84.034604).

79) [Time Projection Chambers for the T2K Near Detectors.](#)

By T2K ND280 TPC Collaboration (N. Abgrall et al.).

[arXiv:1012.0865 [physics.ins-det]], Nucl.Instrum.Meth. A637 (2011) 25-46 [10.1016/j.nima.2011.02.036](https://doi.org/10.1016/j.nima.2011.02.036).

80) [Measurements of Forward Proton Production with Incident Protons and Charged Pions on Nuclear Targets at the CERN Proton Synchroton.](#)

By HARP Collaboration (M. Apollonio et al.).

[arXiv:1006.1223 [hep-ex]], Phys.Rev. C82 (2010) 045208 [10.1103/PhysRevC.82.045208](https://doi.org/10.1103/PhysRevC.82.045208).

81) [HARP and NA61 \(SHINE\) hadron production experiments.](#)

By HARP and NA61 Collaborations (Boris A. Popov for the collaboration).

AIP Conf.Proc. 1189 (2009) no.1, 97-104 [10.1063/1.3274196](https://doi.org/10.1063/1.3274196).

#### 4. Avis motivé de la direction du laboratoire et signature :

Le projet expérimental Hyper-Kamiokande sera un formidable outil d'exploration et de découverte des neutrinos du cosmos grâce à sa grande masse fiduciale et à son excellente sensibilité avec une gamme en énergie allant de quelques MeV à plusieurs dizaines de GeV. Il sera ainsi extrêmement performant pour la détection des neutrinos des supernovas par effondrement gravitationnel, qu'on pourra suivre dans le moindre détail de l'explosion grâce à la détection de plusieurs dizaines de milliers d'événements (pour une SN dans notre galaxie). La détection du fond diffus de supernovas, des neutrinos solaires et des neutrinos atmosphériques, et la recherche de désintégration du proton permettront d'élargir le domaine scientifique couvert par cet instrument astrophysique exceptionnel.

Le groupe Hyper-Kamiokande du laboratoire a une solide expérience dans la détection et l'étude des neutrinos notamment grâce à son activité dans le projet T2K au Japon qui a permis des avancées importantes dans le domaine. Le groupe et les services techniques du LPNHE travaillent à la préparation de l'expérience Hyper-Kamiokande, notamment avec le développement de la synchronisation de l'électronique des photomultiplicateurs, ce qui permettra d'optimiser la sensibilité du détecteur.

La thèse proposée combine une activité expérimentale au plus près du dispositif de détection avec des développements d'analyse des données sur la base des outils informatiques les plus sophistiqués. Cela permettra une formation scientifique solide et complète à l'étudiant.

Je soutiens fortement ce projet.

Marco Zito  
LPNHE UMR 7585

## 5. Suggestion d'experts extérieurs :

Afin d'accélérer le traitement des dossiers, les proposants peuvent suggérer deux noms (avec leurs coordonnées complètes et précises) d'experts localisés en dehors de l'Île-de-France (en France ou à l'étranger) et susceptibles de réaliser l'expertise du projet.

La coordination du DIM se réserve le droit de choisir les experts parmi les propositions qui leur sont faites ou d'autres sources.

### Coordonnées de l'expert n°1

Nom - Prénom : Sanchez Federico

Fonction : Professor

Organisme : Geneva University

Adresse : 24, quai Ernest-Ansermet, CH-1211 Genève 4, Switzerland

Téléphone : +41 223796227

E-mail : Federico Sanchez Nieto <Federico.SanchezNieto@unige.ch>

Spécialité : Neutrino physics

### Coordonnées de l'expert n°2

Nom - Prénom : Mezzetto Mauro

Fonction : Dr

Organisme : INFN-Padova

Adresse : Department of Physics and Astronomy of Padova University, Via Francesco Marzolo, 8, 35131 Padova PD, Italy

Téléphone : +39 0499677147

E-mail : mauro.mezzetto@pd.infn.it

Spécialité : Neutrino physics and astrophysics