

# Hidden Higgs


Adam Falkowski

Rutgers University

Rencontre de Physique des Particules, 25 January 2010

## What do we know about Higgs?


- The Higgs boson is predicted by many theories of electroweak symmetry breaking
- Higgs is the *simplest* mean to unitarize the scattering amplitude of longitudinally polarized W and Z bosons
- A *light* Higgs boson is strongly suggested by electroweak precision observables



- *Hints* that there exists a light scalar field with the coupling  $\sim m_V V_\mu V_\mu h$  to the W and Z gauge bosons
- *But* there is no experimental input as to Higgs coupling to SM fermions!

# Tension

- **Experimental**: tension between the LEP limit  $m_h > 114.4 \text{ GeV}$  and the electroweak fit  $m_h = 80^{+30}_{-23}$ 
  - ▶ Leptonic observables and W mass alone prefer a very light Higgs, of order 60 GeV!
  - ▶ Only the  $Z \rightarrow b\bar{b}$  forward-backward asymmetry pushes the Higgs mass toward larger values



- **Theoretical**: In many extensions of the SM, in particular in the MSSM or simplest little Higgs theories,  $m_{\text{Higgs}} \approx m_Z$  preferred by naturalness, while  $m_{\text{Higgs}} \geq 115 \text{ GeV}$  leads to the *little hierarchy problem*


There is some tension within the minimal Higgs paradigm, which prompts searching for alternatives

## So maybe Higgs IS lighter than 115 GeV?

- **One possibility**: suppressed coupling to Z boson, so that it was not produced at LEP. But then electroweak fit is not improved even if Higgs is light
- **More exciting possibility**: Higgs was copiously produced at LEP, but it escaped our attention due to non-standard decays. This is what I call **Hidden Higgs**


## So maybe Higgs IS lighter than 115 GeV?

- **One possibility:** suppressed coupling to Z boson, so that it was not produced at LEP. But then electroweak fit is not improved even if Higgs is light
- **More exciting possibility:** Higgs was copiously produced at LEP, but it escaped our attention due to non-standard decays. This is what I call **Hidden Higgs**




## Recall the Standard Model Higgs

- LEP constrained  $m_{\text{Higgs}} > 114.4$  GeV by looking for  $H \rightarrow b\bar{b}$  decay



### Higgs decay branching ratios



- SM Higgs couples to mass
- For a light Higgs, the couplings to the relevant SM states are tiny, e.g.  
 $y_b \sim m_b/v_{EW} \sim 0.02$
- Branching ratios for various Higgs decays can easily be altered by new physics

# Higgs limits beyond the SM

Assuming SM production cross section, and  $BR(H \rightarrow xx) = 1$

| <i>Decay Channel</i>                     | <b>Limit</b> |
|--|--------------|
| $h \rightarrow \tilde{\chi}\tilde{\chi}$ | 114 GeV      |
| $h \rightarrow \tau\tau$                 | 115 GeV      |
| $h \rightarrow jj$                       | 113 GeV      |
| $h \rightarrow WW^* \text{ or } ZZ^*$    | 110 GeV      |
| $h \rightarrow AA \rightarrow 4b, 4\tau$ | 110 GeV      |
| $h \rightarrow AA \rightarrow 4c, 4g$    | 86 GeV       |
| $h \rightarrow \text{anything}$          | 82 GeV       |

see [Chang, Fox, Weiner \[hep-ph/0608310\]](#) for review


- Invisible and two-body decay channels very well constrained
- Constraints on four- and more body decay channels typically not much better than the model independent OPAL constraint, with the exception of the  $4b$  and  $4\tau$  channels
- Typically, the multiparticle channels are weakly constrained not because of fundamental reasons but because nobody bothered to look

*Are there any more or less motivated models with Higgs decaying to multiparticle states?*

# Hidden Higgs in NMSSM

Best known example of Hidden Higgs: NMSSM near R-symmetric or PQ-symmetric limit

- NMSSM:  $W = \lambda S H_u H_d + \kappa S^3$ ,  $V_{\text{soft}} = A_\lambda \lambda S H_u H_d + A_\kappa \kappa S^3 + m_S^2 |S|^2$
- Two CP-odd Higgses  $A_{1,2}$ : one in  $S$ , one in  $H_{u,d}$ , that mix due to EW breaking. The  $A_1$  mass for large  $\tan \beta$  is  $m_{A_1}^2 \sim \kappa A_\kappa \mu / \lambda$ . It is light if e.g.  $A_\kappa \sim \text{GeV}$
- $A_1$  has sizable coupling to Higgs via potential, so cascade decay  $h \rightarrow AA \rightarrow 4f$  may easily dominate








# Non-standard Higgs decays in NMSSM

- Much as Higgs, pseudoscalar  $A_1$  couples more strongly to heavier SM particles
- For  $m_{A_1} > 2m_b$  the dominant decay of  $A_1$  is into 2 b quarks

Dobrescu, Landsberg, Matchev [hep-ph/0008192] Dermisek, Gunion [hep-ph/0502105] . Constrained by LEP [hep-ex/0602042]




- For  $2m_\tau < m_{A_1} < 2m_b$  dominant decay into 2 tau leptons Dermisek, Gunion [hep-ph/0611142] . Constrained by Cranmer [20 years of ALEPH]
- For event lighter  $A_1$ , it decays to a pair of gluons; because the 2 gluons are very collimated this case is probably covered by  $H \rightarrow 2j$  analysis

# Other Hidden Higgs models

- The possibility of Hidden Higgs within the NMSSM now seems to be closed
- Nevertheless, a neat example of complicated but finally successful theory-experiment interactions
- Other realizations of Hidden Higgs are still alive
  - ⚡  $H \rightarrow 6j$  in R-parity violating MSSM [Carpenter,Kaplan,Rhee \[hep-ph/0607204\]](#)
  - ⚡  $H \rightarrow 4j$  (Buried Higgs) in SUSY Little Higgs [Bellazzini,Csaki,AA,Weiler \[0906.3026\]](#)
  - ⚡  $H \rightarrow$  lepton jets in MSSM+light hidden sector [AA,Ruderman,Volansky,Zupan \[to appear\]](#)

## Higgs to 6 jets

- Higgs can cascade decay into 6 quarks within the R-parity violating MSSM
- First, Higgs decays into the lightest MSSM neutralino. A large branching fraction requires
  - ▶  $m_{N_1} < m_{Higgs}/2 \sim 50 \text{ GeV}$  (not excluded by experiment if  $N_1$  is mostly bino)
  - ▶  $N_1$  has some (at least 20 percent) higgsino component
- The lightest neutralino can decay into 3 quarks via an off-shell squark,  $N_1 \rightarrow q\tilde{q} \rightarrow qq\bar{q}$ , if the R-parity violating operator  $U^c D^c D^c$  is present in the superpotential
- No bounds on the Higgs mass, except the model independent bound of 82 GeV



# Buried Higgs

- Higgs can cascade decay into four light colored objects within little SUSY (supersymmetric little Higgs models)
- The MSSM extended to include  $SU(3)$  global symmetry spontaneously broken to  $SU(2)$  at the scale  $f \gtrsim v_{EW}$
- Instead of Higgs doublets  $H_{u,d}$ , Higgs triplets  $\mathcal{H}_{u,d}$
- 5 Goldstone bosons from  $SU(3) \rightarrow SU(2)$  breaking, 3 of which get eaten by W and Z after EW breaking



Two physical pGB scalars  $h$  and  $\eta$  embedded in the triplets as

$$\mathcal{H}_u \approx f \sin \beta \begin{pmatrix} 0 \\ \sin((\tilde{v} + h)/f) \\ e^{i\eta/f} \cos((\tilde{v} + h)/f) \end{pmatrix} \quad \mathcal{H}_d^T \approx f \cos \beta \begin{pmatrix} 0 \\ \sin((\tilde{v} + h)/f) \\ e^{-i\eta/f} \cos((\tilde{v} + h)/f) \end{pmatrix}.$$

- The pGB scalar  $h$  identified with the SM Higgs boson
- The pGB pseudoscalar  $\eta$  is a new singlet

## Buried Higgs

- ⚡ Singlet pseudoscalar  $\eta$  is naturally light, thanks to global symmetry protection,
  - ⚡ lives in the 3rd component of the triplet and does not couple to W or Z.
  - ⚡ has derivative couplings to the Higgs,  $\sim f^{-1} h (\partial_\mu \eta)^2$
  - ⚡ couples to SM fermions via their mixing with heavy partner fermions
- Higgs decays dominantly to a pair of PGB pseudoscalars  $\eta$  as long as the scale  $f$  is not too large,  $f \lesssim 400 \text{ GeV}$
  - Couplings of  $\eta$  to SM fermions depend on fermion representations under global SU(3), masses of heavy fermionic partners of SM fermions, etc.
  - Several phenomenologically distinct realizations of Hidden Higgs
    - ▶ Gluophilic Higgs,  $h \rightarrow 4j$  when  $\eta \rightarrow gg$  dominates
    - ▶ Charming Higgs,  $h \rightarrow 4c$  when  $\eta \rightarrow cc$  dominates
  - In most cases, the standard discovery mode  $h \rightarrow \gamma\gamma$  is strongly suppressed




## Higgs to lepton jets

- ⚡ Higgs can decay into multiple collimated leptons (lepton jets) and missing energy, within supersymmetric models with light hidden sector
- Astrophysical observations, especially the PAMELA cosmic ray positron excess, hint at existence of a light, GeV scale hidden sector
- One possibility is that it contains a hidden massive photon  $z_\mu$  that mixes kinetically with the SM hypercharge,


$$\epsilon z_{\mu\nu} B_{\mu\nu} \quad \epsilon \leq 10^{-3}$$

- As a result, the hidden photon can decay into a pair of charged kinematically available SM states: electrons, muons, pions,...




## Higgs to lepton jets

- In a minimal supersymmetric framework hidden photon accompanied by 3 hidden neutralinos and 3 hidden Higgses Baumgart et al [0901.0283]
- The MSSM bino due to SUSY kinetically mixes with the hidden bino, which means that the MSSM LSP is no longer stable but decays into the hidden sector
- If Higgs decays dominantly into a pair of neutralinos or sneutrinos, then it decays into the hidden sector



- A hidden sector cascade follows, producing a number of hidden photons and the lightest hidden neutralino
- Higgs decay products emerge as lepton jets plus missing energy



- Not excluded LEP/Tevatron, though easy to spot by dedicated searches

# Summary

- LEP and Tevatron experiments may have missed a light Higgs if it has non-standard decays
- The gaps can be easily filled by dedicated analysis
- 😊 Ongoing ALEPH, L3 and Tevatron analyses
- 😊 LHC strategies to discover Higgs decaying into light jets or lepton jets in preparation
- A light Higgs could solve the SUSY little hierarchy problem, and improve electroweak fits
- Even if Higgs is heavier than 115 GeV, it is conceivable that non-standard Higgs decay show up at the LHC, as the leading or subleading channel. So better be prepared
- 😊 New interesting theoretical ideas are still being produced