

The background of the slide is a composite of two astronomical images. The left side shows a field of galaxy clusters, with many galaxies appearing as bright, elongated spots. The right side shows a colorful nebula with swirling clouds of gas in shades of red, purple, and blue. A diagonal line separates the two images, running from the top-left towards the bottom-right.

# **Intracluster dust and lensing**

**CALUM MURRAY, CELINE COMBET & CONSTANTIN PAYERNE.**

**-LABORATOIRE DE PHYSIQUE SUBATOMIQUE ET COSMOLOGIE, GRENOBLE**

---

# Intracluster dust


- Dust may be thrown into the intracluster medium by
  - Supernova explosions
  - Ram pressure stripping
  - Tidal interactions
  - Super massive blackholes!
- However once in the medium they will be heated by the intracluster X-ray gas
- Dust can cool the intracluster medium but also inform us of the phenomena that may send dust into the medium



NGC 4402 falling towards the Virgo supercluster - HST

# Cluster Lensing

- Shears galaxy images
- Increases galaxy magnitudes
- Galaxy surveys are magnitude limited, magnification **introduces faint galaxies** into the sample
- Deflects galaxy images away from the cluster centre, which **reduces the number of galaxies** in radial annuli from the centre
- Magnification has different systematics than shear!




# Dust and magnification

- Dust
  - Reduces background galaxy magnitude
- Lensing
  - Galaxies appear further from the cluster centre
  - Increase galaxy magnitudes
- However the dust effects are wavelength dependent!
- Galaxy magnitude is  $m$ ,  $\kappa$  is the lensing convergence,  $\tau_\lambda$  is the optical dust depth at a given wavelength,  $\vec{\theta}$  is the position of a background galaxy and  $\vec{\alpha}$  is the lensing angle

$$m_{obs} \approx m_{int} - \frac{5}{2 \ln 10} (2\kappa - \tau_\lambda)$$

$$\vec{\theta}_{obs} = \vec{\theta}_{int} + \vec{\alpha}_{lens}$$



# Dust and magnification

- Galaxy magnitude is  $m$ ,  $\kappa$  is the lensing convergence,  $\tau_\lambda$  is the optical dust depth at a given wavelength,  $\vec{\theta}$  is the position of a background galaxy and  $\vec{\alpha}$  is the lensing angle


$$m_{obs} \approx m_{int} - \frac{5}{2 \ln 10} (2\kappa - \tau_\lambda)$$

$$\vec{\theta}_{obs} = \vec{\theta}_{int} + \vec{\alpha}_{lens}$$

$$\delta m_{lens} \approx 0.1$$

$$\delta m_{dust} \approx 0.01$$

Towards the centre of a cluster we expect



---


# Objectives

- Joint measure of cluster dust content and cluster masses with **magnification**
- Introduce a **new** (and hopefully improved) likelihood which incorporates galaxy clustering, magnitude and redshift information

---

# Redmapper clusters and Hyper Suprime Cam

- 458 SDSS Redmapper clusters are found in the HSC field
- Red dots are SDSS Redmapper clusters
- White dots are Redmapper clusters within the HSC field
- HSC has a high galaxy density  $n_{gal} \approx 20 [\text{arcmin}^2]$
- 5 magnitude bands (grizy) important for dust searches
- HSC is a good test ground for Rubin (weak lensing)




# Average magnitude profiles

- Using a subsample of 90 clusters in the redshift interval  $0.2 < z_{cluster} < 0.3$
- We measure the average magnitude for a stack of clusters in annuli from the cluster centre
- Clear chromatic signal
- **Attention!** Lensing introduces colour changes, faint galaxies which are introduced to the sample have different colours to bright galaxies
- These profiles have been used to measure dust, not strictly true (Menard et al. 2009)

$$\langle \delta m \rangle = \langle m(\theta) \rangle - \langle m_{field} \rangle$$

$$m_{obs} \approx m_{int} - \frac{5}{2 \ln 10} (2\kappa - \tau_\lambda)$$





# Galaxy number counts

- We need to consider both the shift in galaxy position and the magnitude change
  - Lensing deflects galaxy images away from the cluster centre reducing the number of galaxies
  - Lensing introduces faint galaxies into the sample by increasing their brightness
  - Dust dims galaxies, reducing the number of galaxies
- $n$  is the number density of galaxies,  $\alpha$  is the **slope of the galaxy magnitude function** and as before  $\kappa$  is the lensing convergence,  $\tau_\lambda$  is the optical dust depth at a given wavelength
- $\alpha$  is calculated from the global galaxy distribution

$$\vec{\theta}_{obs} = \vec{\theta}_{int} + \vec{\alpha}_{lens}$$

$$m_{obs} \approx m_{int} - \frac{5}{2 \ln 10} (2\kappa - \tau_\lambda)$$

$$\alpha = 2.5 \frac{d \log_{10} n}{dm} \Big|_{m_{cut}}$$


New term due to dust that we have introduced

$$n_{obs}(\vec{\theta}) \approx n_{int}(\vec{\theta}) \left[ 1 + \alpha(e^{-\tau_\lambda} - 1) + 2\kappa(\alpha - 1) \right]$$

# The usual method

$$n_{obs}(\vec{\theta}) \approx n_{int}(\vec{\theta}) [1 + 2\kappa(\alpha - 1)]$$

- Choose **one magnitude cut** which gives a  $\alpha$  which maximises the signal
  - This is a game between the best value of  $\alpha$  and keeping as many galaxies as possible
  - Does not make much use of galaxy magnitude or redshift information
- Chiu, Umetsu et al. 2020, Schmidt et al. 2010, Broadhurst, Taylor and Peacock 1994
- Marina Ricci has been looking at this in CosmoDC2



---

# Incorporating galaxy magnitude and redshift information

- Bin observed galaxy distribution into **magnitude and redshift bins**
- $\alpha$  must now incorporate galaxies which **leave and enter** the magnitude bin
- The lensed/dusted galaxy distributions can be calculated from the unlensed/undusted distributions
- Dust depends on wavelength and angular separation
- Convergence depends on galaxy redshift and angular separation

$$\alpha = -2.5 \left( \frac{d \log_{10} n}{dm} \Big|_{m_{high}} - \frac{d \log_{10} n}{dm} \Big|_{m_{low}} \right)$$

$$n_{obs}(\vec{\theta}, m, z_{gal}) \approx n_{int}(\vec{\theta}, m, z_{gal}) \left[ 1 + \alpha(m, z_{gal})(e^{-\tau_{\lambda}(\vec{\theta})} - 1) + 2\kappa(\vec{\theta}, z_{gal})(\alpha(m, z_{gal}) - 1) \right]$$

---

# Poisson likelihood


- Analogous to that used in cluster abundance cosmology
- Simply measure the galaxy counts,  $n_{meas,k}$  in bins,  $k$ , of  $\theta$ , redshift, and magnitude and compare to the theoretical expectation  $n_{obs,k}$
- Uses **more information** than the standard method!

$$\ln \mathcal{L} = \sum_k n_{meas,k} \ln n_{obs,k} - \sum_k n_{obs,k}$$

$$n_{obs}(\vec{\theta}, m, z_{gal}) \approx n_{int}(\vec{\theta}, m, z_{gal}) \left[ 1 + \alpha(m, z_{gal})(e^{-\tau_\lambda(\vec{\theta})} - 1) + 2\kappa(\vec{\theta}, z_{gal})(\alpha(m, z_{gal}) - 1) \right]$$

# Mock result validation

- We generate random positions within the HSC footprint and inject a fake cluster signal
  - **Lensed galaxy positions**
- Lensed/dusted galaxy magnitudes




$$\vec{\theta}_{obs} = \vec{\theta}_{int} + \vec{\alpha}_{lens}$$

$$m_{obs} \approx m_{int} - \frac{5}{2 \ln 10} (2\kappa - \gamma_{\lambda})$$

# Mock result validation

- We generate random positions within the HSC footprint and inject a fake cluster signal
  - **Lensed galaxy positions**
  - Lensed/dusted galaxy magnitudes




$$\vec{\theta}_{obs} = \vec{\theta}_{int} + \vec{\alpha}_{lens}$$

$$m_{obs} \approx m_{int} - \frac{5}{2 \ln 10} (2\kappa - \gamma_{\lambda})$$

# Mock result validation

- We generate random positions within the HSC footprint and inject a fake cluster signal
  - **Lensed galaxy positions**
  - **Lensed/dusted galaxy magnitudes**




$$\vec{\theta}_{obs} = \vec{\theta}_{int} + \vec{\alpha}_{lens}$$


$$m_{obs} \approx m_{int} - \frac{5}{2 \ln 10} (2\kappa - \gamma_{\lambda})$$

# Mock results

- For these mocks we have only injected the cluster signal, **no dust**, we still see chromatic effects
- For reference we measure the signal around random points in the HSC footprint



Mocks





Redmapper clusters



# Mock results


- For these mocks we have only injected the cluster signal, **no dust**, we still see chromatic effects
- Galaxy counts in agreement with predictions
  - Here for two different magnitude cuts

$$n_{obs}(\vec{\theta}, m, z_{gal}) \approx n_{int}(\vec{\theta}, m, z_{gal}) [1 + 2\kappa(\alpha - 1)]$$



# Cluster mass estimates

- Work in progress
- 275 synthetic clusters
- 3 different cases, only angular bins is the standard method
  - 16 i-band magnitude bins between 20 and 26.5
  - 5 redshift bins between 0.6 and 3
  - 7 angular bins between 1.5 and 55 arc minutes
- MCMC estimation of the cluster concentration,  $c$ , and the cluster mass,  $M_l$
- **Clearly not working yet**
  - Biased
  - Errors underestimated



---

# Future work

- Estimate stacked cluster masses on mocks
- Incorporate dust models into the mock and our likelihood
- Estimate dust content and cluster masses with Redmapper clusters and HSC galaxies