



ID de Contribution: 53

Type: Non spécifié

## First-forbidden $\beta$ -decay study in the pnQRPA approach

jeudi 21 octobre 2021 14:00 (23 minutes)

First-forbidden beta decays play an important role in several domains of physics. First, in astrophysics, where nuclear data such as the half-life govern stellar evolution and nucleosynthesis [1]. Second, they are of interest for nuclear reactors physics as first highlighted in 2014 [2]. In first-forbidden  $\beta$ -decays, the form factor of the leptonic spectra are not equal to one as for allowed decays. It has been shown that it could have a non negligible impact on the shape of the antineutrino energy spectra. Among the models developed since then, which do not all tend to agree [3, 4, 5, 6], some even state that it could solve the reactor antineutrino shape anomaly.

New theoretical calculations of the first-forbidden form factors associated to summation calculations [7] and dedicated experimental measurements would be useful to corroborate or negate already existing predictions.

Charge-exchange excitations corresponding to beta-decay first forbidden transitions in nuclei have been studied in the self-consistent *proton-neutron quasiparticle random-phase approximation* (pnQRPA) using the finite-range Gogny interaction [8]. No parameters beyond those included in the effective nuclear force are included. Axial deformations are taken into account for both the ground state and charge-exchange excitations. With this formalism, nuclear matrix elements have been computed for operators derived from the multipole expansion of the weak current [9]: spin-dipole, anti-analog dipole and pseudoscalar-axial vector and tensor-polar vector operators. Those operators come to complete the already existing Fermi and Gamow-Teller operators already considered in Ref. [8] in order to have a simultaneous description of the allowed and first-forbidden  $\beta$ -decays.

At this conference, first results of the charge-exchange operators will be presented for both spherical and axially deformed nuclei with a comparison to other theoretical models.

### References:

- [1] M. Arnould, S. Goriely, and K. Takahashi. The r-process of stellar nucleosynthesis: Astrophysics and nuclear physics achievements and mysteries. *Phys. Rept.*, 450:97–213, 2007.
- [2] A. C. Hayes, J. L. Friar, G. T. Garvey, Gerard Jungman, and Guy Jonkmans. Systematic Uncertainties in the Analysis of the Reactor Neutrino Anomaly. *Phys. Rev. Lett.*, 112:202501, 2014.
- [3] Dong-Liang Fang and B. Alex Brown. Effect of first forbidden decays on the shape of neutrino spectra. *Phys. Rev. C*, 91(2):025503, 2015. [Erratum: *Phys.Rev.C* 93, 049903 (2016)].
- [4] X. B. Wang and A. C. Hayes. Weak magnetism correction to allowed  $\beta$  decay for reactor antineutrino spectra. *Phys. Rev. C*, 95(6):064313, 2017.
- [5] X. B. Wang, J. L. Friar, and A. C. Hayes. Nuclear Zemach moments and finite-size corrections to allowed  $\beta$  decay. *Phys. Rev. C*, 94(3):034314, 2016.
- [6] J. Petkovic, T. Marketin, G. Martinez-Pinedo, and N. Paar. Self-consistent calculation of the reactor antineutrino spectra including forbidden transitions. *J. Phys. G*, 46(8):085103, 2019.
- [7] M. Estienne et al. Updated Summation Model: An Improved Agreement with the Daya Bay Antineutrino Fluxes. *Phys. Rev. Lett.*, 123(2):022502, 2019.

[8] M. Martini, S. Peru, and S. Goriely. Gamow-Teller strength in deformed nuclei within the self-consistent charge-exchange quasiparticle random-phase approximation with the Gogny force. *Phys. Rev. C*, 89(4):044306, 2014.

[9] Aage Bohr and Ben R Mottelson. *Nuclear Structure*. World Scientific Publishing Company, 1998.

**Auteur principal:** BELOEUVRE, Arthur (CNRS-IN2P3)

**Co-auteurs:** MARTINI, Marco (IPSA and LPNHE); PERU, Sophie (CEA); ESTIENNE, Magali (IPHC - CNRS/IN2P3); FALLOT, Muriel (Université de Nantes)

**Orateur:** BELOEUVRE, Arthur (CNRS-IN2P3)

**Classification de Session:** Nuclear Physics & Interdisciplinaire

**Classification de thématique:** Nuclear Physics & Interdisciplinaire