

Two-body gravitational scattering in effective field theory


Pierre Vanhove

IPHT
Saclay

Planck 2022, Paris, France

based on work done in collaboration with
N.E.J. Bjerrum-Bohr, Poul Damgaard, Ludovic Planté, Stavros Mougiakakos

Einstein theory of gravity is the main paradigm for understanding the structure and dynamic of our observable Universe



Black holes, compact stars and gravitational waves are amongst the most spectacular predictions of general relativity and natural probes of the fundamental principles of Einstein's theory and its extension

“Gravitational waves enable tests of general relativity in the highly dynamical and strong-field regime. Using events detected by LIGO-Virgo up to 1 October 2019, we evaluate the consistency of the data with predictions from the theory. [...] *We find no evidence for new physics beyond general relativity, for black hole mimickers, or for any unaccounted systematics.*”¹

¹ [LIGO and Virgo collaboration 2010.14529]

Gravity effective field theories

Still it is important to have a better understanding of how Einstein gravity fits within our current EFT approach of particle physics

GW physics requires that one resolves conceptual issues at 5 post-Newtonian order (and higher) and understand better the radiation

By embedding the *classical* Einstein gravity effects in a quantum EFT framework we will shed a new light on subtle gravitational effects

We will be working in the context of an effective field theory assuming :

- ▶ Standard QFT (local, unitary, lorentz invariant, ...)
- ▶ The low-energy DOF: graviton, usual matter fields
- ▶ Standard symmetries: General relativity as we know it

$$\mathcal{S}_{\text{eff}} = \mathcal{S}_{\text{eff}}^{\text{gravity}} + \mathcal{S}_{\text{eff}}^{\text{matter}}$$

Classical gravity from quantum amplitudes

THE GENERATION OF GRAVITATIONAL WAVES. IV. BREMSSTRAHLUNG*†‡

SÁNDOR J. KOVÁCS, JR.

W. K. Kellogg Radiation Laboratory, California Institute of Technology

AND



KIP S. THORNE

Center for Radiophysics and Space Research, Cornell University; and
W. K. Kellogg Radiation Laboratory, California Institute of Technology

Received 1977 October 21; accepted 1978 February 28



g) The Feynman-Diagram Approach

Any classical problem can be solved quantum-mechanically; and sometimes the quantum solution is easier than the classical. There is an extensive literature on the Feynman-diagram, quantum-mechanical treatment of gravitational bremsstrahlung (e.g., Feynman 1961, 1963; Barker, Gupta, and Kaskas 1969; Barker and Gupta



Analytic expressions for the classical gravitational two-body interactions are obtained using techniques from quantum scattering amplitudes

Precison graviton from amplitudes




We need to understand well the coalescence regime in almost all relevant modified gravity theories

The long inspiral phase makes up the major portion of GW signal

EFT are designed for high precision (analytic!) computations applied to the inspiral phase, which is the largest part of the detected signal!


Classical Gravity from Quantum Scattering



One important **new** insight is that the **classical** gravitational two-body interactions (conservative and radiation) can be extracted from **quantum scattering amplitudes**

$$\mathcal{M}(p_1 \cdot p_2, q^2) = \begin{array}{c} \text{Diagram of a central circle with four external lines: } p_1' \text{ (top-left), } p_2' \text{ (top-right), } p_1 \text{ (bottom-left), and } p_2 \text{ (bottom-right).} \end{array} = \sum_{L=0}^{+\infty} G_N^{L+1} \mathcal{M}^{L\text{-loop}}$$

Classical physics from quantum loops



In the limit $\hbar, q^2 \rightarrow 0$ with $\underline{q} = \frac{q}{\hbar}$ fixed at each loop order of the quantum amplitude has the Laurent expansion² $\gamma = \frac{p_1 \cdot p_2}{m_1 m_2}$ and $q^2 = (p_1 - p'_1)^2$

$$\mathfrak{M}_L(\gamma, \underline{q}^2, \hbar) = \frac{\mathcal{M}_L^{(-L-1)}(\gamma, \underline{q}^2)}{\hbar^{L+1} |\underline{q}|^{\frac{L(4-D)}{2} + 2}} + \dots + \frac{\mathcal{M}_L^{(-1)}(\gamma, \underline{q}^2)}{\hbar |\underline{q}|^{\frac{L(4-D)}{2} + 2 - L}} + \mathcal{O}(\hbar^0)$$

- ▶ At all loop orders there is a classical contribution of order $1/\hbar$
- ▶ **classical gravity contributions** are determined by the **unitarity**
- ▶ The quantum amplitude has IR and UV divergences. But the classical amplitude is finite

²

[Iwasaki; Holstein, Donoghue; Bjerrum-Bohr, Damgaard, Planté, Vanhove; Kosower, Maybee, O'Connell]

Classical physics from loops : \hbar counting

One can understand this \hbar scaling by remembering that the mass in the propagator is actually the wave-length $m/(\hbar c)$

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \vec{\nabla}^2 + \frac{m^2 c^2}{\hbar^2} \right) \phi(t, \vec{x}) = 0$$

At the $L + 1$ PM order, the two-body scattering amplitude scales with the masses as

$$\mathcal{M}_L(\gamma, q^2) = \frac{G_N^{L+1} m_1^2 m_2^2}{q^{2 + \frac{(2-D)L}{2}}} \sum_{i=0}^L c_{L-i+2, i+2}(\gamma) m_1^{L-i} m_2^i$$

This piece will emerge from a L quantum amplitude as follows

$$\mathcal{M}^L \Big|_{\text{classical}} \propto \frac{m_1^2 m_2^2}{|q|^2} \hbar^{L-1} G_N^{L+1} \sum_i \left(\frac{m_1 c}{\hbar} \right)^{L-i} \left(\frac{m_2 c}{\hbar} \right)^i \propto \frac{\mathcal{M}_L(\gamma, q^2)}{\hbar}$$

One graviton exchange : tree-level amplitude



$$\mathfrak{M}_1 = -16\pi G_N \hbar \frac{2(p_1 \cdot p_2)^2 - m_1^2 m_2^2 - |\hbar \underline{\vec{q}}|^2 (p_1 \cdot p_2)}{|\hbar \underline{\vec{q}}|^2}$$

The \hbar expansion of the tree-level amplitude

$$\mathfrak{M}_1 = \frac{\mathcal{M}_1^{(-1)}(p_1 \cdot p_2)}{\hbar |\underline{q}|^2} + \hbar 4\pi G_N p_1 \cdot p_2$$

The higher order in q^2 are quantum with powers of \hbar

The *classical* potential is obtained by taking the 3d Fourier transform

$$E_i = \sqrt{p_i^2 + m_i^2}$$

$$\mathcal{V}_1(p_1 \cdot p_2, r) = \int \frac{d^3 \underline{\vec{q}}}{(2\pi)^3} \frac{\mathcal{M}_1^{(-1)}(\underline{\vec{q}}) e^{i\underline{\vec{q}} \cdot \underline{\vec{r}}}}{4E_1 E_2} = \frac{G_N}{E_1 E_2} \frac{m_1^2 m_2^2 - 2(p_1 \cdot p_2)^2}{r}$$

Exponentiation of the \hat{S} -matrix

Using an exponential representation of the \hat{S} matrix³

$$\hat{S} = \mathbb{I} + \frac{i}{\hbar} \hat{T} = \exp\left(\frac{i\hat{N}}{\hbar}\right)$$

doing the Dyson expansion with the conservative and radiation part

$$\hat{T} = G_N \sum_{L \geq 0} G_N^L \hat{T}_L + G_N^{\frac{1}{2}} \sum_{l \geq 0} G_N^L \hat{T}_L^{\text{rad}}, \quad \hat{N} = G_N \sum_{L \geq 0} G_N^L \hat{N}_L + G_N^{\frac{1}{2}} \sum_{l \geq 0} G_N^L \hat{N}_L^{\text{rad}}$$

The classical radial action $\hat{N}^{\text{classical}}$ does not have any \hbar . The higher power of $1/\hbar$ more singular than the classical are needed for the consistency of the full quantum amplitude and the correct exponentiation of the amplitude

$$\mathfrak{M}_L(\gamma, \underline{q}^2, \hbar) = \frac{\mathcal{M}_L^{(-L-1)}(\gamma, \underline{q}^2)}{\hbar^{L+1} |\underline{q}|^{\frac{L(4-D)}{2} + 2}} + \dots + \frac{\mathcal{M}_L^{(-1)}(\gamma, \underline{q}^2)}{\hbar |\underline{q}|^{\frac{L(4-D)}{2} + 2 - L}} + \mathcal{O}(\hbar^0)$$


³[Damgaard, Planté, Vanhove]

Velocity cuts and classical radial action

The precise mass dependence allows to identify the contribution to focus in the amplitude

At L -loop the classical part arises from imposing L -delta functions on the massive propagators

$$\mathcal{M}_L(\gamma, |\underline{q}|^2) \Big|_{\text{classical}} = \frac{1}{\hbar} \frac{\mathcal{M}_L^{(L-2)}(\gamma, D)}{|\underline{q}|^{2 + \frac{(2-D)L}{2}}} \implies N_L^{\text{classical}}(\gamma, D)$$



In practice, we need only evaluate matrix elements in the soft \underline{q}^2 -expansion, this means that we expand genuine unitarity cuts around the velocity cuts

Classical black hole metric from quantum amplitudes

Quantum Tree Graphs and the Schwarzschild Solution

M. J. Duff*

Physics Department, Imperial College, London SW7, England

(Received 7 July 1972)

I. INTRODUCTION

In an attempt to find quantum corrections to solutions of Einstein's equations, the question naturally arises as to whether the $\hbar \rightarrow 0$ limit of the quantum theory correctly reproduces the classical results. Formally, at least, the correspondence between the tree-graph approximation to quantum field theory and the classical solution of the field equations is well known,¹ i.e., the classical field produced by an external source serves as the generating functional for the connected Green's functions in the tree approximation, the closed-loop contributions vanishing in the limit $\hbar \rightarrow 0$. The purpose of this paper is to present an explicit calculation of the vacuum expectation value (VEV) of the gravitational field in the presence of a spherically symmetric source and verify, to second order in perturbation theory, that the result is in agreement with the classical Schwarzschild solution of the Einstein equations. This would appear to be the first step towards tackling the much more ambitious program of including the radiative quantum corrections.


In 1973 Duff asked the question about the classical limit of quantum gravity.^a He showed how to reproduce the Schwarzschild black hole metric from quantum tree graphs to G_N^3 order

The double expansion in G_N and \hbar give a new perspective on the classical limit of gravitational scattering amplitudes

^aM. J. Duff, "Quantum Tree Graphs and the Schwarzschild Solution," Phys. Rev. D **7** (1973), 2317-2326

Black hole metric from amplitudes

$$\frac{GM}{r} + \left(\frac{GM}{r}\right)^2 + \left(\frac{GM}{r}\right)^3 + \left(\frac{GM}{r}\right)^4$$




- ▶ The **tree skeleton graphs** are the one computed by Duff
- ▶ **Reproduces the Schwarzschild-Tangherlini metric in $d \geq 4$ dimensions**⁵

⁵[Mougiakakos, Vanhove]

The scattering angle

One important observable that allows to analytically continue from the scattering regime to the bound state regime is the scattering angle



Scattering angle from the (classical) radial action

$$N_{4\text{PM}}(\gamma, |b|) = \int_{\mathbb{R}^2} e^{iq \cdot b} \frac{N_{4\text{PM}}(\gamma, \underline{q}^2)}{4m_1 m_2 \sqrt{\gamma^2 - 1}} \frac{d^2 \underline{q}}{(2\pi)^2}$$

as $\chi_{4\text{PM}}(\gamma) = -\partial N_{4\text{PM}}(\gamma, J)/\partial J$ with the angular momentum $J = m_1 m_2 \sqrt{\gamma^2 - 1} b/E_{\text{C.M.}}$.

$$\begin{aligned} \frac{\chi}{2} \Big|_{1\text{PM}+2\text{PM}} &= \frac{(2\gamma^2 - 1)}{\gamma^2 - 1} \left(\frac{G_N m_1 m_2}{J} \right) \\ &\quad + \frac{3\pi(m_1 + m_2)(5\gamma^2 - 1)}{8(m_1^2 + m_2^2 + 2m_1 m_2 \gamma)} \left(\frac{G_N m_1 m_2}{J} \right)^2 \end{aligned}$$

Angle for a test mass in the Schwarzschild black hole of mass $M = m_1 + m_2$.

The 3PM scattering angle

$$\begin{aligned} \frac{\chi}{2} \Big|_{3\text{PM}} &= \left(\frac{G_N m_1 m_2}{J} \right)^3 \sqrt{\gamma^2 - 1} \left(\frac{(64\gamma^6 - 120\gamma^4 + 60\gamma^2 - 5)}{3(\gamma^2 - 1)^2} \right. \\ &\quad - \frac{4m_1 m_2}{3\mathcal{E}_{\text{C.M.}}^2} \gamma (14\gamma^2 + 25) + \frac{4m_1 m_2 (3 + 12\gamma^2 - 4\gamma^4) \operatorname{arccosh}(\gamma)}{\mathcal{E}_{\text{C.M.}}^2 \sqrt{\gamma^2 - 1}} \\ &\quad \left. + \frac{2m_1 m_2 (2\gamma^2 - 1)^2}{\mathcal{E}_{\text{C.M.}}^2 \sqrt{\gamma^2 - 1}} \left(-\frac{11}{3} + \frac{d}{d\gamma} \left(\frac{(2\gamma^2 - 1) \operatorname{arccosh}(\gamma)}{\sqrt{\gamma^2 - 1}} \right) \right) \right) \end{aligned}$$

At 3PM (two-loop) new phenomena arise

- ▶ The **conservative part** deviates from Schwarzschild as we have contributions which depends (linearly) on the relative mass⁶


$$\nu = \frac{m_1 m_2}{(m_1 + m_2)^2}$$

- ▶ And the important **Radiation-reaction terms**⁷

⁶[Damour; Bern et al.; di Vecchia et al.; Bjerrum-Bohr et al.]

⁷[Damour; di Vecchia et al.; Bjerrum-Bohr et al.]

Radiation reaction



The problem of **radiation reaction** has been one of the fundamental theoretical issues in general relativity. This is a needed contribution to match the binary-pulsar observations.

At 3PM a consistent derivation of radiation-reaction was missing. The amplitude approach clarified that

- ▶ The radiation-reaction from the soft region of the amplitude (not in the potential region of⁸)
- ▶ The radiation-reaction is needed for restoring a smooth continuity between the non-relativistic, relativistic and ultra-relativistic regimes⁹

The complete classical scattering amplitude gives a clear-cut unified and unambiguous resolution of these issues at 3PM¹⁰

⁸ [Bern et al.]

⁹ [Damour, Veneziano et al.]

¹⁰ [Bjerrum-Bohr, Damgaard, Planté, Vanhove]

Post-Minkowskian expansion

The advantages of the scattering amplitude approach is that

- ▶ No resummation in velocity is needed as this is automatically given by the relativistic expression in the relative velocity u

$$\mathfrak{M}^{\text{GR}}(p_1 \cdot p_2, q) = \sum_{m,n} G_N^n u^m c^{(n,m)}(q) \quad u^2 \sim \frac{G_N(m_1 + m_2)}{r} \ll 1$$

Compare to post-Newtonian expansion known up to 4PN

- ▶ One only gets velocities no higher derivative : acceleration terms & c in the Lagrangian are removed by appropriate coordinate transformation. No need to use equation of motions
- ▶ The amplitude gives a way to analyse the transition between the small velocity regime and the ultra-relativistic regime and the effect of gravitational radiation¹¹

¹¹[Damour; di Vecchia, Heisenber, Russo, Veneziano]

Beyond Einstein gravity

The same formalism can be used to include effects beyond Einstein gravity
From the exponentiation of the S -matrix $S \simeq \exp(N(J, E))$ one can include corrections beyond the classical Einstein gravity and study bulk causality and how classical observables are modified

- ▶ Star light bending angle¹²

$$\theta_s \simeq \frac{\partial N(J, E)}{\partial J} \simeq \frac{4GM}{b} + \frac{15\pi}{4} \left(\frac{GM}{b} \right)^2 + \frac{8bu^S - 48 \log(b/b_0) G\hbar M}{\pi} \frac{1}{b^3}$$

- ▶ Bulk causality and UV completion can be studied using the Shapiro time delay/advance¹³

$$\delta T = 2 \frac{\partial N(J, E)}{\partial E} \geq 0$$

Causality is preserved thanks to various positivity constraints on Wilson coefficients

¹²[Bjerrum-Bohr, Donoghue, Holstein, Planté, Vanhove]

¹³[Donoghue et al.; Camanho et al. ; Arkani-Hamed et al.; Bellazini et al.; Huber et al.]

This approach gives a new understanding on the relation between classical (Einstein) general relativity and the quantum theory of gravity

- 1 The 2-body gravitational scattering amplitude leads to the classical observables : potential, scattering angle, and radiation
- 2 The amplitude approach is much simpler than the traditional approach from solving Einstein's equation, and analytic relativistic expressions. The velocity cut method is a very efficient method for extracting the classical part
- 3 This is a very useful framework for studying subtle effects like radiation-reactions and memory effects where subtle non-linear effects arise from 5PN order [Blanchet; Damour; ...]
- 4 The approach applies to any EFT of gravity where one can compute amplitudes. Therefore this is a powerful approach to derive new constraints for modified gravity scenarios.