The Dark Dimension and the Swampland

Cumrun Vafa Harvard University

Planck 2022 Ancienne Ecole Polytechnique

Based on work W Michiel Monicorc ama Irana Valantia

Consistent theories of quantum gravity:

Very restrictive

Consistency of Att insufficient.

Naive Naturalness ideas fail

Swampland Program

Swampland Program aims to find criteria to distinguish consistent vs. inconsistent act. These criteria replace the notion of naturalness.


The aim of this talk is to combine Swampland criteria together with observations in the context of the "cosmological hierarchy problem"

 $\Lambda \sim 10^{-122}$

to show that we live in a specific corner of the QG landscape.

Plan for the talk: 1-Swampland criteria: Distance/Duality conjecture 2-Application to observed Universe 3-Phenomenological aspects and a unification of hierarchies: Mesoscopic dimension, neutrino masses, Higgs mass, prediction of a new UV scale, ...

Distance/Duality Conjecture [OV '06]; [LPV '19]:



Moreover the tower of light states is either a tower of KK modes, or light string states. Strong evidence from string theory (The String emergence proposal" In the context of dS/AdS the cosmological constant plays the role as an effective field $\Lambda=\pm e^{-\phi}$ thus we have a tower of states whose masses go as


 $m \sim |\Lambda|^{\alpha}$

WARETE

$$\frac{1}{d} \le \alpha \le \frac{1}{2} \quad \text{for } \Lambda > 0$$

(Upper range Higuchi bound, Lower range 1-Loop vacuum energy) Note that this is surprising from EFT: Why do light towers alone dictate Λ ? Example: Non-SUSY $O(16) \times O(16)$ strings in 10d:

 $V_{1-loop} \sim m_s^{10}$ by magic of modular invariance only light tower of string modes contribute. The contribution of higher mass states are screened as they are redundant dof. The sign is positive because there are more light fermions.



Application to our Universe $\Lambda \sim 10^{-122}$


This is an extreme range in field space. We thus predict using distance conjecture that there is either a tower of light string or KK modes whose mass scales as

$$m \sim |\Lambda|^{\alpha}$$
 with $\frac{1}{4} \leq \alpha \leq \frac{1}{2}$.

This in particular means gravity gets modified at the scale of m. The only possibility given the observations that Newtonian force law works up to about $30\mu m$ is the lower bound $\alpha = \frac{1}{d} = \frac{1}{4}$

 $\lambda m = \Lambda^{\frac{1}{4}}$

If $\lambda=1$, this would give $m^{-1}\sim 88\mu m$ which is ruled out. We now estimate $\lambda\sim 10^{-1}-10^{-3}$



Moreover for the asymptotic growth to have set in and not to lead to change in exponent of m, h cannot be too small, i.e. $\lambda \geq m^{\frac{1}{2}}$, which leads to $\lambda \sim 10^{-1} - 10^{-3}$ and $m^{-1} \sim (0.1 - 10) \mu m$

KK Comer or string tower?

Cannot be a string tower, effective theory of gravity valid far above ev

Muse de a kk comer:

How many extra mesoscopic dimensions? Emission of the KK modes leaves a trace in neutron stars and supernova explosions (can potentially cool a proto-neutron star too fast to be compatible with observations if the tower is too light).

This leads to [Hannestad et.al. '20] extra dimensions with length scale in the micron range to be ruled out except 1 extra dimension!

- e For the case of a single extra dimension: L < 44 µm
- e for the case of two extra dimensions: L < .00016 µm

.... So we predict

The Dark Dimension: One extra mesoscopic dimension of length .1-10 micron! This leads to a fundamental Planck scale in higher dimension $\hat{M}=m^{\frac{1}{3}}M_{pl}^{\frac{2}{3}}\sim 10^9-10^{10}GeV$

Comparing to large extra dimension scenarios [A'90,ADD'98,DDG'98,RS'99] the motivation and results are very different: Those approaches bring down the Planck scale to the weak scale to solve the EW hierarchy for example in ADD the number of large extra dimensions is 4 (or more) with Length scale $10^{-12}m$ (or smaller). We are motivated instead by Cosmological hierarchy and Swampland ideas (consistency of QG) and the only option is 1 extra dimension in the micron range.

Phenomenological aspects

GUT brane: Should be localized in the mesocopic dimension for its coupling not to be too weak:

$$\frac{1}{e^2} = V_{SM} \hat{M}^4 \sim 10^2$$

(A nice example is F-theory GUT model:)

$l_{ m GUT}$	
	1

Neutrinos: Given that the bulk must have extra fermions it is natural to identify them as sterile neutrinos 3-components of which can act as right-handed neutrino to give active neutrinos a mass. Indeed this has already been studied in the context of LED [ADDM-R,DDG]. However the scenario we have is more in line with [Carena et.al.'17]:

$$\mathcal{M} = \begin{pmatrix} 0 & \frac{y\langle H \rangle}{\sqrt{l\hat{M}}} \\ \frac{y\langle H \rangle}{\sqrt{l\hat{M}}} & \frac{1}{l} \end{pmatrix} \text{ Leads to } m_{\nu} = \frac{y^2 < H^2 >}{\hat{M}}$$
 With $y \sim 10^{-2} - 10^{-3}$

Assuming that bulk dynamics and CUT dynamics communicate and avoid significant hierarchy in neutrino masses (sterile and active neutrinos having similar masses) this would fix the Higgs vev:

$$\frac{y^2 \langle H \rangle^2}{\hat{M}} \sim \frac{1}{l} \Rightarrow \langle H \rangle \sim \frac{1}{y} \sqrt{\frac{\hat{M}}{l}} \sim \frac{\Lambda^{1/6} M_{pl}^{1/3}}{y \lambda^{2/3}} \sim 10 - 10^3 \, GeV$$

Instability in Higgs potential at $10^{11} GeV$ may be avoided in our scenario decause that is where we hit the higher Planck scale and we cet new cowers of charaed states and black holes.

Other Phenomenological aspects:
-Neutrino oscillations (SBL anomalies with eV sterile neutrino Hannestad et.al.+many other works)

-Ultra High Energy cosmic ray cutoff is close to M. GZK Limit; any observable effects? -Ho tension resolution by the "Early Dark Energy" scenarios may be relevant here (scalar radion field playing a key role). Work in progress with Gonzalo, Montero, Objed and Valenzuela on cosmological aspects.

Summary

Small dark energy + Swampland + observations uniquely lead to a single mesoscopic dimension The Dark Dimension in the micron range (we resisted the temptation of calling it the 'omicron'!).

Possible Unification of hierarchies (Dirac's dream): $l \sim \lambda \Lambda^{-1/4}; \quad m_{\nu} \sim \lambda^{-1} \Lambda^{1/4}$ $\langle H \rangle \sim y^{-1} \lambda^{-2/3} \Lambda^{1/6} M_{pl}^{1/3}$ $\hat{M} \sim \lambda^{-1/3} \Lambda^{1/12} M_{pl}^{2/3}$

Easily falsifiable: improvement on the precision measurement of deviation from Newton's law by a factor of 10-100. Or improvement of astrophysical bounds.

Coincidence of many interesting phenomenological aspects!