

Neutrino Mass and the Early Universe: Dark Matter and Leptogenesis



Bowen Fu


The 24th International Conference From the Planck Scale to the Electroweak Scale

Outline

- ❖ Neutrino portal dark matter in type I seesaw model: limitation
- ❖ Type Ib seesaw model
 - Neutrino portal dark matter
 - Resonant leptogenesis


Neutrino Portal Dark Matter

- General neutrino portal: $y_i \phi \bar{\chi} N_i$ the dark particles are charged under a Z_2 symmetry



- heavy scalar scenario: $\phi \rightarrow \chi N_i$

- Freeze-in production of dark matter:



- ν -Yukawa dominance: sizeable Y


Neutrino Portal Dark Matter in the Littlest Seesaw model

- ❖ Littlest Seesaw model: a highly predictive version of type I seesaw model with 2 RHNs
- ❖ ν -Yukawa interaction can dominate dark matter production when the RHN mass is above 4 TeV
Chianese, King [1806.10606](#)
- ❖ Leptogenesis in the Littlest Seesaw model: $M_{R1} = 5.1 \times 10^{10}$ GeV, $M_{R2} = 3.3 \times 10^{14}$ GeV
King, Sedgwick, Rowley [1808.01005](#)
- ❖ Production through graviton for superheavy particles Chianese, Fu, King [2009.01847](#)
- ❖ Nevertheless, a ν -Yukawa dominant region can be found, but it is very hard to be tested by experiments

**Q: Can we find a model where ν -Yukawa dominance can appear for GeV scale heavy neutrino?
And perhaps compatible with leptogenesis?**

Type Ib Seesaw Model


- ❖ Traditional type I seesaw mechanism (type Ia)



- ❖ At least 2 Majorana RH neutrinos + 1 Higgs
- ❖ 1 Yukawa coupling for each RH neutrino
- ❖ 2 free parameters after considering neutrino mass and mixing: M_{R1} and M_{R2}
- ❖ To have a sizeable coupling, the right-handed neutrino has to be above TeV scale

- ❖ Type Ib seesaw mechanism

Hernandez-Garcia, King (2019)



- ❖ 1 Dirac neutrino + 2 Higgs
- ❖ 1 Yukawa coupling for each Higgs
- ❖ 3 free parameters after considering neutrino mass and mixing: Y_1 , Y_2 and M_N
- ❖ One of Y_1 , Y_2 can be small while the other one is sizeable, providing GeV scale heavy neutrino

Neutrino Portal Dark Matter

Type Ib Seesaw Model with a Neutrino Portal

❖ Particles and symmetries

	Q_α	$u_{R\beta}$	$d_{R\beta}$	L_α	$e_{R\beta}$	Φ_1	Φ_2	N_{R1}	N_{R2}	ϕ	$\chi_{L,R}$
$SU(2)_L$	2	1	1	2	1	2	2	1	1	1	1
$U(1)_Y$	$\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{3}$	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	$-\frac{1}{2}$	0	0	0	0
Z_3	1	ω	ω	1	ω	ω	ω^2	ω^2	ω	ω	ω^2
Z_2	+	+	+	+	+	+	+	+	+	-	-


❖ Seesaw Lagrangian and neutrino portal $\mathcal{N} = (N_{R1}^c, N_{R2})$

Chianese, Fu, King [2102.07780](#)


$$\mathcal{L}_{\text{seesawIb}} = -Y_{1\alpha}^* \bar{L}_\alpha^c \Phi_1^* \mathcal{N}_L - Y_{2\alpha} \bar{L}_\alpha \Phi_2 \mathcal{N}_R - M_N \bar{\mathcal{N}}_L \mathcal{N}_R + \text{h.c.}$$

$$\mathcal{L}_{N_{R}\text{portal}} = y\phi \bar{\chi} \mathcal{N} + \text{h.c.}$$

❖ Freeze-in production of dark matter





Neutrino-Yukawa processes



Dark sector processes

Relation to Experiments



Chianese, Fu, King [2102.07780](https://arxiv.org/abs/2102.07780)

- ❖ 2 key parameters:
 - $\tan\beta$: the ratio of VEVs of the Higgs v_2/v_1
 - m_ϕ/m_χ : For hierarchical mass spectrum, the dark matter production depends on m_ϕ/m_χ
- ❖ U^2 : active-sterile neutrino mixing strength

- ❖ The strongest constraint is given by ν_μ mixing
- ❖ ν -Yukawa dominance is allowed above the coloured dashed lines
- ❖ Less constrained as $\tan\beta$ increases
- ❖ More constrained as m_ϕ/m_χ increases

Leptogenesis

Resonant Leptogenesis in Type Ib Seesaw Model

- ❖ An extended model with a superheavy third RHN and a scalar field



$$\mathcal{L}_{\text{seesawIb}} = -Y_{1\alpha}\bar{\ell}_\alpha\phi_1 N_{R1} - Y_{3\alpha}\bar{\ell}_\alpha\phi_2 N_{R3} - 2Y_{13}\bar{\xi}\overline{N_{R3}^c}N_{R1} - 2Y_{23}\bar{\xi}\overline{N_{R3}^c}N_{R2} \\ - M\overline{N_{R1}^c}N_{R2} - \frac{1}{2}M_3\overline{N_{R3}^c}N_{R3} + \text{h.c.}$$

Fu, King [2107.01486](#)

- ❖ After the scalar field gains a VEV, N_{R1} and N_{R2} gain mass splitting through mixing with N_{R3}

$$M_N = \begin{pmatrix} 0 & M & M_{13} \\ M & 0 & M_{23} \\ M_{13} & M_{23} & M_3 \end{pmatrix} \quad \Delta M_{12} = \frac{\Re[(M_{13} - M_{23})^2]}{2M_{33}}$$

- ❖ Type Ib seesaw mechanism can be realised effectively at low scale



Summary

- ❖ Features of Type Ib Seesaw Model
 - 3 free parameters after considering neutrino mass and mixing
 - Allow a connection between dark matter and neutrino physics through neutrino portal with GeV scale heavy neutrino
 - Resonant leptogenesis can be realised

Thank You!

Vector Portal

Type Ib Seesaw Model with a $U(1)'$ Symmetry

❖ Dirac neutrino — can be charged under gauge symmetry



❖ Particles and symmetries

	\mathcal{N}	$\chi_{L,R}$	ϕ
$SU(2)_L$	1	1	1
$U(1)_Y$	0	0	0
$U(1)'$	1	$\frac{1}{2}$	1

❖ Majorana dark matter candidate $y_\chi^L \bar{\phi} \bar{\chi}_L^c \chi_L + y_\chi^R \bar{\phi} \bar{\chi}_R^c \chi_R + h.c.$

❖ After ϕ gains a VEV, the $U(1)'$ symmetry is broken into a Z_2 symmetry, under which only χ is charged

❖ Freeze-out production of DM



Fu, King 2110.00588


Type Ib Seesaw Model with a $U(1)'$ Symmetry

- ❖ Fourth family of vector-like fermions

	$q_{L\alpha}$	$u_{R\beta}$	$d_{R\beta}$	$\ell_{L\alpha}$	$e_{R\beta}$	q_4	u_4	d_4	ℓ_4	e_4	Φ_1	Φ_2	\mathcal{N}	χ_R	ϕ
$SU(2)_L$	2	1	1	2	1	2	1	1	2	1	2	2	1	1	1
$U(1)_Y$	$\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{3}$	$-\frac{1}{2}$	-1	$\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{3}$	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	$-\frac{1}{2}$	0	0	0
$U(1)'$	0	0	0	0	0	-1	1	1	-1	1	1	-1	1	$\frac{1}{2}$	1


- ❖ Due to the $U(1)'$ charges of the Higgs doublets, the charged fermions can only gain mass from non-renormalisable operators $\bar{q}_{L\alpha} \Phi_2 u_{R\beta} \phi$, $\bar{q}_{L\alpha} \tilde{\Phi}_1 d_{R\beta} \phi$, $\bar{\ell}_\alpha \tilde{\Phi}_1 e_{R\beta} \phi$

- ❖ Fourth family of vector-like fermions: an example of up-type quark mass



Type Ib Seesaw Model with a U(1)' Symmetry


❖ Freeze-out production of DM



❖ Resonance in DM production when $M_{Z'} \sim 2m_\chi$

- Boltzmann suppression below the freeze-out temperature $T_f \sim m_\chi/20$

- Resonant amplitude
$$|\mathcal{M}|^2 \propto \frac{1}{(s - M_{Z'}^2)^2 + M_{Z'}^2 \Gamma_{Z'}^2}$$



Leptogenesis in Type Ib Seesaw Model

❖ In the minimal type Ib seesaw model, the correct asymmetry cannot be produced because the heavy neutrino mass is completely degenerate

❖ Minimal extension of the type Ib seesaw model: a pseudo-Dirac neutrino

$$\mathcal{L}_{\text{seesaw Ib}} = -Y_{1\alpha}\bar{\ell}_\alpha\phi_1 N_{R1} - Y_{2\alpha}\bar{\ell}_\alpha\phi_2 N_{R2} - M\overline{N_{R1}^c} N_{R2} - \Delta M\overline{N_{R2}^c} N_{R2} + \text{h.c.}$$

❖ In the Majorana basis, the phases of Yukawa couplings between RH neutrinos and Higgs bosons differ by $\pi/2$

❖ In a general theory with Yukawa interaction $e^{i\theta_{ik\alpha}} Y_{ik\alpha}\bar{\ell}_\alpha\phi_k n_{Ri}$

$$\left(\epsilon_{n_i}^{\text{wave-function}}\right)_{k\alpha} \propto \sum_{j \neq i} \sum_{l, \beta} Y_{ik\alpha} Y_{jk\alpha} Y_{il\beta} Y_{jl\beta} \sin(\theta_{ik\alpha} - \theta_{jk\alpha} + \theta_{il\beta} - \theta_{jl\beta})$$

❖ In the extension of the type Ib seesaw model with a superheavy third RHN and scalar field, an extra phase difference is developed as $(\theta_2 - \theta_1)$

$$\tan \theta_1 \simeq -\frac{1}{2MM_{33}} \Im \left[(M_{13} + M_{23})^2 \right], \quad \tan \theta_2 \simeq \frac{1}{2MM_{33}} \Im \left[(M_{13} - M_{23})^2 \right].$$