

Abstract

Materials with thermopiezoelectric properties have both theoretical and practical significance in solid-state physics and materials science. Thermopiezoelectric media exhibit coupling among the thermal, electric and elastic fields. Strain, electric displacement and entropy can be written in terms of stress, electric field and temperature. Investigation is concerned with the analysis of nonlinear thermopiezoelectricity using thermodynamic principles. Strain, electric displacement and entropy are expanded into Taylor series. Zeroth to eighth rank tensors are derived for describing material constants. Relationship of material constants of strain, electric displacement and entropy are obtained in thermopiezoelectricity. Due to intrinsic coupling behavior, thermopiezoelectric materials are widely used as sensors and actuators in sensing, actuation, and control of smart structures. Mathematical expressions of material constants may be useful for future investigation of the mechanics and physics of nonlinear thermopiezoelectricity.

Introduction

Mechanical, electrical, and thermal fields are coupled in thermopiezoelectricity. Nonlinear analysis of thermopiezoelectricity has been made for determining their material constants. In the previous studies, Das and co-workers (1998) introduced static, dynamic, and control characteristics of a nonlinear piezoelectric laminated beam subjected to mechanical, temperature, and electric excitations. Wang et al. (1999) studied the nonlinear electrothermal behavior piezoelectric ceramic in a wide electric field and frequency range. Zhou and Tian (2000) developed nonlinear dielectric hysteresis and active control of piezoelectric laminated circular spherical shells. Hall (2001) gave an overview of experimental evidence and understanding of nonlinear piezoelectricity, electric and ferroelectric relationships in piezoelectric ceramics.

Furthermore, Alay and Dokmeci (2002) described a nonlinear rod theory for high-frequency vibrations of thermopiezoelectric materials. Wierzbicki and Linnik (2003) analyzed material constants from zeroth to sixth rank tensors in nonlinear mechanical, electrical and thermal phenomena in piezoelectric crystals. Wagner (2004) presented nonlinear longitudinal vibrations of non-slender piezoelectric rods. Malygina and Chaudhuri (2005) deduced a generalized formulation for nonlinear dynamic analysis of piezoelectric structures. Blackham and Cain (2006) examined nonlinear piezoelectric resonance.

$$\begin{aligned} s_{ij} &= f(\sigma_{ij}, E_i, T) \quad (5) \\ \lambda_i &= f(\sigma_{ij}, E_i, T) \quad (6) \\ S &= f(\sigma_{ij}, E_i, T) \quad (7) \end{aligned}$$

Further, thermodynamic Gibbs function can be written as

$$dG = \frac{\partial G}{\partial \sigma_{ij}} d\sigma_{ij} + \frac{\partial G}{\partial E_i} dE_i + \frac{\partial G}{\partial T} dT \quad (8)$$

In equation (8), indices in the lower part of vertical line on the partial derivatives indicate the variable that may be held constant during differentiation.

Also, equations (4) and (9) yield

$$s_{ij} = -\frac{\partial G}{\partial \sigma_{ij}} \quad (9)$$

$$D_i = -\frac{\partial G}{\partial E_i} \quad (10)$$

$$S = -\frac{\partial G}{\partial T} \quad (11)$$

In equation (1), U , T and S are internal energy, temperature change and entropy, respectively. Subscript indices i, j can assume 1 to 3. It is noted that all other subscript indices denoting in other equations range from 1 to 3 unless otherwise specified. Strain, stress, electric field and electric displacement are denoted by ϵ_{ij} , σ_{ij} , E_i and D_i . Following Einstein's summation convention, the summation sign \sum is omitted. For example, its detail can be found in Water, Balafoutis and Kitzner (2006).

Differential form of Gibbs function is expressed as

$$dG = dU - s_i dE_i - \sigma_{ij} d\epsilon_{ij} - E_i dD_i - D_i dE_i - S dT \quad (2)$$

As far as the first and second law of thermodynamics are concerned, differential form of internal energy is represented by

$$dU = dE_i + E_i dD_i + \sigma_{ij} d\epsilon_{ij} + T dS \quad (3)$$

It follows from equations (2) and (3) that

$$dG = -s_i dE_i - \sigma_{ij} d\epsilon_{ij} - S dT \quad (4)$$

Strain, electric displacement and entropy can be written in the following forms

$$\epsilon_{ij} = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl}} \sigma_{kl} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial E_k} E_k + \frac{\partial^2 S}{\partial \sigma_{ij} \partial T} T \quad (14)$$

Material constants obtained from strain ϵ_{ij}

Taking equation (9), expressions of material constants can be obtained as

Second order elasticity constant (fourth rank tensor)

$$\frac{\partial^2 \epsilon_{ij}}{\partial \sigma_{kl} \partial \sigma_{mn}} = \frac{\partial^2}{\partial \sigma_{kl} \partial \sigma_{mn}} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijkl}^s \quad (15a)$$

In equation (15a) and all other equations of material constants, superscript indices are constant. It should be noted that superscript indices are not taken as powers.

Second order piezoelectric constant (third rank tensor)

$$\frac{\partial \epsilon_{ij}}{\partial E_k} = \frac{\partial}{\partial E_k} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijk}^s \quad (15b)$$

Second order thermo-piezoelectric constant (second rank tensor)

$$\frac{\partial \epsilon_{ij}}{\partial T} = \frac{\partial}{\partial T} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ij}^s \quad (15c)$$

Third order elasticity constant (sixth rank tensor)

$$\frac{\partial^3 \epsilon_{ij}}{\partial \sigma_{kl} \partial \sigma_{mn} \partial \sigma_{pq}} = \frac{\partial^3}{\partial \sigma_{kl} \partial \sigma_{mn} \partial \sigma_{pq}} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijklmn}^s \quad (15d)$$

Third order piezoelectric constant (fifth rank tensor)

$$\frac{\partial^2 \epsilon_{ij}}{\partial \sigma_{kl} \partial E_m} = \frac{\partial^2}{\partial \sigma_{kl} \partial E_m} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijklm}^s \quad (15e)$$

Third order thermo-piezoelectric constant (fourth rank tensor)

$$\frac{\partial \epsilon_{ij}}{\partial \sigma_{kl} \partial T} = \frac{\partial}{\partial \sigma_{kl} \partial T} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijkl}^s \quad (15f)$$

$$\text{Fourth order elasticity constant (eighth rank tensor)}$$

$$\frac{\partial^4 \epsilon_{ij}}{\partial \sigma_{kl} \partial \sigma_{mn} \partial \sigma_{pq} \partial \sigma_{rs}} = \frac{\partial^4}{\partial \sigma_{kl} \partial \sigma_{mn} \partial \sigma_{pq} \partial \sigma_{rs}} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijklmnrs}^s \quad (15g)$$

Fourth order piezoelectric constant (seventh rank tensor)

$$\frac{\partial^3 \epsilon_{ij}}{\partial \sigma_{kl} \partial \sigma_{mn} \partial E_p} = \frac{\partial^3}{\partial \sigma_{kl} \partial \sigma_{mn} \partial E_p} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijklmnp}^s \quad (15h)$$

Fourth order thermo-piezoelectric constant (sixth rank tensor)

$$\frac{\partial^2 \epsilon_{ij}}{\partial \sigma_{kl} \partial \sigma_{mn} \partial T} = \frac{\partial^2}{\partial \sigma_{kl} \partial \sigma_{mn} \partial T} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijklmnp}^s \quad (15i)$$

Third order electrostriction constant (fourth rank tensor)

$$\frac{\partial^2 \epsilon_{ij}}{\partial E_k \partial E_l} = \frac{\partial^2}{\partial E_k \partial E_l} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijkl}^e \quad (15j)$$

Third order electro-thermo-elastic constant (third rank tensor)

$$\frac{\partial \epsilon_{ij}}{\partial E_k \partial T} = \frac{\partial}{\partial E_k \partial T} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijk}^e \quad (15k)$$

Fourth order electrostriction constant (sixth rank tensor)

$$\frac{\partial^3 \epsilon_{ij}}{\partial \sigma_{kl} \partial \sigma_{mn} \partial E_p} = \frac{\partial^3}{\partial \sigma_{kl} \partial \sigma_{mn} \partial E_p} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijklmnp}^e \quad (15l)$$

Fourth order electro-thermo-elastic constant (fifth rank tensor)

$$\frac{\partial^2 \epsilon_{ij}}{\partial \sigma_{kl} \partial \sigma_{mn} \partial T} = \frac{\partial^2}{\partial \sigma_{kl} \partial \sigma_{mn} \partial T} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ijklmnp}^e \quad (15m)$$

Third order thermal expansion constant (second rank tensor)

$$\frac{\partial \epsilon_{ij}}{\partial T} = \frac{\partial}{\partial T} \left(\frac{\partial G}{\partial \sigma_{ij}} \right) = e_{ij}^t \quad (15n)$$

$$\frac{\partial^2 D_i}{\partial \sigma_{ij} \partial \sigma_{kl}} = \frac{\partial^2}{\partial \sigma_{ij} \partial \sigma_{kl}} \left(\frac{\partial G}{\partial E_i} \right) = e_{ijkl}^D \quad (16a)$$

Fourth order thermo-electro-elastic constant (third rank tensor)

$$\frac{\partial D_i}{\partial \sigma_{ij} \partial T} = \frac{\partial}{\partial \sigma_{ij} \partial T} \left(\frac{\partial G}{\partial E_i} \right) = e_{ij}^D \quad (16b)$$

Material constants obtained from entropy S

Using equation (11), the material constants can be expressed as

Second order piezoelectric constant (third rank tensor)

$$\frac{\partial S}{\partial \sigma_{ij} \partial E_k} = \frac{\partial}{\partial \sigma_{ij} \partial E_k} \left(-\frac{\partial G}{\partial T} \right) = -e_{ijk}^S \quad (17a)$$

Second order electric heat constant (first rank tensor)

$$\frac{\partial S}{\partial E_i} = \frac{\partial}{\partial E_i} \left(-\frac{\partial G}{\partial T} \right) = -e_i^S \quad (17b)$$

Second order thermal capacity constant (zeroth rank tensor)

$$\frac{\partial S}{\partial T} = \frac{\partial}{\partial T} \left(-\frac{\partial G}{\partial T} \right) = -c^S \quad (17c)$$

Third order piezo-electric constant (fourth rank tensor)

$$\frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} = \frac{\partial^2}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \left(-\frac{\partial G}{\partial T} \right) = -e_{ijklm}^S \quad (17d)$$

Third order thermo-electro-elastic constant (third rank tensor)

$$\frac{\partial S}{\partial \sigma_{ij} \partial T} = \frac{\partial}{\partial \sigma_{ij} \partial T} \left(-\frac{\partial G}{\partial T} \right) = -e_{ij}^S \quad (17e)$$

Third order thermal expansion constant (second rank tensor)

$$\frac{\partial S}{\partial T} = \frac{\partial}{\partial T} \left(-\frac{\partial G}{\partial T} \right) = -c^S \quad (17f)$$

$$\frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl}} = \frac{\partial^2}{\partial \sigma_{ij} \partial \sigma_{kl}} \left(-\frac{\partial G}{\partial T} \right) = -e_{ijkl}^S \quad (17g)$$

Fourth order piezoelectric constant (sixth rank tensor)

$$\frac{\partial^3 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial \sigma_{mn} \partial E_p} = \frac{\partial^3}{\partial \sigma_{ij} \partial \sigma_{kl} \partial \sigma_{mn} \partial E_p} \left(-\frac{\partial G}{\partial T} \right) = -e_{ijklmnp}^S \quad (17h)$$

Fourth order thermo-electro-elastic constant (fifth rank tensor)

$$\frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} = \frac{\partial^2}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \left(-\frac{\partial G}{\partial T} \right) = -e_{ijklmnp}^S \quad (17i)$$

Third order electrostriction constant (fourth rank tensor)

$$\frac{\partial^2 S}{\partial E_k \partial E_l} = \frac{\partial^2}{\partial E_k \partial E_l} \left(-\frac{\partial G}{\partial T} \right) = -e_{ijkl}^S \quad (17j)$$

Third order electro-thermo-elastic constant (third rank tensor)

$$\frac{\partial S}{\partial E_k \partial T} = \frac{\partial}{\partial E_k \partial T} \left(-\frac{\partial G}{\partial T} \right) = -e_{ijk}^S \quad (17k)$$

Fourth order electrostriction constant (sixth rank tensor)

$$\frac{\partial^3 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} = \frac{\partial^3}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \left(-\frac{\partial G}{\partial T} \right) = -e_{ijklmnp}^S \quad (17l)$$

Fourth order thermo-electro-elastic constant (fifth rank tensor)

$$\frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} = \frac{\partial^2}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \left(-\frac{\partial G}{\partial T} \right) = -e_{ijklmnp}^S \quad (17m)$$

Third order thermal capacity constant (zeroth rank tensor)

$$\frac{\partial S}{\partial T} = \frac{\partial}{\partial T} \left(-\frac{\partial G}{\partial T} \right) = -c^S \quad (17n)$$

Nonlinearity and linearity of thermopiezoelectricity

It is convenient to write nonlinear constitutive equations (19) to (21) in matrix form as

$$\begin{bmatrix} \epsilon_{ij} \\ D_i \\ S \end{bmatrix} = \begin{bmatrix} A & D & E \\ D & D & E \\ E & E & C \end{bmatrix} \begin{bmatrix} \sigma_{ij} \\ E_i \\ T \end{bmatrix} \quad (22)$$

In equation (22), A, D, E, C are tensor effects and D, E, C are conjugate coefficients respectively. The base and conjugate effects are given by

$$A = \frac{\partial^2 \epsilon_{ij}}{\partial \sigma_{kl} \partial \sigma_{mn}} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 \epsilon_{ij}}{\partial \sigma_{kl} \partial E_m} E_m + \frac{\partial^2 \epsilon_{ij}}{\partial \sigma_{kl} \partial T} T \quad (23a)$$

$$D = \frac{\partial^2 D_i}{\partial \sigma_{ij} \partial \sigma_{kl}} \sigma_{kl} + \frac{\partial^2 D_i}{\partial \sigma_{ij} \partial E_m} E_m + \frac{\partial^2 D_i}{\partial \sigma_{ij} \partial T} T \quad (23b)$$

$$E = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23c)$$

$$C = \frac{\partial^2 S}{\partial T \partial T} T^2 + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial \sigma_{ij} \partial T} \sigma_{ij} T \quad (23d)$$

$$D = \frac{\partial^2 D_i}{\partial \sigma_{ij} \partial \sigma_{kl}} \sigma_{kl} + \frac{\partial^2 D_i}{\partial \sigma_{ij} \partial E_m} E_m + \frac{\partial^2 D_i}{\partial \sigma_{ij} \partial T} T \quad (23e)$$

$$E = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23f)$$

$$C = e_{ijklmnp}^S \sigma_{kl} \sigma_{mn} + e_{ijklm}^S E_m + e_{ij}^S T \quad (23g)$$

$$D = \frac{\partial^2 D_i}{\partial \sigma_{ij} \partial \sigma_{kl}} \sigma_{kl} + \frac{\partial^2 D_i}{\partial \sigma_{ij} \partial E_m} E_m + \frac{\partial^2 D_i}{\partial \sigma_{ij} \partial T} T \quad (23h)$$

$$E = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23i)$$

$$F = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23j)$$

$$G = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23k)$$

$$H = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23l)$$

$$I = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23m)$$

$$J = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23n)$$

$$K = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23o)$$

$$L = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23p)$$

$$M = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23q)$$

$$N = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23r)$$

$$O = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23s)$$

$$P = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23t)$$

$$Q = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23u)$$

$$R = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23v)$$

$$S = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23w)$$

$$T = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23x)$$

$$U = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23y)$$

$$V = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23z)$$

$$W = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23aa)$$

$$X = \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial E_m} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial \sigma_{ij} \partial \sigma_{kl} \partial T} \sigma_{kl} \sigma_{mn} + \frac{\partial^2 S}{\partial E_i \partial E_j} E_i E_j + \frac{\partial^2 S}{\partial E_i \partial T} E_i T + \frac{\partial^2 S}{\partial T \partial T} T^2 \quad (23ab)$$

$$Y$$