

Edge modes and counting statistics in the 2d and 4d Quantum Hall effect

Jean-Marie Stéphan

Camille Jordan Institute & CNRS, University of Lyon 1, France

DPP-fermions, Lyon 2022

[Estienne & JMS, Phys. Rev. B 101, 2019]

[Estienne, Oblak & JMS, Scipost Physics 11, 2021]

[Estienne, JMS & Witczak-Krempa, Nat. Comm. 13, 2022]


Slides + Chalk

Outline

- 1 Simple 2d Integer Quantum Hall wavefunctions
- 2 Simple 4d Integer Quantum Hall wavefunctions
- 3 Counting statistics and related problems

Integer Quantum Hall effect

[von Klitzing, Dorda & Pepper 1980]



Resistivities

$$\rho_{xx} = 0 \quad , \quad \rho_{xy} = \frac{h}{e^2} \times \frac{1}{\text{integer} \pm 10^{-8}}$$

Hamiltonian in a magnetic field + trapping potential

$$H = \frac{1}{2} (\mathbf{p} - \mathbf{A})^2 + \frac{k}{2} (x^2 + y^2)$$

Symmetric gauge

$$\mathbf{A} = \frac{B}{2} \begin{pmatrix} -y \\ x \end{pmatrix}$$


naturally leads to single particle wave functions

$$\phi_m(z) = \frac{z^m}{\sqrt{\pi m!}} e^{-|z|^2/2}$$


with single particle energies $\epsilon_m = \hbar\omega m$ with $\omega = \frac{k}{B}$.

$$K_N(z, z') = \sum_{m=0}^{N-1} \frac{(z^* z')^m}{\pi m!} e^{-(|z|^2 + |z'|^2)/2}$$


Density profile




Two-point function (correlation kernel)




Two-point function (correlation kernel)




Two-point function (correlation kernel)




Two-point function (correlation kernel)




Two-point function (correlation kernel)




Two-point function (correlation kernel)




Two-point function (correlation kernel)



Two-point function (correlation kernel)



Two-point function (correlation kernel)



4d Quantum Hall effect

$(x, y, u, v) \in \mathbb{R}^4$. Simplest generalization of the previous model:

$$H = (\mathbf{p} - \mathbf{A})^2 + \frac{k}{2} (x^2 + y^2) + \frac{k'}{2} (u^2 + v^2)$$

with vector potential

$$\mathbf{A} = \frac{B}{2} \begin{pmatrix} -y \\ x \\ -v \\ u \end{pmatrix}$$

leads to ($z = x + iy$, $w = u + iv$ up to some units)

$$\phi_{m,n}(z, w) = \frac{z^m w^n}{\sqrt{\pi^2 m! n!}} e^{-(|z|^2 + |w|^2)/2}$$

with energies $\epsilon_{m,n} = \hbar(\omega m + \omega' n)$ with $\omega = \frac{k}{B}$, $\omega' = \frac{k'}{B}$

Generalizations of QHE to 4d predict anisotropic edge modes

[Zhang & Hu, Science 2001] [Karabali & Nair, Nucl. Phys. B 2002]

[Helvang & Polchinski, CRP 2003]

Possible experimental realisations by engineering synthetic dimensions (quasicrystals, internal states, photonics) [Kraus, Ringel & Zilberberg, PRL 2013]

[Price, Zilberberg, Ozawa, Carusotto & Goldman, PRL 2015]


[Lohse, Schweizer, Price, Zilberberg & Bloch Nature 2018]

...

$$K_N(z, w|z', w') = \sum_{m+\Delta n < N} \frac{(z^* z')^m (w^* w')^n}{\pi^2 m! n!} e^{-(|z|^2 + |w|^2 + |z'|^2 + |w'|^2)/2}$$


$$\Delta = \frac{\omega'}{\omega}$$

Edge modes on the torus




$$\Delta = 1$$

Edge modes on the torus




$$\Delta = 3/2$$

Edge modes on the torus




$$\Delta = 5/3$$

Edge modes on the torus




$$\Delta = 7/5$$


Edge modes on the torus




$$\Delta = 41/29$$

1d slices at $\beta = 0$ 

$$\Delta = 17/12$$

1d slices at $\beta = 0$ 

$$\Delta = \sqrt{2}$$

1d slices at $\beta = 0$ 

$$\Delta = \sqrt{2}$$

Counting statistics and related problems

Thank you!