



ID de Contribution: 13

Type: Non spécifié

Modeling of High Column Density systems in the Ly- α Forests Correlation Function

vendredi 6 mai 2022 09:45 (15 minutes)

The Lyman- α forest is detected as the series of absorption lines in the quasar spectra, caused by the Lyman- α transitions of neutral hydrogen in the low-density, high-redshift intergalactic medium (IGM). It is a biased continuous tracer of the quasi-linear matter density field, and the auto (cross) correlation function of the forests (with quasars) has been used to detect the Baryon Acoustic Oscillations (BAO) signal. The Damped Lyman- α System (DLAs) is one of the most important systematics in the Lyman- α BAO analysis. DLAs are strong absorption regions in Lyman-alpha forests caused by neutral hydrogen along the sightline with extremely high column densities, usually $\log(N_{\text{HI}}) \geq 20$. We present an accurate model to characterize the impact of DLAs on the measurement of the Lyman- α correlation function, as well as the BAO fitting.

Auteurs principaux: BALLAND, Christophe (LPNHE Paris - UPMC -ILP); TAN, ting; RICH, James (SPP); LE GOFF, Jean-Marc (CEA Saclay DRF/Irfu/DPhP)

Orateur: TAN, ting

Classification de Session: Neutrinos