

Heavy-meson transport coefficients in a thermal medium



Juan M. Torres-Rincon
Universitat de Barcelona
Institut de Ciències del Cosmos



in collaboration with
G. Montaña, À. Ramos and L. Tolos


DFG Deutsche
Forschungsgemeinschaft



Strong and ElectroWeak Matter 2022
IPhT, Saclay - Université Paris VI, Paris, Jun 20 – 24, 2022


Introduction: Heavy flavor

(A. Bazavov *et al.*, 1904.09951)



- ▶ Infer QCD properties at high temperatures through final state of RHICs
- ▶ Find clean and solid observables to connect detections to early stages
- ▶ **Hard Probes:** Jets, hard electromagnetic emission, heavy flavor (quarkonia, open-heavy flavor hadrons...)

Heavy quarks: formed at the initial stage of the collision (short formation time) and difficult to equilibrate along their evolution (large relaxation time)



Interactions in a thermal medium?

Transport coefficients?


Introduction: Thermal propagation

- ▶ Heavy mesons (D, D^*, \bar{B}, \dots) at $T < 150 \text{ MeV}/k_B$
- ▶ Interacting with an equilibrated light-meson gas ($\Phi = \{\pi, K, \bar{K}, \eta\}$)
- ▶ **Heavy-hadron mass is the dominant scale**

$$M_D \gg m_\Phi, T, \Lambda_{QCD}$$

- ▶ Picture: Brownian particle in a thermal bath
B. Svetitsky, Phys. Rev. D37, 484 (1988)
- ▶ Transport properties: (Heavy-flavor) **diffusion coefficient, D_s .**

$$\vec{j} = -D_s(T, \mu_i) \vec{\nabla} n$$



Effective field theory

Effective Lagrangian based on **chiral** and **heavy-quark spin-flavor** symmetries.

- ▶ **Chiral expansion** up to NLO
 - : also explicitly broken due to light-meson masses (π, K, \bar{K}, η).
- ▶ **Heavy-quark mass expansion** up to LO
 - : broken by heavy meson masses (D, D_s, D^*, D_s^*).

E.E. Kolomeitsev and M.F.M. Lutz *Phys.Lett. B582 (2004) 39*, J. Hofmann and M.F.M. Lutz *Nucl.Phys. A733 (2004) 142*,
F.K.Guo, C.Hanhart, S. Krewald, U.G. Meissner *Phys.Lett. B666 (2008) 251*, L.S. Geng, N. Kaiser, J. Martin-Camalich and W. Weise *Phys.Rev.D82,05422 (2010)*, L.M. Abreu, D. Cabrera, F.J. Llanes-Estrada and JMT-R. *Annals Phys. 326 (2011) 2737...*

$$\mathcal{L}_{\text{LO}} = Tr[\nabla^\mu \textcolor{red}{D} \nabla_\mu \textcolor{red}{D}^\dagger] - m_D^2 Tr[\textcolor{red}{D} \textcolor{red}{D}^\dagger] - Tr[\nabla^\mu \textcolor{red}{D}^{*\nu} \nabla_\mu \textcolor{red}{D}_\nu^{*\dagger}] + m_{D^*}^2 Tr[\textcolor{red}{D}^{*\mu} \textcolor{red}{D}_\mu^{*\dagger}]$$

$$+ ig Tr \left[\left(\textcolor{red}{D}^{*\mu} \textcolor{blue}{u}_\mu \textcolor{red}{D}^\dagger - \textcolor{red}{D} \textcolor{blue}{u}^\mu \textcolor{red}{D}_\mu^{*\dagger} \right) \right] + \frac{g}{2m_D} Tr \left[\left(\textcolor{red}{D}_\mu^* \textcolor{blue}{u}_\alpha \nabla_\beta \textcolor{red}{D}_\nu^{*\dagger} - \nabla_\beta \textcolor{red}{D}_\mu^* \textcolor{blue}{u}_\alpha \textcolor{red}{D}_\nu^{*\dagger} \right) \epsilon^{\mu\nu\alpha\beta} \right]$$

$$\textcolor{red}{D} = (D^0, D^+, D_s^+)$$

$$\nabla^\mu = \partial^\mu - \frac{1}{2}(\textcolor{blue}{u}^\dagger \partial^\mu \textcolor{blue}{u} + \textcolor{blue}{u} \partial^\mu \textcolor{blue}{u}^\dagger)$$

$$\textcolor{blue}{u}^\mu = i(\textcolor{blue}{u}^\dagger \partial^\mu \textcolor{blue}{u} - \textcolor{blue}{u} \partial^\mu \textcolor{blue}{u}^\dagger)$$

$$\textcolor{blue}{u} = \exp \left[\frac{i}{\sqrt{2}F} \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & K^0 \\ K^- & \bar{K}^0 & -\frac{2\eta}{\sqrt{6}} \end{pmatrix} \right]$$

Perturbative potential

Perturbative amplitude

$$\begin{aligned} V(s, t, u) = & \frac{C_0}{4f_\pi^2}(s - u) + \frac{2C_1}{f_\pi^2} h_1 + \frac{2C_2}{f_\pi^2} h_3(k_2 \cdot k_3) \\ & + \frac{2C_3}{f_\pi^2} h_5[(k \cdot k_3)(k_1 \cdot k_2) + (k \cdot k_2)(k_1 \cdot k_3)] \end{aligned}$$

f_π : pion decay constant

Isospin coefficients: fixed by symmetry


Low-energy constants: fixed by experiment
or by underlying theory

Z.-H. Guo *et al.* Eur. Phys. J.C79, 1, 13 (2019)

Elastic processes:

$D\pi$, DK , $D\bar{K}$, $D\eta$

$D_s\pi$, D_sK , $D_s\bar{K}$, $D_s\eta$ and their inelastic channels.



Resummation at finite temperature

At $T \neq 0$ **Imaginary Time Formalism** in a self-consistent approximation

$$D_i \quad D_j = D_i \quad D_j + D_i \quad D_k \quad D_j$$


$$T_{ij} = [1 - G_{D\Phi} V]_{ik}^{-1} V_{kj}$$

$$G_{D\Phi}(E, \vec{p}; T) = \int \frac{d^3 k}{(2\pi)^3} \int d\omega \int d\omega' \frac{S_D(\omega, \vec{k}; T) S_\Phi(\omega', \vec{p} - \vec{k}; T)}{E - \omega - \omega' + i\varepsilon} [1 + f(\omega, T) + f(\omega', T)]$$

$$S_D(\omega, \vec{k}; T) = -\frac{1}{\pi} \text{Im } \Delta_D(\omega, \vec{k}; T) = -\frac{1}{\pi} \text{Im} \left(\frac{1}{\omega^2 - \vec{k}^2 - m_D^2 - \Pi_D(\omega, \vec{k}; T)} \right)$$

$$\Pi_D(\omega_n, \vec{k}; T) = T \int \frac{d^3 p}{(2\pi)^3} \sum_m \Delta_\Phi(\omega_m - \omega_n, \vec{p} - \vec{k}) T_{D\Phi}(\omega_m, \vec{p})$$

$$D \rightarrow D + D \rightarrow D \rightarrow \pi$$



D -meson kinetic theory

Kinetic theory with off-shell effects



Kadanoff-Baym equations

L. Kadanoff, G.Baym, "Quantum statistical mechanics" 1962, P. Danielewicz, Annals Phys. 152, 239 (1984), W. Botermans and R. Malfliet, Phys. Rept. 198, 115 (1990), J.-P. Blaizot and E. Iancu, Nucl. Phys. B557, 183 (1999), J. Rammer "Quantum field theory of non-equilibrium states" (2007), W. Cassing, Eur. Phys. J.168, 3 (2009)

$$\overbrace{\left(k^\mu - \frac{1}{2} \frac{\partial \text{Re } \Pi^R(X, k)}{\partial k_\mu} \right) \frac{\partial iG_D^<(X, k)}{\partial X^\mu}}^{\text{Advective term}} = \underbrace{\frac{1}{2} i\Pi^<(X, k) iG_D^>(X, k)}_{\text{Gain term}} - \underbrace{\frac{1}{2} i\Pi^>(X, k) iG_D^<(X, k)}_{\text{Loss term}}$$

Kadanoff-Baym ansatz:

$$iG_D^<(X, k) = 2\pi S_D(X, k) f_D(X, k^0)$$

$$iG_D^>(X, k) = 2\pi S_D(X, k) [1 + f_D(X, k^0)]$$


T -matrix approximation

Close kinetic equation employing the **T -matrix approximation**

(L. Kadanoff, G. Baym, "Quantum statistical mechanics" 1962, P. Danielewicz, Annals Phys. 152, 239 (1984), W. Botermans and R. Malfliet, Phys. Rept. 198, 115 (1990))

T-matrix approximation

$$i\Pi^<(X, k) = \sum_{\{a,b,c\}} \int \frac{d^4 k_1}{(2\pi)^4} \int \frac{d^4 k_2}{(2\pi)^4} \int \frac{d^4 k_3}{(2\pi)^4} (2\pi)^4 \delta^{(4)}(k_1 + k_2 - k_3 - k) \\ \times |T(k_1^0 + k_2^0 + i\epsilon, \mathbf{k}_1 + \mathbf{k}_2)|^2 iG_{D_a}^<(X, k_1) iG_{\Phi_b}^<(X, k_2) iG_{\Phi_c}^>(X, k_3)$$



Transport equation

$$\left(k^\mu - \frac{1}{2} \frac{\partial \text{Re } \Pi^R(X, k)}{\partial k_\mu} \right) \frac{\partial iG_D^<(X, k)}{\partial X^\mu} = \frac{1}{2} i\Pi^<(X, k) iG_D^>(X, k) - \frac{1}{2} i\Pi^>(X, k) iG_D^<(X, k)$$

Kadanoff-Baym Ansätze:

$$iG_D^<(X, k) = 2\pi S_D(X, k) f_D(X, k^0)$$

$$iG_\Phi^<(X, k) = 2\pi S_\Phi^{(0)}(X, k) f_\Phi^{(0)}(X, k^0)$$

Approximation on light sector [Schenk (1993), Toublan (1997)]:

$$S_\Phi^{(0)}(k^0, \mathbf{k}) = \frac{1}{2E_k} [\delta(k^0 - E_k) - \delta(k^0 + E_k)]$$

and

$$f_\Phi^{(0)}(k^0) = \frac{1}{e^{k^0/T} - 1}$$

using the faster equilibration of light sector

Fokker-Planck equation


Off-shell Fokker-Planck equation, $m_D \gg m_\Phi, T$

$$\frac{\partial}{\partial t} G_D^<(t, k) = \frac{\partial}{\partial k^i} \left\{ \hat{A}(k; T) k^i G_D^<(t, k) + \frac{\partial}{\partial k^j} \left[\hat{B}_0(k; T) \Delta^{ij} + \hat{B}_1(k; T) \frac{k^i k^j}{k^2} \right] G_D^<(t, k) \right\}$$

where $\Delta^{ij} = \delta^{ij} - k^i k^j / k^2$

Off-shell Transport Coefficients

$$\begin{cases} \hat{A}(k^0, \mathbf{k}; T) & \equiv \left\langle 1 - \frac{\mathbf{k} \cdot \mathbf{k}_1}{k^2} \right\rangle \\ \hat{B}_0(k^0, \mathbf{k}; T) & \equiv \frac{1}{4} \left\langle \mathbf{k}_1^2 - \frac{(\mathbf{k} \cdot \mathbf{k}_1)^2}{k^2} \right\rangle \\ \hat{B}_1(k^0, \mathbf{k}; T) & \equiv \frac{1}{2} \left\langle \frac{[\mathbf{k} \cdot (\mathbf{k} - \mathbf{k}_1)]^2}{k^2} \right\rangle \end{cases}$$





with


$$\begin{aligned} \langle \mathcal{F}(\mathbf{k}, \mathbf{k}_1) \rangle &\equiv \frac{1}{2k^0} \sum_{\lambda, \lambda'=\pm} \lambda \lambda' \int dk_1^0 \int \prod_{i=1}^3 \frac{d^3 k_i}{(2\pi)^3} \frac{1}{2E_2 2E_3} S_D(k_1^0, \mathbf{k}_1) (2\pi)^4 \delta^{(3)}(\mathbf{k} + \mathbf{k}_3 - \mathbf{k}_1 - \mathbf{k}_2) \\ &\times \delta(k^0 + \lambda' E_3 - \lambda E_2 - k_1^0) |\mathcal{T}(k^0 + \lambda' E_3, \mathbf{k} + \mathbf{k}_3)|^2 f^{(0)}(\lambda' E_3) \tilde{f}^{(0)}(\lambda E_2) f^{(0)}(k_1^0) \mathcal{F}(\mathbf{k}, \mathbf{k}_1) \end{aligned}$$

“Unitary” contribution: $\lambda' = +$

2 processes evaluated at $|T(k^0 + E_3, \mathbf{k} + \mathbf{k}_3)|^2$:





3 → 1 process
only effective at $T \neq 0$
for off-shell D meson




“Landau” contribution: $\lambda' = -$

2 processes evaluated at $|T(k^0 - E_3, \mathbf{k} + \mathbf{k}_3)|^2$:



Not suppressed at high temperatures!

2 \rightarrow 2 process **only effective** at $T \neq 0$ where Landau cut of $\text{Im } G$ emerges




Spatial diffusion coefficient

Spatial diffusion coefficient

$$2\pi T D_s(T) = \frac{2\pi T^3}{B_0(k^0 = E_k, \mathbf{k} \rightarrow 0; T)}$$

JMT-R, G. Montaña, À. Ramos, L. Tolos,
Phys. Rev. C 105, 025203 (2022)



Lattice-QCD calculations



- ▶ N. Brambilla *et al.*
Phys. Rev. D102, 074503 (2020)
- ▶ D. Banerjee *et al.*
Phys. Rev. D85, 014510 (2012)
- ▶ A. Francis *et al.*
Phys. Rev. D92, 116003 (2015)
- ▶ L. Altenkort *et al.*
Phys. Rev. D103, 014511 (2021)

Bayesian study of RHICs

- ▶ W. Ke *et al.*
Phys. Rev. C98, 064901 (2018)

\bar{B} -meson diffusion coefficient

JMT-R, G. Montaña, À. Ramos, L. Tolos (to be published soon)
G. Montaña (PhD thesis 2022)



Bayesian study of RHICs

- W. Ke *et al.*
Phys. Rev. C98, 064901 (2018)

Quasiparticle model for QGP

- S.K. Das *et al.*
Phys. Rev. D 94, 114039 (2016)

Conclusions and Outlook

Summary

1. D/\bar{B} -meson **EFT** extended to **finite temperature** in a self-consistent way
2. Heavy-meson **kinetic theory** studied via the Kadanoff-Baym equations.
We derived an **off-shell Fokker-Planck** equation
3. Heavy-flavor **transport coefficients** below T_c including thermal modifications and off-shell effects.
Apparent matching to lattice-QCD and Bayesian analyses at $T \sim T_c$

Outlook

1. Analysis of energy loss for *Mesonenstrahlung* processes
2. Thermal evolution of $X(3872)$ as a bound state of $D\bar{D}^*/D^*\bar{D}$
3. Transport coefficients of causal heavy-flavor diffusion

Heavy-meson transport coefficients in a thermal medium



Juan M. Torres-Rincon
Universitat de Barcelona
Institut de Ciències del Cosmos



in collaboration with
G. Montaña, À. Ramos and L. Tolos

DFG Deutsche
Forschungsgemeinschaft



Strong and ElectroWeak Matter 2022
IPhT, Saclay - Université Paris VI, Paris, Jun 20 – 24, 2022