

# **Jet quenching in glasma**


**Stanisław Mrówczyński**

*Jan Kochanowski University, Kielce, Poland  
and National Centre for Nuclear Research, Warsaw, Poland*


In collaboration with **Margaret Carrington & Alina Czajka**

# Motivation

- ▶ We consider the earliest stages of relativistic heavy-ion collisions.
- ▶ According to CGC, color charges confined in the colliding nuclei generate **glasma** – the system of strong mostly classical chromodynamic fields.



- ▶ How hard probes propagate through the glasma?



$$\frac{dE}{dx}, \hat{q} ?$$

# Fokker-Planck Equation

- Transport of hard probes can be described using the Fokker-Planck equation.

$$\left( \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla \right) n(t, \mathbf{r}, \mathbf{p}) = \left( \nabla_p^i X^{ij}(\mathbf{v}) \nabla_p^j + \nabla_p^i Y^i(\mathbf{v}) \right) n(t, \mathbf{r}, \mathbf{p})$$

$n(t, \mathbf{r}, \mathbf{p})$  - distribution function of hard probes

$$\mathbf{v} \equiv \frac{\mathbf{p}}{E_p}, \quad \nabla^i_p \equiv \frac{\partial}{\partial p_i}$$

$$X^{ij}(\mathbf{v}), \quad Y^i(\mathbf{v}) \quad \Rightarrow \quad \left\{ \begin{array}{l} \frac{dE}{dx} = -\frac{\mathbf{v}^i}{\mathbf{v}} Y^i(\mathbf{v}) \quad \text{collisional energy loss} \\ \hat{q} = \frac{2}{\mathbf{v}} \left( \delta^{ij} - \frac{\mathbf{v}^i \mathbf{v}^j}{\mathbf{v}^2} \right) X^{ji}(\mathbf{v}) \quad \text{momentum broadening} \end{array} \right.$$


$$n(t, \mathbf{r}, \mathbf{p}) = n_{\text{eq}}(\mathbf{p}) \sim e^{-\frac{E_{\mathbf{p}}}{T}}$$

$\Leftrightarrow$

solves FK equation

$$Y^j(\mathbf{v}) = \frac{\mathbf{v}^i}{T} X^{ij}(\mathbf{v})$$

# Origin of Fokker-Planck Equation



- ▶ How to obtain a Fokker-Planck equation for plasma?  
Apply the *quasilinear* method known in plasma physics.

# Derivation of Fokker-Planck Equation

The dynamics is assumed to be dominated by strong classical fields.

Vlasov equation

$$p_\mu D^\mu Q(t, \mathbf{r}, \mathbf{p}) - \frac{g}{2} p^\mu \{ F_{\mu\nu}(t, \mathbf{r}), \partial_p^\nu Q(t, \mathbf{r}, \mathbf{p}) \} = 0$$

free streaming

mean-field force

$Q(t, \mathbf{r}, \mathbf{p})$  - exact distribution function of hard probes which is the  $N_c \times N_c$  matrix

$$D^\mu \equiv \partial^\mu - ig[A^\mu, \dots]$$

chromodynamic strength tensor

$$\{A, B\} \equiv AB + BA,$$

$$F^{\mu\nu} \equiv \partial^\mu A^\nu - \partial^\nu A^\mu - ig[A^\mu, A^\nu]$$

# Derivation of Fokker-Planck Equation

Regular and fluctuating quantities

fluctuating part

$$Q(t, \mathbf{r}, \mathbf{p}) = \langle Q(t, \mathbf{r}, \mathbf{p}) \rangle + \delta Q(t, \mathbf{r}, \mathbf{p})$$

regular colorless part

$$\langle Q(t, \mathbf{r}, \mathbf{p}) \rangle = n(t, \mathbf{r}, \mathbf{p}) I$$

$n(t, \mathbf{r}, \mathbf{p})$  - averaged distribution function


- ▶  $|n| \gg |\delta Q|, \quad |\nabla_p n| \gg |\nabla_p \delta Q|$
- ▶  $|\frac{\partial n}{\partial t}| \ll |\frac{\partial \delta Q}{\partial t}|, \quad |\nabla n| \ll |\nabla \delta Q|$
- ▶  $\langle \mathbf{E} \rangle = 0, \quad \langle \mathbf{B} \rangle = 0, \quad \mathbf{E}, \mathbf{B}, A^\mu \sim \delta Q$

# Derivation of Fokker-Planck Equation

$$Q(t, \mathbf{r}, \mathbf{p}) = n(t, \mathbf{r}, \mathbf{p})I + \delta Q(t, \mathbf{r}, \mathbf{p})$$

Lorentz force


Vlasov equation



$$\mathbf{F} \equiv g(\mathbf{E} + \mathbf{v} \times \mathbf{B})$$

$$(D^0 + \mathbf{v} \cdot \mathbf{D})Q - \mathbf{F} \cdot \nabla_p Q = 0$$

ensemble averaging



collision term

$$\left( \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla \right) n(t, \mathbf{r}, \mathbf{p}) = \overbrace{\frac{1}{N_c} \text{Tr} \langle \mathbf{F}(t, \mathbf{r}) \cdot \nabla_p \delta Q(t, \mathbf{r}, \mathbf{p}) \rangle}^{\text{collision term}}$$

Fluctuations provide a collision term.

# Derivation of Fokker-Planck Equation

How to compute the collision term?

$$C \equiv \frac{1}{N_c} \text{Tr} \left\langle \mathbf{F} \cdot \nabla_p \delta Q \right\rangle = ?$$

$$Q(t, \mathbf{r}, \mathbf{p}) = n(t, \mathbf{r}, \mathbf{p}) I + \delta Q(t, \mathbf{r}, \mathbf{p})$$

Vlasov equation

$$(D^0 + \mathbf{v} \cdot \mathbf{D}) Q - \mathbf{F} \cdot \nabla_p Q = 0$$

$$\begin{aligned} n &>> |\delta Q| \\ |\nabla_p n| &>> |\nabla_p \delta Q| \end{aligned}$$

linearization

$$\begin{aligned} \left| \frac{\partial n}{\partial t} \right| &<< \left| \frac{\partial \delta Q}{\partial t} \right| \\ |\nabla n| &<< |\nabla \delta Q| \end{aligned}$$

$$\left( \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla \right) \delta Q = \mathbf{F} \cdot \nabla_p n$$

# Derivation of Fokker-Planck Equation

Solution of the linearized transport equation

$$\left( \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla \right) \delta Q(t, \mathbf{r}, \mathbf{p}) = \mathbf{F}(t, \mathbf{r}) \cdot \nabla_p n(\mathbf{p})$$

initial value

$$\delta Q(t, \mathbf{r}, \mathbf{p}) = \int_0^t dt' \mathbf{F}(t', \mathbf{r} - \mathbf{v}(t-t')) \cdot \nabla_p n(\mathbf{p}) + \delta Q_0(\mathbf{r} - \mathbf{v}t, \mathbf{p})$$



$$C \equiv \frac{1}{N_c} \text{Tr} \langle \mathbf{F}(t, \mathbf{r}) \cdot \nabla_p \delta Q(t, \mathbf{r}, \mathbf{p}) \rangle = (\nabla_p^i X^{ij}(\mathbf{v}) \nabla_p^j + \nabla_p^i Y^i(\mathbf{v})) n(\mathbf{p})$$

$$\left\{ \begin{array}{l} X^{ij}(\mathbf{v}) = \frac{1}{N_c} \int_0^t dt' \langle F^i(t, \mathbf{r}) F^j(t', \mathbf{r} - \mathbf{v}(t-t')) \rangle \\ Y^i(\mathbf{v}) = \frac{1}{N_c} \langle F^i(t, \mathbf{r}) \delta Q_0(\mathbf{r} - \mathbf{v}t, \mathbf{p}) \rangle \frac{1}{n(\mathbf{p})} \end{array} \right.$$

# Fokker-Planck Equation

$$\left[ \left( \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla \right) n(t, \mathbf{r}, \mathbf{p}) = \left( \nabla_p^i X^{ij}(\mathbf{v}) \nabla_p^j + \nabla_p^i Y^i(\mathbf{v}) \right) n(t, \mathbf{r}, \mathbf{p}) \right]$$

$$\left\{ \begin{array}{l} X^{ij}(\mathbf{v}) = \frac{1}{N_c} \int_0^t dt' \left\langle F^i(t, \mathbf{r}) F^j(t', \mathbf{r} - \mathbf{v}(t-t')) \right\rangle \\ Y^j(\mathbf{v}) = \frac{\mathbf{v}^i}{T} X^{ij}(\mathbf{v}) \quad \mathbf{F} \equiv g(\mathbf{E} + \mathbf{v} \times \mathbf{B}) \end{array} \right.$$

► The collision term is given by field correlators  $\langle E^i E^j \rangle, \langle B^i E^j \rangle, \langle B^i B^j \rangle$

► Gauge covariance is lost due to the linearization!

To restore gauge covariance:  $\langle E_a^i(t, \mathbf{r}) E_a^j(t', \mathbf{r}') \rangle \rightarrow \langle E_a^i(t, \mathbf{r}) \Omega_{ab}(t, \mathbf{r} | t', \mathbf{r}') E_b^j(t', \mathbf{r}') \rangle$

$$\Omega(t, \mathbf{r} | t', \mathbf{r}') \equiv P \exp \left[ i g \int_{(t', \mathbf{r}')}^{(t, \mathbf{r})} ds_\mu A^\mu(s) \right]$$

# Field correlators in Equilibrium QGP

space-time translational invariance

$$\boxed{\left\langle E_a^i(t, \mathbf{r}) E_b^j(t', \mathbf{r}') \right\rangle = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \int \frac{d^3 k}{(2\pi)^3} e^{-i(\omega(t-t') - \mathbf{k}(\mathbf{r}-\mathbf{r}'))} \overbrace{\left\langle E_a^i E_b^j \right\rangle}_{\omega, \mathbf{k}}}$$

flucuation spectrum

►  $\left\langle E_a^i E_b^j \right\rangle_{\omega, \mathbf{k}} = 2\delta^{ab} \frac{\omega^4}{e^{\beta|\omega|} - 1} \left[ \frac{k^i k^j}{\mathbf{k}^2} \frac{\text{Im} \varepsilon_L(\omega, \mathbf{k})}{|\omega^2 \varepsilon_L(\omega, \mathbf{k})|^2} + \left( \delta^{ij} - \frac{k^i k^j}{\mathbf{k}^2} \right) \frac{\text{Im} \varepsilon_T(\omega, \mathbf{k})}{|\omega^2 \varepsilon_T(\omega, \mathbf{k}) - \mathbf{k}^2|^2} \right]$

►  $\left\langle B_a^i B_b^j \right\rangle_{\omega, \mathbf{k}} = 2\delta^{ab} \frac{\omega^2 \mathbf{k}^2}{e^{\beta|\omega|} - 1} \left( \delta^{ij} - \frac{k^i k^j}{\mathbf{k}^2} \right) \frac{\text{Im} \varepsilon_T(\omega, \mathbf{k})}{|\omega^2 \varepsilon_T(\omega, \mathbf{k}) - \mathbf{k}^2|^2}$

►  $\left\langle B_a^i E_b^j \right\rangle_{\omega, \mathbf{k}} = \left\langle E_a^j B_b^i \right\rangle_{\omega, \mathbf{k}} = 2\delta^{ab} \frac{\omega^3}{e^{\beta|\omega|} - 1} \varepsilon^{imj} k^m \frac{\text{Im} \varepsilon_T(\omega, \mathbf{k})}{|\omega^2 \varepsilon_T(\omega, \mathbf{k}) - \mathbf{k}^2|^2}$

$\varepsilon_{L,T}(\omega, \mathbf{k})$  - chromodielectric functions

# Fokker-Planck Equation of Equilibrium QGP

$$\left( \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla \right) n(t, \mathbf{r}, \mathbf{p}) = \left( \nabla_p^i X^{ij}(\mathbf{v}) \nabla_p^j + \nabla_p^i Y^i(\mathbf{v}) \right) n(t, \mathbf{r}, \mathbf{p})$$

Isotropy

$$X^{ij}(\mathbf{v}) \equiv X_L(v) \frac{\mathbf{v}^i \mathbf{v}^j}{\mathbf{v}^2} + X_T(v) \left( \delta^{ij} - \frac{\mathbf{v}^i \mathbf{v}^j}{\mathbf{v}^2} \right), \quad Y^j(\mathbf{v}) = \frac{\mathbf{v}^i}{T} X^{ij}(\mathbf{v}) = \frac{\mathbf{v}^i}{T} X_L(v)$$



$$\begin{cases} X_L(v) = \dots \\ X_T(v) = \dots \end{cases}$$

For heavy quarks

$$v \ll 1, \quad g \ll 1$$

$$X_L(v) = X_T(v) \approx \frac{g^2 C_F}{12\pi} m_D^2 T \log\left(\frac{T}{m_D}\right) \quad C_F \equiv \frac{N_c^2 - 1}{2N_c}$$

# Fokker-Planck Equation of Equilibrium QGP



Quantitative agreement with  $X_L(v)$  &  $X_T(v)$  obtained from the Boltzmann collision term by means of the diffusive approximation.


The standard FP equation is reproduced!

# Transport of hard probes in plasma

$$X^{ij}(\mathbf{v}) = \frac{g}{N_c} \int_0^t dt' \left\{ \left\langle E^i(t, \mathbf{r}) E^j(t', \mathbf{r}') \right\rangle + \epsilon^{jkl} v^k \left\langle E^i(t, \mathbf{r}) B^l(t', \mathbf{r}') \right\rangle \right. \\ \left. + \epsilon^{ikl} v^k \left\langle B^l(t, \mathbf{r}) E^j(t', \mathbf{r}') \right\rangle + \epsilon^{ikl} \epsilon^{jmn} v^k v^m \left\langle B^l(t, \mathbf{r}) B^n(t', \mathbf{r}') \right\rangle \right\}$$

$$\mathbf{r}' \equiv \mathbf{r} - \mathbf{v}(t - t')$$

$$\left\{ \begin{array}{l} \hat{q} = \frac{2}{v} \left( \delta^{ij} - \frac{v^i v^j}{v^2} \right) X^{ji}(\mathbf{v}) \\ \frac{dE}{dx} = - \frac{v^i v^j}{vT} X^{ij}(\mathbf{v}) \end{array} \right.$$



- M. Carrington, A. Czajka & St. Mrówczyński, Nuclear Physics A **1001**, 121914 (2020),  
 M. Carrington, A. Czajka & St. Mrówczyński, arXiv: 2112.06812,  
 M. Carrington, A. Czajka & St. Mrówczyński, arXiv:2202.00357, to appear in Physical Review C

# Glasma from AA collisions


The earliest stage of relativistic heavy-ion collisions

Classical Yang-Mills equation

$$D_\mu F^{\mu\nu}(x) = j^\nu(x)$$


$$j^\mu(x) = j_1^\mu(x) + j_2^\mu(x)$$

$$j_{1,2}^\mu(x) = \pm \delta^{\mu\mp} \delta(x^\pm) \rho_{1,2}(\mathbf{x}_\perp)$$



Ansatz of gauge potentials

$$\left\{ \begin{array}{l} A^+(x) = \Theta(x^+) \Theta(x^-) x^+ \alpha(\tau, \mathbf{x}_\perp) \\ A^-(x) = -\Theta(x^+) \Theta(x^-) x^- \alpha(\tau, \mathbf{x}_\perp) \\ A^i(x) = \Theta(x^+) \Theta(x^-) \alpha_\perp^i(\tau, \mathbf{x}_\perp) \\ \quad + \Theta(-x^+) \Theta(x^-) \beta_1^i(\mathbf{x}_\perp) + \Theta(x^+) \Theta(-x^-) \beta_2^i(\mathbf{x}_\perp) \end{array} \right.$$



Boundary condition

$$\left\{ \begin{array}{l} \alpha(0, \mathbf{x}_\perp) = \beta_1^i(\mathbf{x}_\perp) + \beta_2^i(\mathbf{x}_\perp) \\ \alpha_\perp^i(0, \mathbf{x}_\perp) = -\frac{ig}{2} [\beta_1^i(\mathbf{x}_\perp), \beta_2^i(\mathbf{x}_\perp)] \end{array} \right.$$

Gauge condition

$$x^+ A^+ + x^- A^- = 0$$

# Proper time expansion

$$\alpha(\tau, \mathbf{x}_\perp) = \sum_{n=0}^{\infty} \tau^n \alpha_{(n)}(\mathbf{x}_\perp), \quad \alpha_\perp^i(\tau, \mathbf{x}_\perp) = \sum_{n=0}^{\infty} \tau^n \alpha_{\perp(n)}^i(\mathbf{x}_\perp)$$

The Yang-Mills equations for the expanded potentials are solved recursively.

$$E^i = F^{i0}, \quad B^i = \frac{1}{2} \epsilon^{ijk} F^{kj}$$


0th order

$$\mathbf{E}_{(0)} = (0, 0, E^z), \quad \mathbf{B}_{(0)} = (0, 0, B^z)$$


$$E_{(0)}^z(\mathbf{x}_\perp) = -ig[\beta_1^i(\mathbf{x}_\perp), \beta_2^i(\mathbf{x}_\perp)],$$

$$B_{(0)}^z(\mathbf{x}_\perp) = -ig \epsilon^{zij} [\beta_1^i(\mathbf{x}_\perp), \beta_2^j(\mathbf{x}_\perp)],$$

At higher orders transverse fields show up



*E & B fields along the axis z*



# Field correlators

The correlators

$$\left\langle E_a^i(t, \mathbf{r}) E_b^j(t', \mathbf{r}') \right\rangle, \quad \left\langle E_a^i(t, \mathbf{r}) B_b^j(t', \mathbf{r}') \right\rangle, \quad \left\langle B_a^i(t, \mathbf{r}) B_b^j(t', \mathbf{r}') \right\rangle$$

are expressed through

$$\sum \partial^i \partial^j \left\langle \beta_a^k(\mathbf{x}_\perp) \beta_b^l(\mathbf{y}_\perp) \dots \beta_c^m(\mathbf{z}_\perp) \right\rangle$$

IR regulator  $m = \Lambda_{\text{QCD}}$

In covariant gauge  $\partial_\mu \beta^\mu = 0$

$$-\nabla^2 \beta^+(\mathbf{x}_\perp) = \rho(\mathbf{x}_\perp) \Rightarrow \beta^+(\mathbf{x}_\perp) = \frac{1}{2\pi} \int d^2 x'_\perp K_0(m | \mathbf{x}_\perp - \mathbf{x}'_\perp |) \rho(\mathbf{x}'_\perp)$$

The potentials are transformed from the covariant to light-cone gauge

Wick theorem

$$\left\langle \rho_a^k(\mathbf{x}_\perp) \rho_b^l(\mathbf{y}_\perp) \dots \rho_c^m(\mathbf{z}_\perp) \right\rangle = \sum \prod \left\langle \rho_a^i(\mathbf{x}_\perp) \rho_b^j(\mathbf{y}_\perp) \right\rangle$$

Glasma graph approximation

$$\left\langle \beta_a^k(\mathbf{x}_\perp) \beta_b^l(\mathbf{y}_\perp) \dots \beta_c^m(\mathbf{z}_\perp) \right\rangle = \sum \prod \left\langle \beta_a^i(\mathbf{x}_\perp) \beta_b^j(\mathbf{y}_\perp) \right\rangle = \sum \prod B_{ab}^{ij}(\mathbf{x}_\perp - \mathbf{y}_\perp)$$

# Basic correlator

$$B_{ab}^{ij}(\mathbf{x}_\perp, \mathbf{y}_\perp) \equiv \langle \beta_a^i(\mathbf{x}_\perp) \beta_b^j(\mathbf{y}_\perp) \rangle = \int d^2x'_\perp d^2y'_\perp \dots \dots \langle \rho_a^i(\mathbf{x}'_\perp) \rho_b^j(\mathbf{y}'_\perp) \rangle$$

$$\langle \rho_a^i(\mathbf{x}_\perp) \rho_b^j(\mathbf{y}_\perp) \rangle = g^2 \mu \delta^{ab} \delta^{(2)}(\mathbf{x}_\perp - \mathbf{y}_\perp)$$

color charge surface density  
 $\mu = g^{-4} Q_s^2$

$$B_{ab}^{ij}(\mathbf{x}_\perp, \mathbf{y}_\perp) \equiv \delta^{ab} \left( \delta^{ij} C_1(r) - \hat{r}^i \hat{r}^j C_2(r) \right)$$

$$\mathbf{r} \equiv \mathbf{x}_\perp - \mathbf{y}_\perp, \quad r \equiv |\mathbf{r}|, \quad \hat{r}^i \equiv \frac{\mathbf{r}^i}{r}$$

$$\left\{ \begin{array}{l} C_1(r) \equiv \frac{m^2 K_0(mr)}{g^2 N_c (mr K_1(mr) - 1)} \left\{ \exp \left[ \frac{g^4 N_c \mu (mr K_1(mr) - 1)}{4\pi m^2} \right] - 1 \right\} \\ \\ C_2(r) \equiv \frac{m^3 r K_1(mr)}{g^2 N_c (mr K_1(mr) - 1)} \left\{ \exp \left[ \frac{g^4 N_c \mu (mr K_1(mr) - 1)}{4\pi m^2} \right] - 1 \right\} \end{array} \right.$$

$\approx_{r \ll m^{-1}} \# \log(mr)$   
 UV regularization required  
 $r > Q_s^{-1}$

# Fokker-Planck Equation


$$\left( \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla \right) n(t, \mathbf{r}, \mathbf{p}) = \left( \nabla_p^i X^{ij}(\mathbf{v}) \nabla_p^j + \nabla_p^i Y^i(\mathbf{v}) \right) n(t, \mathbf{r}, \mathbf{p})$$

$$\left\{ \begin{array}{l} X^{ij}(\mathbf{v}) = \frac{g}{N_c} \int_0^t dt' \left\{ \left\langle E^i(t, \mathbf{r}) E^j(t', \mathbf{r}') \right\rangle + \epsilon^{jkl} v^k \left\langle E^i(t, \mathbf{r}) B^l(t', \mathbf{r}') \right\rangle \right. \\ \left. + \epsilon^{ikl} v^k \left\langle B^l(t, \mathbf{r}) E^j(t', \mathbf{r}') \right\rangle + \epsilon^{ikl} \epsilon^{jmn} v^k v^m \left\langle B^l(t, \mathbf{r}) B^m(t', \mathbf{r}') \right\rangle \right\} \\ Y^j(\mathbf{v}) = \frac{v^i}{T} X^{ij}(\mathbf{v}) \end{array} \right.$$

The correlators are computed order by order.

$$\left\{ \begin{array}{l} \hat{q} = \frac{2}{v} \left( \delta^{ij} - \frac{v^i v^j}{v^2} \right) X^{ji}(\mathbf{v}) \\ \frac{dE}{dx} = - \frac{v^i}{v} Y^i(\mathbf{v}) \end{array} \right.$$



# Hard probes in plasma



$$N_c = 3, \quad g = 1$$

$$Q_s = 2 \text{ GeV}$$

$$m = 0.2 \text{ GeV}$$



# Glasma impact on jet quenching

Glasma

$$\hat{q}_{\max} = 6 \text{ GeV}^2 / \text{fm}$$

$$t_{\max} = 0.06 \text{ fm}$$

Equilibrium QGP

$$\hat{q} = 3T^3$$


$$t_0 = 0.6 \text{ fm}$$

$$T_0 = 450 \text{ MeV}$$

$$\hat{q}_0 = 1.4 \text{ GeV}^2 / \text{fm}$$

$$T = T_0 \left( \frac{t_0}{t} \right)^{1/3}$$

$$L = 10 \text{ fm}$$



$$\Delta p_T^2 \Big|_{\text{non-eq}} = \int_0^{t_0} dt \hat{q}(t)$$

$$\Delta p_T^2 \Big|_{\text{eq}} = \int_{t_0}^L dt \hat{q}(t)$$

$$\frac{\Delta p_T^2 \Big|_{\text{non-eq}}}{\Delta p_T^2 \Big|_{\text{eq}}} = 0.93$$

# Summary & Conclusions

- ▶ The Fokker-Planck equation of hard probes interacting with classical chromodynamic fields rather than with plasma constituents is derived.
- ▶ The known case of equilibrium plasma is reproduced.
- ▶ The field correlators are computed up to  $\tau^5$ .
- ▶ The momentum broadening and energy loss in the glasma are significantly bigger than in equilibrated QGP.
- ▶ In spite of its short lifetime the glasma provides a significant contribution to the jet quenching.