



Studying the Beginning of the Universe from the Bottom of the World

Clem Pryke – APC Colloquium, Paris – June 27 2022

Modern cosmology in a nutshell:




Edwin Hubble

1) The universe is expanding.
(Hubble, 1920s)

2) It was once hot and dense, like the inside of the Sun.

(Alpher, Gamow, Herman, 1940s)

3) We can see the (redshifted) glow!
The *Cosmic Microwave Background*
(Penzias & Wilson, 1964)




Bob Wilson & Arno Penzias
1978 Nobel Prize

⇒ acceptance of the “HOT BIG BANG”

Cosmic Microwave Background Surface of Last Scattering

All sky temperature map
projected on a sphere




We are at the center


CMB temperature is a sample of the density structure on a shell cut
through the 380,000 year old Universe

Perturbations are one part in 10,000 at that time – and Gaussian!

Power Spectrum (Blob size histogram)




Power Spectrum (Blob size histogram)




Triumphant/Embarrassing Contemporary Cosmology

CMB and other data fits GR based LCDM model *beautifully* – but it demands that 96% of the Universe is invisible to us




And it implies that the future is runaway expansion...



Also it doesn't explain horizon/flatness etc...


The Horizon Problem



How did points A and B “know” to be at the same temperature in the distant past when they had never been in causal contact?

(They still aren’t today!)

Inflation solves the Flatness Problem



→ Inflation... →

If you take some curved space and blow it up enough pretty soon it is no longer curved on a local scale – where “local scale” here means our entire observable Universe!

History of the Universe

Inflation posits a pre-phase of exponential expansion




Alan Guth



Andrei Linde

Radius of the Visible Universe



What Does Inflation Do For Us?

Solves the horizon problem:
Why is the CMB nearly uniform?
How do apparently causally disconnected regions of space get set to the same temperature?



A volume much larger than our entire observable universe today was once a causally connected sub atomic speck.

Solves the flatness problem:
Why is the net spatial curvature so close to zero?



Any initial spatial curvature is diluted away to undetectability by the hyper expansion.

Explains the initial perturbations:
Why Gaussian with close to flat power law spectrum? ($n_s \approx 1$)



Equal amounts of perturbations are injected by quantum fluctuations at each step in the exponential expansion.

Solves the monopole problem:
Why do we not observe magnetic monopoles in the Universe today?



Monopoles are diluted away to undetectability.

Inflation is controversial

Inflationary Paradigm after Planck 2013

Alan H. Guth,¹ David I. Kaiser,¹ and Yasunori Nomura²

¹*Center for Theoretical Physics, Laboratory for Nuclear Science, and Department of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

²*Berkeley Center for Theoretical Physics, Department of Physics and Theoretical Physics Group, Lawrence Berkeley National Laboratory, University of California, Berkeley, CA 94720, USA*

(Dated: December 29, 2013, revised January 13, 2014)

[arxiv/1312.7619](https://arxiv.org/abs/1312.7619)



Inflationary schism after Planck2013

Anna Ijjas,^{1,2} Paul J. Steinhardt,³ and Abraham Loeb⁴

¹*Max-Planck-Institute for Gravitational Physics (Albert-Einstein-Institute), 14476 Potsdam, Germany*

²*Rutgers University, New Brunswick, NJ 08901, USA*


³*Department of Physics and Princeton Center for Theoretical Science, Princeton University, Princeton, NJ 08544, USA*

⁴*Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA*
(Dated: March 14, 2014)


[arxiv/1402.6980](https://arxiv.org/abs/1402.6980)




History of the Universe




History of the Universe




History of the Universe




History of the Universe



History of the Universe




History of the Universe



A "remote sensing" gravitational wave experiment

CMB polarization: arises at last scattering from local radiation quadrupole



CMB Polarization, B-modes and r


- The CMB is partially polarized (due to local radiation quadrupoles at last scattering)
- Any polarization pattern can be decomposed into E-modes (gradient modes) and B-modes (curl modes)
- Basic LCDM makes only E-modes at last scattering – although lensing deflections in flight produce a bit of a B-mode
- Primordial gravitational waves produce both E-modes and B-modes – but best to look for the B-modes since most distinct there
- Theory gives us a good template shape for the gravitational wave signal – but it does *not* tell us the amplitude
- The amplitude is parameterized by a single number r
- A wide range of inflation theories exist – the simplest are already ruled out – more complex ones can produce r which is undetectably small
- The experimental mission is to obtain the best possible sensitivity to r
- If we can detect r we determine the energy scale of inflation – if not we can rule out additional inflationary models

CMB Polarization, B-modes and r


- The CMB is partially polarized (due to local radiation quadrupoles at last scattering)
- Any polarization pattern can be decomposed into E-modes (gradient modes) and B-modes (curl modes)
- Basic LCDM makes only E-modes at last scattering – although lensing deflections in flight produce a bit of a B-mode
- Primordial gravitational waves produce both E-modes and B-modes – but best to look for the B-modes since most distinct there
- Theory gives us a good template shape for the gravitational wave signal – but it does *not* tell us the amplitude
- The amplitude is parameterized by a single number r
- A wide range of inflation theories ruled out – more complex ones small
- The experimental mission is to
- If we can detect r we determine we can rule out additional inflation

Warning: It's a bit like the search for proton decay – a well motivated physics target to look for, but theories can be adjusted to make the amplitude arbitrarily small...



CMB Polarization power spectra




CMB space missions map the full sky



Ground based telescopes map part of the sky more deeply





INIST

CARDIFF
UNIVERSITY

UNIVERSITY OF
Cincinnati




SLAC
JPL

BICEP/Keck Basic Experimental Strategy




- Small aperture telescopes (cheap, fast, low systematics)
- Target the 2 degree peak of the PGW B-mode
- Integrate continuously from South Pole
- Observe order 1% patch of sky (smaller is actually better!)
- Scan and pair difference modulation

Unfortunately we are in a galaxy




The interstellar space within our galaxy contains cold dust grains which glow thermally in microwaves, and relativistic electrons which emit synchrotron radiation

Unfortunately we are in a galaxy




When CMB people talk about “foregrounds” it is analogous to what HEP people call “backgrounds” – something which gets in the way of the thing one is trying to measure.

Polarized Foreground Contamination from Our Galaxy



(Plot from Dunkley et al
arxiv/0811.3915)

Polarized Foreground Contamination from Our Galaxy




The BICEP/Keck Telescopes

Telescope as compact as possible while still having the angular resolution to observe degree-scale features.



On-axis, refractive optics allow the entire telescope to rotate around boresight for polarization modulation.

Pulse tube cooler cools the optical elements to 4 K.

3-stage helium sorption refrigerator further cools the detectors to 0.3 K.



Mass-produced Superconducting Detectors




Planar antenna array



Transition edge sensor

Slot antennas



Microstrip filters


BICEP/Keck Band Passes

The dry South Pole atmosphere provides excellent observing conditions most of the year.


The approx. 30% fractional bandpasses fit within atmospheric transmission windows straddled by oxygen and water lines.

In these windows, the atmosphere is quite transparent to microwaves.

The detector passbands are defined by a filter printed directly onto the focal plane wafers.



Journey to the South Pole



Minneapolis -> California -> New Zealand -> McMurdo -> South Pole

Antarctic Continent



Larger than the US – Ice sheet 3000 metres thick!



Big Program!



Arrival in Antarctica



McMurdo – base on the coast



On to the Pole – over the Transantarctic Mountains




Unloading at Pole



The Actual South Pole



Nothing Out There!



Why do this at the Pole?

South Pole CMB telescopes



- High and *dry* – excellent atmospheric transmission
- On Earth's rotational axis - One day/night cycle per year
 - Long night makes for great quality data
- Good support infrastructure – power, cargo, data comm
- Food and accommodation provided
- Even Tuesday night bingo...

Stage 2

Stage 3



BICEP2
(2010-2012)

Keck Array
(2012-2019)




BICEP3
(2016-present)

BICEP Array
(2020-present)





Telescope and Mount




Focal Plane



Beams on Sky






Raw Data - Perfect Weather

Time 50 mins



Run 20120622C01_dk293, scan block 1; Rx: 0



➤ Cover the whole field in 60 such scansets then start over at new boresight rotation


➤ Scanning modulates the CMB signal to freqs < 4 Hz

Raw Data - Worse Weather

Time 50 mins





Run 20120622D01_dk293, scan block 1; Rx: 0




➤ Scanning over lumpy atmosphere
→ “clouds”

➤ Pair difference still clean
→ atmosphere is unpolarized





BK18 95GHz Maps


95 GHz T signal





95 GHz T noise




95 GHz Q signal





95 GHz Q noise



95 GHz U signal




95 GHz U noise




Right ascension [deg.]

Declination [deg.]

BK18 150GHz Maps



BK18 220GHz Maps




BK18 auto/cross spectra between:
 BICEP3 95GHz,
 BICEP2/Keck 150GHz,
 Keck 220GHz,
 and Planck 353GHz

Black lines are
 LCDM
 Red lines are
 LCDM+dust

Blue panels are
 BB
 spectra


Green panels are
 EE spectra




Multicomponent parametric likelihood analysis

Take the joint likelihood of all the spectra simultaneously vs. model for BB that is the Λ CDM lensing expectation + 7 parameter foreground model + r

foreground model = dust + synchrotron






$r_{.05} < 0.09$

BKP


arxiv/1502.00612



$r_{.05} < 0.07$

BK14


arxiv/1510.09217



$r_{.05} < 0.06$

BK15

arxiv/1810.05216




$r_{.05} < 0.035$

BK18


arxiv/2110.00483

Per bandpower CMB component extraction




Stage 2

Stage 3




Stage 2

Stage 3





Stage 2


Stage 3




Delensing with SPT-3G data



High resolution maps




Can be used to reconstruct the lensing deflection map...





...which can then be used to calculate the lensing signal enabling a deeper search for inflationary gravitational waves

Latest Generation Experiment "BICEP Array"



2018-19: Built New Telescope at UMN












Lifting on part of new telescope



Working in the snow





Feb 2020 – the finished product



Summary

- The Universe is expanding – it was once a hot dense “fireball”.
- We understand its development all the way back to a very high energy state.
- The theory of “Inflation” says that our entire observable Universe today all came from a single sub-atomic spec in a hyper expansion lasting a tiny fraction of a second
 - If this “Inflation” really happened it will have made a background of gravitational waves
 - We may be able to detect the imprint of these by measuring the polarization pattern of the Cosmic Microwave Background – if we can build a sensitive enough telescope
 - BICEP/Keck set the world’s best upper limits to date ruling out multiple previously popular classes of inflationary models (monomial and natural)
 - And the search goes on with bigger and better experiments...



Stage IV CMB experiment: CMB-S4

- CMB-S4: a next generation ground-based program building on CMB stage 2 & 3 projects to pursue inflation, neutrino properties, dark energy and new discoveries.
- Targeting to deploy O(500,000) detectors spanning 30 - 300 GHz using multiple telescopes and sites to map most of the sky to provide sensitivity to cross critical science thresholds.
- Multi-agency effort (DOE & NSF). Complementary with balloon and space-based instruments.
- Broad participation of the US CMB community, including the existing NSF CMB groups, DOE National Labs and the High Energy Physics community.
- U.S. led program; international partnerships expected.



Recommended by P5 & NRC Antarctic reports



A science driven program combining the deep CMB experience of the university groups with the expertise and resources at the national labs.