


# Primordial black holes as dark matter

**Alexander Kusenko**  
(UCLA and Kavli IPMU)  
Paris, LPTHE 2022

# Nobel Prize 2020: Black holes' existence confirmed

Milky Way, Sagittarius A\*



R. Penrose  
R. Genzel  
A. Ghez




A. Ghez (UCLA)

Observations: BHs exist!

⇒ **PBH is a plausible dark matter candidate, the only candidate known to exist in nature**

# Experimental constraints






## A - Dark matter

B - candidate events from HSC, OGLE  
[1701.02151, 1901.07120]

C - interesting for GW, as well as  
transmuted NS  $\rightarrow$  BH population  
[1707.05849; 2008.12780]

D - seeds of supermassive black holes  
[astro-ph/0204486,  
arXiv:1202.3848, 2008.11184]


# First candidate events [Takada et al., Kavli IPMU]



**Figure 13.** One remaining candidate that passed all the selection criteria of microlensing event. The images in the upper plot show the postage-stamped images around the candidate as in Fig. 7: the reference image, the target image, the difference image and the residual image after subtracting the best-fit PSF image, respectively. The lower panel shows that the best-fit microlensing model gives a fairly good fitting to the measured light curve.

## First candidate events from HSC and OGLE

[Niikura et al.. Nature Astron., arXiv:1701.02151, 1901.07120]




# How to make PBHs

Need a ~30% or higher overdensity early enough in the history of the universe.

- Primordial fluctuations enhanced on small scales (inflation model)
- Yukawa interactions, “long-range” forces, radiative cooling => PBH
- Supersymmetry: Q-balls as building blocks of PBH
- Supersymmetry: Q-balls with long-range scalar forces
- Multiverse => PBHs

## THE BEST Simple RECIPES

America's Test Kitchen



200 Flavorful Recipes that Cook in 30 Minutes or Less


# How to make PBHs

Need a ~30% or higher overdensity early enough in the history of the universe.

- **Primordial fluctuations enhanced on small scales (inflation model)**
- Yukawa interactions, “long-range” forces, radiative cooling => PBH
- Supersymmetry: Q-balls as building blocks of PBH
- Supersymmetry: Q-balls with long-range scalar forces
- Multiverse => PBHs

# THE BEST Simple RECIPES

America's Test Kitchen



200 Flavorful Recipes that Cook in 30 Minutes or Less


# How to make PBHs

Need a ~30% or higher overdensity early enough in the history of the universe.

- Primordial fluctuations enhanced on small scales (inflation model)
- **Yukawa interactions, “long-range” forces, radiative cooling => PBH**
- Supersymmetry: Q-balls as building blocks of PBH
- Supersymmetry: Q-balls with long-range scalar forces
- Multiverse => PBHs

## THE BEST Simple RECIPES

America's Test Kitchen



200 Flavorful Recipes that Cook in **30 Minutes or Less**

# PBH formation mechanism: Yukawa “fifth force”

**Yukawa interactions:**

$$V(r) = \frac{y^2}{r} e^{-m_\chi r}$$

$$y\chi\bar{\psi}\psi$$

**a heavy fermion interacting  
with a light scalar**

A light scalar field  $\Rightarrow$  long-range attractive force,  $\Rightarrow$  instability similar to  
stronger than gravity gravitational instability,  
only stronger

$\Rightarrow$  **halos form** even in radiation dominated universe

[Amendola et al., 1711.09915; Savastano et al., 1906.05300; Domenech, Sasaki, 2104.05271]

Same Yukawa coupling provides a source of **radiative cooling** by emission of  
gravitational radiation  $\Rightarrow$  **halos collapse to black holes**

[Flores, AK, 2008.12456, PRL 126 (2021) 041101; 2008.12456]



# Strong long-range force: instability and structure formation

$$\delta(x, t) = \delta\rho/\rho$$

energy density perturbations (radiation)

$$\Delta(x, t) = \Delta n_\psi/n_\psi$$

density perturbations of a kinetically decoupled particle

$$\ddot{\delta}_k + \frac{1}{t}\dot{\delta}_k - \frac{3}{8t^2}(\Omega_r\delta_k + \Omega_m\Delta_k) = 0$$

$$\ddot{\Delta}_k + \frac{1}{t}\dot{\Delta}_k - \frac{3}{8t^2}[\Omega_r\delta_k + \Omega_m(1 + \beta^2)\Delta_k] = 0$$

$\Rightarrow$

$$\Delta_k(a) \approx \Delta_{k,\text{in}} \left( \frac{t}{t_{\text{in}}} \right)^{p/2}, \quad p = \sqrt{\frac{3}{2}(1 + \beta^2)\Omega_\psi}$$

$$\beta \equiv y(M_P/m_\psi) \gg 1$$

$p = \text{huge} \Rightarrow$



[Flores, AK, 2008.12456, PRL]

**fast growth, even in the radiation-dominated era!**

# Growth of structures due to Yukawa force: N-body simulations

Inman, PRELIMINARY


Domenech, Inman, Sasaki, AK  
work in progress




# Rapid growth of structures... plus radiative cooling!

Same Yukawa fields allow particles moving with acceleration emit scalar waves

⇒ radiative cooling and collapse to black holes



Flores, AK, Phys.Rev.Lett. 126 (2021) 4, 041101;  
2008.12456



# PBH DM abundance natural for $m_\psi \sim 1-100$ GeV

Asymmetric dark matter models: Asymmetry in the dark sector = baryon asymmetry

In our case, all these particles end up in black holes:


$$f_{\text{PBH}} = \frac{\Omega_{\text{PBH}}}{\Omega_{\text{DM}}} = 0.2 \frac{m_\psi}{m_p} \frac{\eta_\psi}{\eta_B} = \left( \frac{m_\psi}{5 \text{ GeV}} \right) \left( \frac{\eta_\psi}{10^{-10}} \right)$$

Similar to asymmetric dark matter [Petraki]

[Flores, AK, 2008.12456, PRL 126 (2021) 041101]

Natural explanation for the ratio

**(dark matter density) / (ordinary matter density)**  
**for  $\sim 1-100$  GeV masses**




# How to make PBHs

Need a ~30% or higher overdensity early enough in the history of the universe.

- Primordial fluctuations enhanced on small scales (modify inflation)
- Yukawa interactions, “long-range” forces, radiative cooling => PBH
- **Supersymmetry: Q-balls as building blocks of PBH**
- Supersymmetry: Q-balls with long-range scalar forces
- Multiverse => PBHs

## THE BEST Simple RECIPES

America's Test Kitchen




200 Flavorful Recipes that Cook in **30 Minutes or Less**

# Scalar fields in de Sitter space (used by Affleck-Dine)

A scalar with a small mass develops a VEV


[Chernikov, Tagirov; Starobinsky, Zeldovich; Bunch, Davies; Linde; Affleck, Dine; Starobinsky, Yokoyama]



# Scalar fields in de Sitter space (used by Affleck-Dine)

A scalar with a small mass develops a VEV

[Chernikov, Tagirov; Starobinsky, Zeldovich; Bunch, Davies; Linde; Affleck, Dine; Starobinsky, Yokoyama]





# Scalar fields: an instability (Q-balls)

**Gravitational instability** can occur due to the attractive force of gravity.

**Similar instability** can occur due to scalar self-interaction which is **attractive**:

$$U(\phi) \supset \lambda_3 \phi^3 \quad \text{or} \quad \lambda_{\chi\phi\phi} \chi \phi^\dagger \phi$$



[AK, Shaposhnikov, hep-ph/9709492]




# Scalar fields: an instability (Q-balls)

homogeneous solution  $\varphi(x, t) = \varphi(t) \equiv R(t)e^{i\Omega(t)}$

$$\delta R, \delta \Omega \propto e^{S(t) - i\vec{k}\vec{x}}$$

$$\delta\ddot{\Omega} + 3H(\delta\dot{\Omega}) - \frac{1}{a^2(t)}\Delta(\delta\Omega) + \frac{2\dot{R}}{R}(\delta\dot{\Omega}) + \frac{2\dot{\Omega}}{R}(\delta\dot{R}) - \frac{2\dot{R}\dot{\Omega}}{R^2}\delta R = 0,$$

$$\delta\ddot{R} + 3H(\delta\dot{R}) - \frac{1}{a^2(t)}\Delta(\delta R) - 2R\dot{\Omega}(\delta\dot{\Omega}) + U''\delta R - \dot{\Omega}^2\delta R = 0.$$




$$(\dot{\Omega}^2 - U''(R)) > 0 \Rightarrow \text{growing modes: } 0 < k < k_{\max}$$

$$k_{\max}(t) = a(t)\sqrt{\dot{\Omega}^2 - U''(R)}$$


Also of interest: oscillons

AK, Shaposhnikov, hep-ph/9709492


# Numerical simulations of scalar field fragmentation



[Multamaki].




SUSY Q-balls



[Kasuya, Kawasaki]

# Affleck - Dine baryogenesis (SUSY): scalars are flat directions



Inflation

radiation dominated

matter dominated

modern era  
(dark energy  
dominated)

origin of  
primordial  
perturbations

$$p = \frac{1}{3} \rho$$


$$\rho \propto a^{-4}$$

structures don't grow


$$p = 0$$

$$\rho \propto a^{-3}$$

structures grow




# Scalar lump (Q-ball) formation can lead to PBHs



Early matter dominated epoch in the middle of radiation dominated era

[Cotner, AK, Phys.Rev.Lett. 119 (2017) 031103]

# Size of “particles” affects Poisson fluctuations



many particles  $\Rightarrow$   
small (poisson) fluctuations



FEW GIANT PARTICLES  $\Rightarrow$   
LARGE POISSON FLUCTUATIONS

# Affleck-Dine process and scalar fragmentation in SUSY

[Cotner, AK, Sasaki, Takhistov et al., 1612.02529, 1706.09003, 1801.03321, 1907.10613]


Flat directions lifted by SUSY breaking terms, which determine the scale of fragmentation.

$$M_{\text{hor}} \sim r_f^{-1} \left( \frac{M_{\text{Planck}}^3}{M_{\text{SUSY}}^2} \right) \sim 10^{23} \text{g} \left( \frac{100 \text{ TeV}}{M_{\text{SUSY}}} \right)^2$$

$$M_{\text{PBH}} \sim r_f^{-1} \times 10^{22} \text{g} \left( \frac{100 \text{ TeV}}{M_{\text{SUSY}}} \right)^2$$

$$10^{17} \text{g} \lesssim M_{\text{PBH}} \lesssim 10^{22} \text{g}$$

Cotner, AK, Phys.Rev.Lett. 119 (2017) 031103  
 Cotner, AK, Sasaki, Takhistov, JCAP 1910 (2019) 077




# How to make PBHs

Need a ~30% or higher overdensity early enough in the history of the universe.

- Primordial fluctuations enhanced on small scales (modify inflation)
- Yukawa interactions, “long-range” forces, radiative cooling => PBH
- Supersymmetry: Q-balls as building blocks of PBH
- **Supersymmetry: Q-balls with long-range scalar forces**
- Multiverse => PBHs

## THE BEST Simple America's Test Kitchen RECIPES



200 Flavorful Recipes that Cook in 30 Minutes or Less

# Yet another way to get PBHs from SUSY: long-range forces

A SUSY flat direction  $\varphi$  can couple to another SUSY scalar,  $\chi$ , which can mediate long-range forces between SUSY Q-balls, leading to Yukawa long-range potential

$$V(\varphi, \chi) = U(\varphi) + \frac{1}{2}m_\chi^2\chi^2 - y\chi\varphi^\dagger\varphi + \frac{\lambda}{4}\chi^4$$


Long-range forces  
work as in the case of  
Yukawa interaction  
but  
**individual Q-balls**  
grow until they reach  
the mass/size of a BH

$$f_{\text{PBH}} = \frac{\Omega_{\text{DM}}}{\Omega_{\text{DM}}}$$

$$\simeq \left( \frac{e^{-1/2\epsilon}}{2 \times 10^{-13}} \right) \left( \frac{\Lambda}{10^5 \text{ GeV}} \right)^2 \left( \frac{10^6 \text{ GeV}}{T_f} \right)$$

$$M_{\text{PBH}} \simeq 10^{23} \text{ g} \left( \frac{100 \text{ TeV}}{\Lambda} \right)^2$$

Flores, AK, 2108.08416






# How to make PBHs

Need a ~30% or higher overdensity early enough in the history of the universe.

- Primordial fluctuations enhanced on small scales (modify inflation)  
Yukawa interactions, “long-range” forces, radiative cooling => PBH
  - Supersymmetry: Q-balls as building blocks of PBH
  - Supersymmetry: Q-balls with long-range scalar forces
- **Multiverse => PBHs**



# THE BEST Simple RECIPES

America's Test Kitchen



200 Flavorful Recipes that Cook in **30 Minutes or Less**

# And yet another mechanism: inflationary multiverse




Tunneling events lead to nucleation of baby universes, which appear to outside observer as black holes.

Deng, Vilenkin JCAP 12 (2017) 044

AK, Sasaki, Sugiyama, Takada, Takhistov, Vitagliano, Phys Rev Lett 125 (2020) 181304

# Tail of the mass the function $\propto M^{-1/2}$ , accessible to HSC



[AK, Sasaki, Sugiyama, Takada, Takhistov, Vitagliano, Phys.Rev.Lett. 125 (2020) 181304  
arXiv:2001.09160]

# PBH masses, spins, and a *new window on the early universe*

Formation mechanism	Mass range	PBH spin
Inflationary perturbations [review: 2007.10722]	DM, LIGO, supermassive	small
Yukawa “fifth force” [2008.12456]	DM, LIGO, supermassive	small
Long-range forces between SUSY Q-balls [2108.08416]	DM (mass range: $10^{-16}$ - $10^{-6} M_{\odot}$ )	small
Supersymmetry flat directions, Q-balls [1612.02529, 1706.09003, 1907.10613]	DM (mass range: $10^{-16}$ - $10^{-6} M_{\odot}$ )	large
Light scalar field Q-balls (not SUSY) [1612.02529, 1706.09003, 1907.10613]	DM, LIGO, supermassive	large
Oscillons [1801.03321]	DM, LIGO, supermassive	large
Multiverse bubbles [1512.01819, 1710.02865, 2001.09160]	DM, LIGO, supermassive	small


# PBH and neutron stars

- Neutron stars can capture PBH, which consume and destroy them from the inside.
- Capture probability high enough in DM rich environments, e.g. Galactic Center
- Missing pulsar problem...  
[e.g. Dexter, O'Leary, arXiv:1310.7022]
- What happens if NSs really are systematically destroyed by PBH?

## Neutron star destruction by black holes


⇒ r-process nucleosynthesis, 511 keV, FRB

[Fuller, AK, Takhistov, Phys.Rev.Lett. 119 (2017) 061101 ]




Fast-spinning millisecond pulsar.

Image: NASA/Dana Berry




# MSP spun up by an accreting PBH



- MSP with a BH inside, spinning near mass shedding limit: elongated spheroid
- Rigid rotator: viscosity sufficient even without magnetic fields [Kouvaris, Tinyakov]; more so if magnetic field flux tubes are considered
- Accretion leads to a decrease in the radius, increase in the angular velocity (by angular momentum conservation)
- Equatorial regions gain speed in excess of escape velocity: ejection of cold neutron matter


[Fuller, AK, Takhistov, Phys. Rev. Lett. 119 (2017) 061101] also, *Viewpoint* by H.-T. Janka

# r-process nucleosynthesis: site unknown



- s-process cannot produce peaks of heavy elements
- Observations well described by r-process
- Neutron rich environment needed
- Site? SNe? NS-NS collisions?..

# r-process nucleosynthesis: site unknown



- **SN?** Problematic: neutrinos
- **NS mergers?** Can account for all r-process?




Image: Los Alamos, Nuclear Data Group



# NS-NS might not be not enough...

SCIENTISTS DAZED AND CONFUSED BY EXTRAORDINARY AMOUNT OF GOLD IN THE UNIVERSE


There's too much gold in the universe. No one knows where it came from.

By Rafi Letzter - Staff Writer 12 days ago

Something is showering gold across the universe. But no one knows what it is.

Kobayashi,

THE ASTROPHYSICAL JOURNAL, 900:179 (33pp), 2020 September 10



**Figure 39.** The time evolution (in Gyr) of the origin of elements in the periodic table: Big Bang nucleosynthesis (black), AGB stars (green), core-collapse supernovae including SNe II, HNe, ECSNe, and MRSNe (blue), SNe Ia (red), and NSMs (magenta). The amounts returned via stellar mass loss are also included for AGB stars and core-collapse supernovae depending on the progenitor mass. The dotted lines indicate the observed solar values.

[Kobayashi et al.,  
ApJ 900:179, 2020]

# r-process material: observations

Milky Way (total):  $M \sim 10^4 M_{\odot}$


Ultra Faint Dwarfs (UFD): most of UFDs show no enhancement of r-process abundance.

However, **Reticulum II** shows an enhancement by factor  $10^2$ - $10^3$ !

*"Rare event"* consistent with the UFD data: one in ten shows r-process material  
[Ji, Frebel et al. Nature, 2016]

# NS disruptions by PBHs


- Centrifugal ejection of cold neutron-rich material ( $\sim 0.1 M_{\odot}$ )  
MW:  $M \sim 10^4 M_{\odot}$  ✓
- UFD: a rare event, only one in ten UFDs could host it in 10 Gyr ✓
- Globular clusters: low/average DM density, but high density of millisecond pulsars. Rates OK. ✓



[Fuller, AK, Takhistov, PRL 119 (2017) 061101]  
also, a *Viewpoint* PRL article by Hans-Thomas Janka

# NS disruptions by PBHs

- Weak/different GW signal
- No significant neutrino emission
- Fast Radio Bursts
- Kilonova event **without** a GW counterpart, but with a possible coincident FRB (LSST, ZTF,...)
- 511 keV line



[Fuller, AK, Takhistov, Phys. Rev. Lett. 119 (2017) 061101 ]


# 511-keV line in Galactic Center

Origin of positrons unknown. Need to produce  $10^{50}$  positrons per year. Positrons must be produced with energies below 3 MeV to annihilate at rest. [Beacom, Yuksel '08]

Cold, neutron-rich material ejected in PBH-NS events is heated by  $\beta$ -decay and fission to  $T \sim 0.1$  MeV

→ **generate  $10^{50}$   $e^+$ /yr** for the rates needed to explain r-process nucleosynthesis.

Positrons are non-relativistic.



ESA/Bouchet et al.

$$\Gamma(e^+e^- \rightarrow \gamma\gamma) \sim 10^{50} \text{yr}^{-1}$$

Fuller, AK, Takhistov, Phys. Rev. Lett. 119 (2017) 061101

# Fast Radio Bursts (FRB)

Origin unknown. One repeater, others: non-repeaters.  $\tau \sim$  ms.

PBH - NS events: final stages dynamical time scale  $\tau \sim$  ms.

NS magnetic field energy available for release:  $\sim 10^{41}$  erg


Massive rearrangement of magnetic fields at the end of the NS life, on the time scale  $\sim$ ms produces an FRB.

**Consistent with observed FRB fluence.**

Fuller, AK, Takhistov, Phys.Rev.Lett. 119 (2017) 6, 061101; 1704.01129

Abramowicz, Bejger, Wielgus, Astrophys. J. 868, 17 (2018); 1704.05931

Kainulainen, Nurmi, Schiappacasse, Yanagida, arXiv:2108.08717



# GW detectors can discover small PBH...


**PBH + NS**



**BH of 1-2  $M_{\odot}$**

[Takhistov et al., 1704.01129, 1707.05849;  
2008.12780]

...if it detects mergers of  
**1-2  $M_{\odot}$  black holes**  
(not expected from evolution of stars)



# Conclusion

- Simple, generic formation scenarios in the early universe:  
PBH from scalar forces, PBH from a scalar field fragmentation, PBH from vacuum bubbles...
- PBH with masses  $10^{-16} - 10^{-10} M_{\odot}$ , motivated by 1-100 TeV scale **supersymmetry**, can make up 100% (or less) of dark matter. **PBH is a generic dark matter candidate in SUSY**
- PBH from  $\sim 1$ -100 GeV scale particles can naturally explain DM abundance
- Microlensing (HSC) can detect the tail of DM mass function.
- PBH can contribute to r-process nucleosynthesis
- Signatures of PBH:
  - Kilonova without a GW counterpart, or with a weak/unusual GW signature
  - An unexpected population of 1-2  $M_{\odot}$  black holes (GW)
  - Galactic positrons, FRB, etc.