



ID de Contribution: 176

Type: Non spécifié

J/ψ photoproduction in Pb–Pb collisions with nuclear overlap measured in ALICE at LHC.

mardi 25 octobre 2022 15:00 (30 minutes)

The presence of an unexpected significant excess of low p_T J/ψ over the expected hadronic J/ψ production was confirmed in peripheral Pb–Pb collisions and observed for the first time in semi-central Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with ALICE at LHC. The measurements were performed in the dimuon decay channel at forward rapidity ($2.5 < y < 4$) and in the dielectron decay channel at mid-rapidity. Surprisingly, a large increase in the J/ψ nuclear modification factor was observed in peripheral Pb–Pb collisions at low p_T , below 0.3 GeV/c. Most of the excess is believed to originate from a coherent photoproduction mechanism, which was unexpected until few years ago in peripheral collisions with nuclear overlap and is well known in ultra-peripheral Pb–Pb collisions. The coherent photoproduction implies the interaction of a quasi-real photon coherently with the full Pb target nucleus, which remains intact. Many theoretical models developed for ultra-peripheral collisions have tried to extend the description of the coherent J/ψ photoproduction mechanism to Pb–Pb collisions with nuclear overlap. In this presentation, I will show the results obtained with ALICE, and will compare them to theoretical calculations. And I will introduce the most recent analysis of the rapidity differential coherent J/ψ photoproduction cross section measurement at forward rapidity in ALICE.

Auteur principal: SHATAT, Afnan (IJCLab, Université Paris Saclay)

Orateur: SHATAT, Afnan (IJCLab, Université Paris Saclay)

Classification de Session: Hadronic Physics

Classification de thématique: Hadronic Physics