


Cosmic Neutrinos in Louvain-la-Neuve

francis halzen



- ANTARES, Baikal GVD, IceCube
- the diffuse high-energy neutrino flux
- observation of the first sources
- multimessenger astronomy: plan B


ν and γ beams : heaven and earth




accelerator is powered by large gravitational energy

supermassive black hole

nearby radiation or hydrogen, or...




\sim cosmic ray + neutrino




\sim cosmic ray + gamma

- muon produced by neutrino near IceCube
- comes through the Earth
- 2,600 TeV inside detector
- not atmospheric




- muon produced by neutrino near IceCube
- comes through the Earth
- 2,600 TeV inside detector
- not atmospheric
- angular resolution: astronomy




neutrinos interacting
inside the detector

muon neutrinos
filtered by the Earth



date: **June 11, 2014**
most probable energy: **9 PeV**
topology: **track**




superior total energy
measurement
to 10%, all flavors, all sky

astronomy: superior
angular resolution
superior (0.3°)

muon neutrino flux filtered by the Earth: atmospheric vs cosmic


Number of Events per Bin

Data/MC




electron and tau neutrinos (showers)

$$E^2 dN/dE \sim E^{-2.5}$$




muon neutrinos through Earth (tracks)

update: multi-year cascade ($\nu e + \nu \tau$) analysis



update: multi-year starting ν_μ track analysis



All-sky search for HE cascades

[arXiv:2211.09447](https://arxiv.org/abs/2211.09447)

Additional selection requirements:

($N_{\text{hit}_\mu} = 0, E_{\text{rec}} \geq 70 \text{ TeV}$) or

($N_{\text{hit}_\mu} = 1, E_{\text{rec}} \geq 100 \text{ TeV}$)

N_{hit_μ} is number of hits in time interval
where hits from muons are expected

Expected:

7.4 events from atm. muons

0.8 events from atm. neutrinos



5.8 events for Baikal-GVD best fit

$E^{-2.58}$ astrophysical flux

Found in real data: 16 events

Probability for the background-only hypothesis (stat.+sys.)

P-value = 0.026 (2.22 σ)



Search for upward moving events [arXiv:2211.09447](https://arxiv.org/abs/2211.09447)

Additional selection requirements:

$$E > 15 \text{ TeV} \text{ & } N_{\text{hit}} > 11 \text{ & } \cos\theta < -0.25$$

Expected:

0.5 events from atm. muons


2.7 events from atm. neutrinos

6.3 events for Baikal-GVD best fit $E^{-2.58}$
astrophysical flux

Found in data: 11 events

Probability for the background-only
hypothesis (stat.+sys.)

P-value = 0.0024 (3.05 σ)





ANTARES

Updated data sample @ ICRC2019: 2007-2015 (2450 days) → [2007-2018 \(3330 days\)](#)

All-sky / All-flavor neutrino search

- Selection cuts optimized with MRF procedure (assumed spectral index $\Gamma = 2.5$)
- Look for excess above a given E_{th}
- Combine track & shower samples



Data: 50 events (27 tracks + 23 showers)



Background expectation (atm. flux, incl. prompt) :

36.1 ± 8.7 (19.9 tracks and 16.2 showers) – stat. + syst.


Results not really constraining... but fully compatible with IceCube

Combined (tracks+showers) likelihood fitting:

Cosmic: $\Phi_{100 \text{ TeV}} = (1.5 \pm 1.0) \times 10^{-18} \text{ GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$
 $\Gamma = 2.3 \pm 0.4$





Results not really constraining... but fully compatible with IceCube




cosmic neutrinos: four independent observations exceeding 5σ

- muon neutrinos through the Earth
- starting neutrinos: all flavors
- tau neutrinos
- Glashow event



partially contained event with energy 6.3 PeV




resonant production of a weak intermediate boson by an anti-electron neutrino interacting with an atomic electron




- energy measurement understood
- shower consistent with the hadronic decay of a weak intermediate boson W
- identification of anti-electron neutrino





in the extreme universe the energy in neutrinos is larger than the energy in gamma rays




the neutrino sources are likely opaque to gamma rays

- we do not see our own Galaxy, we see the Universe
- Our Galaxy is a neutrino desert
- in the extreme universe more energy is emitted in neutrinos than in gamma rays
- neutrinos are produced in obscured sources
- this was totally unexpected and represents a great opportunity for a “new” astronomy


ANTARES: Galactic plane (preliminary)

- **Data period:** 2007–2020
- **Events:** tracks + showers, using existing diffuse neutrino selections
- **Signal hypothesis:** looking for signal in the region $|\ell| < 30^\circ$ and $|b| < 2^\circ$ assuming a simple power-law



$$\frac{dN}{dEdtd\Omega} = \phi_0 \times \left(\frac{E}{40 \text{ TeV}} \right)^{-\gamma} \text{ in } [\text{GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}] \text{ per flavor}$$



Best-fit: $\gamma = 2.45$, $\phi_0(40 \text{ TeV}) = 4.0 \text{e-16}$



Likelihood analysis of Cygnus X region



Source position is consistent with either HAWC or LHAASO extended source position.

Neutrino flux level is consistent with the γ -ray flux of the extended Cocoon source.

High-Energy Cosmic Neutrinos


francis halzen



- ANTARES, Baikal-GVD,IceCube
- the diffuse high-energy neutrino flux
- observation of the first sources
- multimessenger astronomy: plan B



one year of IceCube neutrinos >100 GeV

(reaches neutrino purity of 97% but overwhelmingly atmospheric)



~ 200 cosmic neutrinos

~12 separated from atmospheric background with $E>60$ TeV




- maximize the likelihood L at each point in the sky
- usually, add energy term to the signal likelihood S

$$L(n_s, x_s, \gamma) = \prod_i^{events} \left(\frac{n_s}{N} S_i(|x_i - x_s| \sigma_i, E_i, \gamma) + \frac{N - n_s}{N} B_i(\delta_i, E_i) \right)$$

\downarrow
 $S_i(|\vec{x}_i - \vec{x}_s|, \sigma_i) = \frac{1}{2\pi\sigma_i^2} \exp\left(-\frac{|\vec{x}_i - \vec{x}_s|^2}{2\sigma_i^2}\right)$

evidence for non-uniform sky map in 10 years of IceCube data : mostly resulting from 4 extragalactic source candidates



pre-trial p-value for clustering of high energy neutrinos

IceCube, PRL 124, 051103 (2020)

Source List Results							
Name	Class	α [deg]	δ [deg]	\hat{n}_s	$\hat{\gamma}$	$-\log_{10}(p_{local})$	$\phi_{90\%}$
PKS 2320-035	FSRQ	350.88	-3.29	4.8	3.6	0.45	3.3
3C 454.3	FSRQ	343.50	16.15	5.4	2.2	0.62	5.1
TXS 2241+406	FSRQ	341.06	40.96	3.8	3.8	0.42	5.6
RGB J2243+203	BLL	340.99	20.36	0.0	3.0	0.33	3.1
CTA 102	FSRQ	338.15	11.73	0.0	2.7	0.30	2.8
BL Lac	BLL	330.69	42.28	0.0	2.7	0.31	4.9
OX 169	FSRQ	325.89	17.73	2.0	1.7	0.69	5.1
B2 2114+33	BLL	319.06	33.66	0.0	3.0	0.30	3.9
PKS 2032+107	FSRQ	308.84	-0.94	0.0	2.4	0.33	3.5
2HVS J0814+42	GAL	307.91	13.1	3.4	0.8	0.47	9.0
Gamma Cygni	GAL	305.56	40.26	7.4	3.7	0.59	6.9
MGRO J2019+37	GAL	304.85	36.80	0.0	3.1	0.33	4.0
MG2 J201534+3710	FSRQ	303.92	37.19	4.4	4.0	0.40	5.6
MG4 J200112+4352	BLL	300.30	43.89	6.1	2.3	0.67	7.8
1ES 1959+650	BLL	300.01	65.15	12.6	3.3	0.77	12.3
1RXS J194246.3+1	BLL	295.70	10.56	0.0	2.7	0.33	2.6
RX J1931.1+0937	BLL	292.78	9.63	0.0	2.9	0.29	2.8
NVSS J190836-012	UNIDB	287.20	-1.53	0.0	2.9	0.22	2.3
MGRO J1908+06	GAL	287.17	6.18	4.2	2.0	1.42	5.7
TXS 1902+556	BLL	285.80	55.68	11.7	4.0	0.85	9.9
HESS J1857+026	GAL	284.30	2.67	7.4	3.1	0.53	3.5
GRS 1285.0	UNIDB	283.15	0.69	1.7	3.8	0.27	2.3
HESS J1852-000	GAL	283.00	0.00	3.3	3.7	0.38	2.6
HESS J1849-000	GAL	282.26	-0.02	0.0	3.0	0.28	2.2
HESS J1843-033	GAL	280.75	-3.30	0.0	2.8	0.31	2.5
OT 081	BLL	267.87	9.65	12.2	3.2	0.73	4.8
S4 1749+70	BLL	267.15	70.10	0.0	2.5	0.37	8.0
1H 1720+117	BLL	261.27	11.88	0.0	2.7	0.30	3.2
PKS 1717+177	BLL	259.81	17.75	19.8	3.6	1.32	7.3
Mkn 501	BLL	253.47	39.76	10.3	4.0	0.61	7.3
4C +38.41	FSRQ	248.82	38.14	4.2	2.3	0.66	7.0
PG 1553+113	BLL	238.93	11.19	0.0	2.8	0.32	3.2
GB6 J1542+6129	BLL	235.75	61.50	29.7	3.0	2.74	22.0
B2 1520+31	FSRQ	230.55	31.74	7.1	2.4	0.83	7.3
PKS 1502+036	AGN	226.26	3.44	0.0	2.7	0.28	2.9
PKS 1502+106	FSRQ	226.10	10.50	0.0	3.0	0.33	2.6
PKS 1441+25	FSRQ	220.99	25.03	7.5	2.4	0.94	7.3
PKS 1424+240	BLL	216.76	23.80	41.5	3.9	2.80	12.3
NVSS J141826-023	BLL	214.61	-2.56	0.0	3.0	0.25	2.0
B3 1343+451	FSRQ	206.40	44.88	0.0	2.8	0.39	5.0
S4 1250+53	BLL	193.31	53.02	2.2	2.5	0.39	5.9
PG 1246+586	BLL	192.08	58.34	0.0	2.8	0.35	6.4
MG1 J123931+0443	FSRQ	189.89	4.73	0.0	2.6	0.28	2.4
M 87	AGN	187.71	12.39	0.0	2.8	0.29	3.1
ON 246	BLL	187.56	25.30	0.9	1.7	0.37	4.2
3C 273	FSRQ	187.27	2.04	0.0	3.0	0.28	1.9
4C +21.35	FSRQ	186.23	21.38	0.0	2.6	0.32	3.5
W Comae	BLL	185.38	28.24	0.0	3.0	0.32	3.7
PG 1218+304	BLL	185.34	30.17	11.1	3.9	0.70	6.7
PKS 1216-010	BLL	184.64	-1.33	6.9	4.0	0.45	3.1
B2 1215+30	BLL	184.48	30.12	18.6	3.4	1.09	8.5
Ton 599	FSRQ	179.88	29.24	0.0	2.2	0.29	4.5

avoid $>10^5$ trials \rightarrow search 110 preselected source candidates

PKS B1130+008	BLL	173.20	0.58	15.8	4.0	0.96	4.4
Mkn 421	BLL	166.12	38.21	2.1	1.9	0.38	5.3
4C +01.28	BLL	164.61	1.56	0.0	2.9	0.26	2.4
1H 1013+498	BLL	153.77	49.43	0.0	2.6	0.29	4.5
4C +55.17	FSRQ	149.42	55.38	11.9	3.3	1.02	10.6
M 82	SBG	148.95	69.67	0.0	2.6	0.36	8.8
PMN J0948+0022	AGN	147.24	0.37	9.3	4.0	0.76	3.9
OJ 287	BLL	133.71	20.12	0.0	2.6	0.32	3.5
PKS 0829+046	BLL	127.97	4.49	0.0	2.9	0.28	2.1
S4 0814+42	BLL	124.56	42.38	0.0	2.3	0.30	4.9
OJ 014	BLL	122.87	1.78	16.1	4.0	0.99	4.4
1H 0808+042	BLL	122.40	52.01	0.0	2.8	0.20	4.7
PKS 0730+01	FSRQ	114.82	1.62	0.0	2.8	0.20	2.4
PKS 0735+17	BLL	114.54	17.71	0.0	2.8	0.30	3.5
4C +14.23	FSRQ	111.33	14.42	8.5	2.9	0.60	4.8
S5 0716+71	BLL	110.49	71.34	0.0	2.5	0.38	7.4
PSR B0656+14	GAL	104.95	14.24	8.4	4.0	0.51	4.4
1ES 0647+250	BLL	102.70	25.06	0.0	2.9	0.27	3.0
B3 0609+413	BLL	93.22	41.37	1.8	1.7	0.42	5.3
Crab nebula	GAL	83.63	22.01	1.1	2.2	0.31	3.7
OG +050	FSRQ	83.18	7.55	0.0	3.2	0.28	2.9
TXS 0518+211	BLL	80.44	21.21	15.7	3.8	0.92	6.6
TXS 0506+056	BLL	77.35	5.70	12.3	2.1	3.72	10.1
PKS 0502+049	FSRQ	76.34	5.00	11.2	3.0	0.66	4.1
S3 0458-02	FSRQ	75.30	-1.97	5.5	4.0	0.33	2.7
PKS 0440-00	FSRQ	70.66	-0.29	7.6	3.9	0.46	3.1
MG2 J043337+2905	BLL	68.41	29.10	0.0	2.7	0.28	4.5
PKS 0422+00	BLL	66.19	0.60	0.0	2.9	0.27	2.3
PKS 0420-01	FSRQ	65.83	-1.33	9.3	4.0	0.52	3.4
PKS 0336-01	FSRQ	54.88	-1.77	15.5	4.0	0.99	4.4
NGC 1275	AGN	49.96	41.51	3.6	3.1	0.41	5.5
NGC 1068	SBG	40.67	-0.01	50.4	3.2	4.74	10.5
PKS 0235+164	BLL	39.67	16.62	0.0	3.0	0.28	3.1
4C +28.07	FSRQ	39.48	28.80	0.0	2.8	0.30	3.6
3C 66A	BLL	35.67	43.04	0.0	2.8	0.30	3.9
B2 0218+357	FSRQ	35.28	35.94	0.0	3.1	0.33	4.3
PKS 0215+015	FSRQ	34.46	1.74	0.0	3.2	0.27	2.3
MG1 J021114+1051	BLL	32.81	10.86	1.6	1.7	0.43	3.5
TXS 0141+268	BLL	26.15	27.09	0.0	2.5	0.31	3.5
B3 0133+388	BLL	24.14	39.10	0.0	2.6	0.28	4.1
NGC 598	SBG	23.52	30.62	11.4	4.0	0.63	6.3
S2 0109+22	BLL	18.03	22.75	2.0	3.1	0.30	3.7
4C +01.02	FSRQ	17.16	1.59	0.0	3.0	0.26	2.4
M 31	SBG	10.82	41.24	11.0	4.0	1.09	9.6
PKS 0019+058	BLL	5.64	6.14	0.0	2.9	0.29	2.4
PKS 2233-148	BLL	339.14	-14.56	5.3	2.8	1.26	21.4
HESS J1841-055	GAL	280.23	-5.55	3.6	4.0	0.55	4.8
HESS J1837-069	GAL	279.43	-6.93	0.0	2.8	0.30	4.0
PKS 1510-089	FSRQ	228.21	-9.10	0.1	1.7	0.41	7.1
PKS 1329-049	FSRQ	203.02	-5.16	6.1	2.7	0.77	5.1
NGC 4945	SBG	196.36	-49.47	0.3	2.6	0.31	50.2
3C 279	FSRQ	194.04	-5.79	0.3	2.4	0.20	2.7
PKS 0805-07	FSRQ	122.07	-7.86	0.0	2.7	0.31	4.7
PKS 0727-11	FSRQ	112.58	-11.69	1.9	3.5	0.59	11.4
LMC	SBG	80.00	-68.75	0.0	3.1	0.36	41.1
SMC	SBG	14.50	-72.75	0.0	2.4	0.37	44.1
PKS 0048-09	BLL	12.68	-9.49	3.9	3.3	0.87	10.0
NGC 253	SBG	11.90	-25.29	3.0	4.0	0.75	37.7

Source List Results							
Name	Class	α [deg]	δ [deg]	\hat{n}_s	$\hat{\gamma}$	$-\log_{10}(p_{local})$	$\phi_{90\%}$
PKS 2320-035	FSRQ	350.88	-3.29	4.8	3.6	0.45	3.3
3C 454.3	FSRQ	343.50	16.15	5.4	2.2	0.62	5.1
TXS 2241+406	FSRQ	341.06	40.96	3.8	3.8	0.42	5.6
RGB J2243+203	BLL	340.99	20.36	0.0	3.0	0.33	3.1
CTA 109	FSRQ	228.15	11.72	0.0	2.7	0.20	2.8
BL L							
OX 1							
B2 211							
PKS 203							
2HVS 100							
Gamma							
MGRO J2							
MG2 J2015							
MG4 J2001							
1ES 195							
1RXS J194							
RX J1931							
NVSS J190							
MGRO J1							
TXS 190							
HESS J18							
GRS 12							
HESS J18							
HESS J18							
HESS J18							
OT 0							
S4 174							
1H 1720							
PKS 171							
Mkn 1							
4C +3							
PG 1553							
GB6 J154							
B2 1520							
PKS 150							
PKS 150							
PKS 144							
PKS 142							
NVSS J141							
B3 1343							
S4 1250							
PG 1246							
MG1 J1239							
M 8							
ON 2							
3C 2							
4C +2							
W Col							
PG 1218+304	BLL	185.34	30.17	11.1	3.9	0.70	6.7
PKS 1216-010	BLL	184.64	-1.33	6.9	4.0	0.45	3.1
B2 1215+30	BLL	184.48	30.12	18.6	3.4	1.09	8.5
Ton 599	FSRQ	179.88	29.24	0.0	2.2	0.29	4.5

PKS B1130+008	BLL	173.20	0.58	15.8	4.0	0.96	4.4
Mkn 421	BLL	166.12	38.21	2.1	1.9	0.38	5.3
4C +01.28	BLL	164.61	1.56	0.0	2.9	0.26	2.4
1H 1013+498	BLL	153.77	49.43	0.0	2.6	0.29	4.5
4C +55.17	FSRQ	149.42	55.38	11.9	3.3	1.02	10.6
M 82	SBG	148.95	69.67	0.0	2.6	0.36	8.8
PKS 10048+0029	AGN	147.24	0.27	0.2	4.0	0.76	3.9

Based on 4th Fermi catalog of gamma-ray sources: 4FGL-2DR

Selected a priori based on gamma-ray brightness and IceCube sensitivity at object's declination


95 are blazars

14 are AGN / other galaxies

1 Galactic source (MGRO J1908+06)

PG 1218+304	BLL	185.34	30.17	11.1	3.9	0.70	6.7
PKS 1216-010	BLL	184.64	-1.33	6.9	4.0	0.45	3.1
B2 1215+30	BLL	184.48	30.12	18.6	3.4	1.09	8.5
Ton 599	FSRQ	179.88	29.24	0.0	2.2	0.29	4.5

F180 U121-11	FSRQ	112.30	-11.03	1.9	3.0	0.99	11.4
LMC	SBG	80.00	-68.75	0.0	3.1	0.36	41.1
SMC	SBG	14.50	-72.75	0.0	2.4	0.37	44.1
PKS 0048-09	BLL	12.68	-9.49	3.9	3.3	0.87	10.0
NGC 253	SBG	11.90	-25.29	3.0	4.0	0.75	37.7




limits and interesting fluctuations ?



a rotating supermassive
black hole


Illustr.

NGC 1068: an obscured cosmic accelerator




neutrino production in obscured cores of active galaxies


- electrons and protons are accelerated in the high field regions associated with the black hole and the accretion disk
- produce neutrinos in the optically thick corona



NGC 1068 neutrinos: the disk corona model




the radiatively obscured core of an active galaxy: opaque to γ -rays



[PS: the neutrinos are not produced by star formation because they are not accompanied by gamma rays]

NEUTRINO BEAMS



the p γ efficiency dilemma

- efficiency for producing the neutrinos in the photon target:

$$\tau_{p\gamma} \simeq \frac{\kappa_{p\gamma} R_{\text{escape}}}{\lambda_{p\gamma}} \simeq R_{\text{escape}} \sigma_{p\gamma} n_{\text{photons}}$$

- likelihood of the multimessenger photons to be absorbed in target

$$\tau_{\gamma\gamma} \simeq R_{\text{target}} \sigma_{\gamma\gamma} n_{\text{photons}}$$


→ therefore, with $R_{\text{escape}} \sim R_{\text{target}}$

$$\tau_{\gamma\gamma} \sim \frac{\sigma_{\gamma\gamma}}{\sigma_{p\gamma}} \tau_{p\gamma} \sim 300 \tau_{p\gamma}$$

→ do not expect high energy gamma rays to accompany cosmic neutrinos

→ blazar jets are out

neutrinos produced in the gamma-ray obscured core of NGC 1068



the cocoon absorbs the protons to produce neutrinos

by dimensional analysis:

$$\tau_{\gamma\gamma} \simeq 10^5 \left[\frac{R_s}{R} \right] \left[\frac{1 \text{ keV}}{E_X} \right] \left[\frac{L_X}{L_{edd}} \right] \sim 300 \tau_{p\gamma}$$

NGC 1068:

$$E_X = 1 \text{ keV}; L_X \sim 10^{43} \text{ ergs}^{-1} \text{ and } M = 10^7 M_{\text{sun}}$$


the size of the neutrino production region (or the gamma ray absorption region for the case of pp) is $10 \sim 100$ Schwarzschild radii!

$$R_s = [2GM]/c^2 = 3 \times 10^5 \text{ cm} \left[\frac{M}{M_{\text{sun}}} \right] \simeq 0.1 R$$

$$L_{edd} = \frac{4\pi GMm_p c}{\sigma_T} = 1.2 \times 10^{38} \frac{\text{erg}}{\text{s}} \left[\frac{M}{M_{\text{sun}}} \right] \simeq 10^2 L_X$$



interesting fluctuations or neutrino sources?

- improved detector geometry and calibration (each PMT calibrated individually)
- improved muon angular resolution and energy reconstruction
 - DNN (energy) and BDT (pointing) reconstruction
 - *point spread function consistent with simulation*
 - insensitive to systematics
 - improved characterization of the optics of the ice



applied to 10 years of archival data (pass 2),
data unblinded, answer...


pointing with neutrinos:



directional distributions:


- better modeling of the directional distribution of neutrinos
- consistent with full Monte-Carlo simulations of the detector

the new IceCube neutrino map



1% of scrambled data sets have a spot $\geq 5.3\sigma$

is the hot spot coincident with one of the 110 preselected sources?



at the NGC1068 location:


astrophysical neutrino events = 79^{+22}_{-20}

spectral index = 3.2 ± 0.2


single source significance 5.2σ


1 in 100,000 scrambled data sets have object $\geq 5.2\sigma = 4.2\sigma$ evidence

another look at the result




- measured astrophysical neutrino events = 79^{+22}_{-20}
- the angular distribution of the events matches simulation






limits and interesting fluctuations ?

search of a list of candidate sources
selected prior to the analysis



Catalog of 110 Objects

Binomial Test



1 in 10,000 scrambled data sets have
binomial test result $-\log_{10}(p_{min}) \geq 5.3$

= 3.7σ evidence

Source Name	Source Type	α [°]	δ [°]	\hat{n}_s	$\hat{\gamma}$	$-\log_{10} p_{local}$	$\Phi_{90\%}$
NGC 1068	SBG/AGN	40.67	-0.01	79	3.2	7.0 (5.2 σ)	9.6
PKS 1424+240	BLL	216.76	23.80	77	3.5	4.0 (3.7 σ)	11.4
TXS 0506+056	BLL/FSRQ	77.36	5.70	5	2.0	3.6 (3.5 σ)	7.5
PKS 0019+058	BLL	5.64	6.13	1	2.4	0.4 (0.2 σ)	2.6
1ES 0033+595 (*)	BLL	8.98	59.83	0	4.3	0.0 (0.0 σ)	5.0
M 31	GAL	10.82	41.24	13	3.3	0.8 (1.0 σ)	6.2
4C +01.02	FSRQ	17.17	1.58	0	4.3	0.0 (0.0 σ)	2.1
S2 0109+22	BLL	18.03	22.75	10	2.8	0.7 (0.8 σ)	4.8
B3 0133+388	BLL	24.14	39.10	0	4.3	0.0 (0.0 σ)	3.8
TXS 0141+268	BLL	26.15	27.09	0	4.3	0.0 (0.0 σ)	3.2
MITG J021114+1051	BLL	32.81	10.86	0	4.3	0.0 (0.0 σ)	2.6
PKS 0015+015	FSRQ	34.46	1.72	2	3.9	0.2 (0.0 σ)	1.9

Source Name	Source Type	α [°]	δ [°]	\hat{n}_s	$\hat{\gamma}$	$-\log_{10} p_{local}$	$\Phi_{90\%}$
PG 1218+304	BLL	185.34	30.17	0	3.1	0.0 (0.0 σ)	3.4
W Comae	BLL	185.38	28.24	0	4.3	0.0 (0.0 σ)	3.3
4C+21.35	FSRQ	186.23	21.38	0	4.3	0.0 (0.0 σ)	3.2
3C 273	FSRQ	187.27	2.03	21	4.3	0.6 (0.7 σ)	3.0
ON 246	BLL	187.56	25.30	0	4.3	0.0 (0.0 σ)	3.2
M 87	RDG	187.71	12.39	0	0.6	0.0 (0.0 σ)	2.8
MITG J123931+0443	FSRQ	189.89	4.73	0	4.3	0.0 (0.0 σ)	2.4
PG 1246+586	BLL	192.08	58.34	0	4.3	0.0 (0.0 σ)	4.8
S4 1250+53	BLL	193.31	53.02	0	4.3	0.0 (0.0 σ)	4.0
B3 1343+451	FSRQ	206.39	44.88	5	2.9	0.4 (0.3 σ)	4.7
NGC 5380 (*)	GAL	209.33	37.50	4	2.4	0.9 (1.2 σ)	6.4
NVSS J141826-023336	BLL	214.61	-2.56	6	3.9	0.4 (0.4 σ)	2.3
PKS 1441+25	FSRQ	220.99	25.03	3	2.1	0.7 (0.9 σ)	5.0
TXS 1452+516 (*)	BLL	223.62	51.41	0	2.3	0.0 (0.0 σ)	4.4
PKS 1502+106	FSRQ	226.10	10.50	1	1.8	0.5 (0.5 σ)	3.4
NLSY1	PKS 1502+036	226.27	3.45	0	4.3	0.0 (0.0 σ)	2.1
B2 1520+31	FSRQ	230.55	31.74	35	4.3	1.0 (1.3 σ)	6.2
Arp 220 (*)	SBG	233.70	23.53	0	4.3	0.0 (0.0 σ)	3.1
GB6 J1542+6129	BLL	235.76	61.50	16	3.0	1.9 (2.2 σ)	13.0
PG 1553+113	BLL	238.93	11.19	2	4.3	0.2 (0.0 σ)	2.3
4C+15.54 (*)	BLL	241.77	15.84	0	4.3	0.0 (0.0 σ)	2.9
4C+38.41	FSRQ	248.82	38.14	4	2.3	0.9 (1.1 σ)	6.2
Mkn 501	BLL	253.47	39.76	15	4.3	0.5 (0.5 σ)	5.0
PKS 1717+177	BLL	259.81	17.75	34	4.3	1.0 (1.2 σ)	5.1
IH 1720+117	BLL	261.27	11.87	0	4.3	0.0 (0.0 σ)	2.7
S4 1749+70	BLL	267.16	70.10	0	4.3	0.0 (0.0 σ)	6.6
OT 081	BLL	267.88	9.65	0	2.9	0.0 (0.0 σ)	2.7
RX J1754.1+3212 (*)	BLL	268.55	32.20	0	4.3	0.0 (0.0 σ)	3.4
S5 1803+784 (*)	BLL	270.17	78.47	0	2.7	0.0 (0.0 σ)	7.5
NVSS J184425+154646 (*)	BLL	281.12	15.79	11	4.3	0.4 (0.2 σ)	3.1
LQAC 284+003 (*)	BCU	284.48	3.22	12	2.5	2.0 (2.2 σ)	5.2
TXS 1902+556	BLL	285.81	55.68	3	4.3	0.3 (0.0 σ)	4.6
MGRO J1908+06	UID	286.91	6.32	2	1.8	1.4 (1.7 σ)	4.8
RX J1931.1+0937	BLL	292.78	9.63	15	4.3	0.5 (0.4 σ)	3.1
87GB 194024.3+102612 (*)	BLL	295.70	10.56	0	4.3	0.0 (0.0 σ)	2.6
IES 1959+650	BLL	300.01	65.15	8	3.4	0.5 (0.4 σ)	7.2

MITG J200112+4352	BLL	300.30	43.89	3	4.3	0.3 (0.0 σ)	3.6
7C 2010+4619 (*)	BLL	303.02	46.49	4	2.5	0.7 (0.9 σ)	6.4
MITG J201534+3710	FSRQ	303.89	37.18	19	3.6	0.7 (0.9 σ)	5.5
PKS 2032+107	FSRQ	308.85	10.94	0	4.3	0.0 (0.0 σ)	2.8
B2 2114+33	BLL	319.06	33.66	12	2.9	0.8 (0.9 σ)	5.7
OX 169	FSRQ	325.89	17.73	4	4.3	0.3 (0.0 σ)	2.7
BL Lac	BLL	330.69	42.28	11	4.3	0.4 (0.3 σ)	4.7
CTA 102	FSRQ	338.15	11.73	0	4.3	0.0 (0.0 σ)	2.6
B2 2234+28A (*)	FSRQ	339.10	28.48	8	3.2	0.4 (0.3 σ)	4.1
RGB J2243+203	BLL	340.99	20.36	5	3.6	0.3 (0.0 σ)	2.8
TXS 2241+406	FSRQ	341.06	40.96	0	4.3	0.0 (0.0 σ)	3.9
3C 454.3	FSRQ	343.50	16.15	1	1.5	1.2 (1.6 σ)	5.5
B2 2308+34 (*)	FSRQ	347.77	34.42	19	3.6	0.7 (0.9 σ)	5.6

B2 1215+30	BLL	184.48	30.12	15	3.1	0.9 (1.1 σ)	5.7
PKS 1216-010	BLL	184.64	-1.33	0	3.7	0.0 (0.0 σ)	2.0

top 5 hot spots for 3 searches performed

TXS 0506+056 ➔

	α [°]	δ [°]	$\hat{\mu}_{ns}$	$\hat{\gamma}$	$-\log_{10}(p_{local})$
$\gamma = 2.0$					
#1	76.93	12.90	13.4	2.00	6.08
#2	9.76	7.50	4.9	2.00	5.04
#3	77.37	5.57	6.2	2.00	4.88
#4	179.25	52.44	5.5	2.00	4.87
#5	202.63	33.89	7.1	2.00	4.74

NGC 1068 ➔

	α [°]	δ [°]	$\hat{\mu}_{ns}$	$\hat{\gamma}$	$-\log_{10}(p_{local})$
$\gamma = 2.5$					
#1	40.65	0.09	36.8	2.50	5.84
#2	177.91	23.24	21.4	2.50	5.45
#3	105.78	1.03	23.6	2.50	5.17
#4	182.46	39.52	22.2	2.50	4.91
#5	180.16	42.21	26.0	2.50	4.86

NGC 1068 ➔

TXS 0506+056 ➔

	α [°]	δ [°]	$\hat{\mu}_{ns}$	$\hat{\gamma}$	$-\log_{10}(p_{local})$
Free γ					
#1	40.69	0.09	80.7	3.20	7.30
#2	297.27	27.45	69.8	3.24	5.51
#3	76.93	12.90	11.2	1.81	5.37
#4	180.20	42.19	47.8	3.03	4.80
#5	208.15	23.16	55.5	3.19	4.60

NUCLEAR EMISSION IN SPIRAL NEBULAE*

CARL K. SEYFERT†

1943

ABSTRACT

Spectrograms of dispersion 37–200 Å/mm have been obtained of six extragalactic nebulae with high-excitation nuclear emission lines superposed on a normal G-type spectrum. All the stronger emission lines from $\lambda 3727$ to $\lambda 6731$ found in planetaries like NGC 7027 appear in the spectra of the two brightest spirals observed, NGC 1068 and NGC 4151.

How are neutrinos produced in non-jetted AGN?

We conclude that active galactic accelerating particles to cosmic galactic cosmic rays is likely to particular, in the Virgo supercluster. NGC 4151 and NGC 1068 are likely "local" metagalactic cosmic rays, the ultra-high energy ($E \geq 10^{19}$ eV) air showers. The energy density of photons in the immediate vicinity of a black hole may be too high (Blumenthal, 1970) to permit the acceleration of protons beyond $\sim 10^{14}$ eV, (except by beaming processes). The highest energy protons hence are accelerated somewhat farther out, or else by beaming (Lovelace, 1976). Gamma rays from the ergosphere of a black hole are degraded at energies above ~ 1 MeV, and from a spinar, above ~ 1 GeV. Neutrinos are not thus affected and would provide information on very high energy particles in active galactic nuclei.

1982

R. Silberberg and M. M. Shapiro

Laboratory for Cosmic Ray Physics
Naval Research Laboratory
Washington, D.C. 20375

High-Energy Cosmic Neutrinos

francis halzen



- ANTARES, Baikal-GVD, IceCube
- the diffuse high-energy neutrino flux
- observation of the first sources
- multimessenger astronomy: plan B

RESEARCH ARTICLE SUMMARY

NEUTRINO ASTROPHYSICS

Multimessenger observations of a flaring blazar coincident with high-energy neutrino IceCube-170922A


The IceCube Collaboration, *Fermi-LAT*, **MAGIC**, *AGILE*, **ASAS-SN**, **HAWC**, **H.E.S.S.**, *INTEGRAL*, **Kanata**, **Kiso**, **Kapteyn**, **Liverpool Telescope**, **Subaru**, *Swift/NuSTAR*, **VERITAS**, and **VLA/17B-403 teams***†

RESEARCH ARTICLE

NEUTRINO ASTROPHYSICS

Neutrino emission from the direction of the blazar TXS 0506+056 prior to the IceCube-170922A alert


IceCube Collaboration*†



MAGIC
detects emission of
 > 100 GeV gammas

IceCube 170922
290 TeV

Fermi
detects a flaring
blazar within 0.06°



MASTER robotic optical telescope network: after 73 seconds

Follow-up detections of IC170922 based on public telegrams



IceCube

September 22



Swift

September 26



Fermi, ASAS-SN

September 28



SALT, Kapteyn

October 7



MAGIC

October 4



Liverpool, AGILE

September 29



Kanata, NuSTAR

October 12



VLA


October 17



Subaru


October 25





search in archival
IceCube data:

- 100-day flare in 2014
- spectrum $E^{-2.2}$
- $L_v > 10^{47}$ erg/s
- no gamma ray flare!



global robotic network of
optical telescopes
connects TXS 0506+056
to IC170922A in the time
domain




“MASTER found the blazar in the off-state *after one minute* and then switched to on-state two hours after the event. The effect is observed at a 50-sigma significance level”


Optical Observations Reveal Strong Evidence for High Energy Neutrino Progenitor

V.M. Lipunov^{1,2}, V.G. Kornilov^{1,2}, K.Zhirkov¹, E. Gorbovskoy², N.M. Budnev⁴, D.A.H.Buckley³, R. Rebolo⁵, M. Serra-Ricart⁵, R. Podesta^{9,10}, N.Tyurina², O. Gress^{4,2}, Yu.Sergienko⁸, V. Yurkov⁸, A. Gabovich⁸, P.Balanutsa², I.Gorbunov², D.Vlasenko^{1,2}, F.Balakin^{1,2}, V.Topolev¹, A.Pozdnyakov¹, A.Kuznetsov², V.Vladimirov², A. Chasovnikov¹, D. Kuvshinov^{1,2}, V.Grishpun^{1,2}, E.Minkina^{1,2}, V.B.Petkov⁷, S.I.Svertilov^{2,6}, C. Lopez⁹, F. Podesta⁹, H.Levato¹⁰, A. Tlatov¹¹, B. Van Soelen¹², S. Razzaque¹³, M. Böttcher¹⁴


neutrino




gamma ray




optical



TXS 0506+056



multimessenger observations of TXS 0506 + 056



Upward-going cascade **GVD210418CA**

MJD = 59322.94855324

Energy E = 224 TeV ($\pm 30\%$)

Distance from central string r = 70 m


Zenith angle = 115°

RA = 82.4° , Dec = 7.1°

Probability for the background-only hypothesis (stat.+sys.)


P-value = 0.0033 (2.93σ)

Signalness: 97.1%




A high energy neutrino from the direction of TXS 0506+056

GVD210418CA (97% signalness) lies within 90% error circle from TXS 0506+056



The chance probability for such an association to occur randomly due to the background is $p = 0.0074$

Radio and gamma-ray light curves of TXS 0506+056.




Analysis of RATAN-600 radiotelescope data (11GHz) showed increased activity

- IC event registered during γ flare and radio activity
- Baikal-GVD event during radio activity
- Probability of IC non-observation: 11%

arXiv:2210.01650

```
[ I3EventHeader:  
  StartTime: 2022-09-18 12:46:05.322,758,760,8 UTC  
  EndTime : 2022-09-18 12:46:05.322,778,378,4 UTC  
  RunID : 137065  
  SubrunID : 0  
  EventID : 22012496  
  SubEventID : 0  
  SubEventStream : InIceSplit  
 ]
```




IC220918


within 90% error circle of TXS 0506+056

TXS 0506+056

- two statistically independent observations above the $> 3\sigma$ level
- it is also the second source in the all-sky search at 3.7σ
- high-statistic association of IC170922 with optical variation in time domain
- TXS 0506+056 within the error circle of IC220918 and the highest energy event observed by Baikal-GVD
- supported by TeV gamma ray and by radio imaging of the core (jet loses its tight collimation after 5 milliarcseconds)



NGC 1068




neutrino astronomy 2022

- it exists
- more neutrinos, better neutrinos, more telescopes
- closing in on cosmic ray sources
- [are active galaxies with obscured cores the sources of cosmic rays?]




overflow slides

Neutrinos? Perfect Messengers




- electrically neutral
- massless (in this talk)
- unabsorbed
- unlike γ rays, neutrinos are solely created in processes involving cosmic rays
- ... but difficult to detect


multimessenger astronomy




- PeV γ 's lose energy on CMB photons
- gamma rays may also lose energy in the target that produces neutrinos even before reaching the EBL
- efficient neutrino production sites are likely to be optically thick to gamma rays; expect no correlation between gamma-ray and neutrino activity




10,000 times too small to
do neutrino astronomy...





- 3 km deep glacier at geographic South Pole
- we transformed 1 km³ of Antarctic ice below 1.5 km into a Cherenkov detector





IceCube
5160 photomultipliers
instrument one km³ of
Antarctic ice between
1.4 and 2.4 km depth

photomultiplier
tube -10 inch



the IceCube Neutrino Observatory







ice: step by step

- hole ice ?





- birefringence




- o air bubbles/hydrates
below 1350 m

- ice layers





A



B

Event 135440/3139778-0
Time 2021-06-29 18:09:44 UTC
Duration 22320.7 ns



HIGH-ENERGY EVENTS NOW PUBLIC ALERTS!


We send our high-energy events in real-time as public GCN alerts now!

47

TITLE: GCN/AMON NOTICE
NOTICE_DATE: Wed 27 Apr 16 23:24:24 UT
NOTICE_TYPE: AMON ICECUBE HESE
RUN_NUM: 127853
EVENT_NUM: 67093193
SRC_RA: 240.5683d {+16h 02m 16s} (J2000),
240.7644d {+16h 03m 03s} (current),
239.9678d {+15h 59m 52s} (1950)
SRC_DEC: +9.3417d {+09d 20' 30"} (J2000),
+9.2972d {+09d 17' 50"} (current),
+9.4798d {+09d 28' 47"} (1950)
SRC_ERROR: 35.99 [arcmin radius, stat+sys, 90% containment]
SRC_ERROR50: 0.00 [arcmin radius, stat+sys, 50% containment]
DISCOVERY_DATE: 17505 TJD; 118 DOY; 16/04/27 (yy/mm/dd)
DISCOVERY_TIME: 21152 SOD {05:52:32.00} UT
REVISION: 2
N_EVENTS: 1 [number of neutrinos]
STREAM: 1
DELTA_T: 0.0000 [sec]
SIGMA_T: 0.0000 [sec]
FALSE_POS: 0.0000e+00 [s^-1 sr^-1]
PVALUE: 0.0000e+00 [dn]
CHARGE: 18883.62 [pe]
SIGNAL_TRACKNESS: 0.92 [dn]
SUN_POSTN: 35.75d {+02h 23m 00s} +14.21d {+14d 12' 45"}


GCN notice for starting track sent Apr 27

We send **rough reconstructions** first and then **update them**.




from light in the ice to astronomer in less than one minute

electron and tau neutrinos (showers)



muon neutrinos through Earth (tracks)

NEUTRINO BEAMS: HEAVEN & EARTH



- a target efficient at converting protons into neutrinos is unlikely to be transparent to high energy photons.
- IC170922? TXS 0506+056 is not a blazar when neutrinos are emitted as confirmed by gamma ray, optical and radio observations

RADIO INTERFEROMETRY



1912.01743v1
[astro-ph.GA]


2017.88

- core brightening observed in a radio burst that started 5 years ago
- beyond 5 milliarcseconds the jet loses its tight collimation




Peak: 1256.0, RMS: 0.09 mJy/beam



beyond 5 milliarcseconds the jet loses its tight collimation



- jet found a target after tens of pc to produce neutrinos
- obscures the gamma rays
- a massive star in the host galaxy, the jet of a merging galaxy, warped jet, structured jet...?





gamma rays in 2017 at the time the neutrino is produced ?
 a few \sim 10 GeV photons and not much else, consistent with
 an obscured source, not a blazar



- MAGIC, HESS and VERITAS: no TeV gamma rays at the time the neutrino was produced
- MAGIC: onset of the TeV flux 5 days after IC170922
- confirmed by MASTER: the blazar switches from the “off” to “on” state 2 hours after the neutrino

a second cosmic ray source ?



[Previous | Next]

Neutrino candidate source FSRQ PKS 1502+106 at highest flux density at 15 GHz

ATel #12996; **S. Kiehlmann (IoA FORTH, OVRO), T. Hovatta (FINCA), M. Kadler (Univ. WÄ4rzburg), W. Max-Moerbeck (Univ. de Chile), A. C.S. Readhead (OVRO)**
on 7 Aug 2019; 12:31 UT

Credential Certification: Sebastian Kiehlmann (skiehlmann@mail.de)

Subjects: Radio, Neutrinos, AGN, Blazar, Quasar



On 2019/07/30.86853 UT IceCube detected a high-energy astrophysical neutrino candidate (Atel #12967). The FSRQ PKS 1502+106 is located within the 50% uncertainty region of the event. We report that the flux density at 15 GHz measured with the OVRO 40m Telescope shows a long-term outburst that started in 2014, which is currently reaching an all-time high of about 4 Jy, since the beginning of the OVRO measurements in 2008. A similar 15 GHz long-term outburst was seen in TXS 0506+056 during the neutrino event [IceCube-170922A](#).


Related

- 12996 [Neutrino candidate source FSRQ PKS 1502+106 at highest flux density at 15 GHz](#)
- 12985 [IceCube-190730A: Swift XRT and UVOT Follow-up and prompt BAT Observations](#)
- 12983 [Optical fluxes of candidate neutrino blazar PKS 1502+106](#)
- 12981 [ASKAP observations of blazars possibly associated with neutrino events IC190730A and IC190704A](#)
- 12974 [Optical follow-up of IceCube-190730A with ZTF](#)
- 12971 [IceCube-190730A: MASTER alert observations and analysis](#)
- 12967 [IceCube-190730A an astrophysical neutrino candidate in spatial coincidence with FSRQ PKS 1502+106](#)
- 12926 [VLA observations reveal increasing brightness of 1WHSP J104516.2+275133, a potential source of IC190704A](#)

γ -ray

radio

PKS 1502+106



300 TeV
neutrino
produced

target
moves
through
the jet:
blocks
photons

THE ICECUBE COLLABORATION

