

Test of Lepton Flavor Universality using $B_d \rightarrow D^{*-} \tau \nu$ decays at LHCb

GDR - Intensity frontiere 2022

G. BENANE[†], C. CHEN[†], B. FANG, D. FAZZINI, D. GERSTEL, O. LEROY[†],
A. MORRIS, P. K. RESMI, A. ROMERO VIDAL, G. WORMSER

[†] Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France
November 3rd 2022

Introduction

$R(D^*)$ hadronic analysis (Run 2)


Results

Conclusions

Introduction

Lepton Flavor Universality

The weak interaction in the Standard Model treats identically the three charged leptons: e , μ , and τ except for their different masses.




This property is referred as **Lepton Flavor Universality (LFU)**.

To test the LFU hypothesis we measure

$$R(D^*) \equiv \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)}$$

$R(D^{(*)})$ measurements \rightarrow charged flavor changing current $b \rightarrow c \ell \nu_\ell$:




Previous measurements of $R(D^*)$:

- BaBar (2012)
 - $\frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \ell \nu_\mu)}$ with $\ell = \mu, e$
- Belle (2015 and 2017)
 - Hadronic and leptonic τ
 - One-prong hadronic $\tau \rightarrow \pi \nu_\tau$ and $\tau \rightarrow \rho \nu_\tau$
- LHCb (2015 and 2018)
 - Muonic $\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau$
 - 3-prong hadronic τ (Run 1)

$R(D^*)$ hadronic analysis (Run 2)

Signal & Normalisation mode


$$B^0 \rightarrow D^{*-} \tau^+ \nu_\tau \text{ and } \tau^+ \rightarrow 3\pi^\pm (\pi^0) \bar{\nu}_\tau$$



Signal mode

- Same final states in signal and normalisation modes
- Signal mode partially reconstructed (missing neutrino $\bar{\nu}_\tau$)

$$B^0 \rightarrow D^{*-} 3\pi^\pm$$




Normalisation mode

- Normalisation mode fully reconstructed
- Helps to cancel out systematic uncertainties


Signal & Normalisation mode

$$B^0 \rightarrow D^{*-} \tau^+ \nu_\tau \text{ and } \tau^+ \rightarrow 3\pi^\pm (\pi^0) \bar{\nu}_\tau$$



Signal mode

$$B^0 \rightarrow D^{*-} 3\pi^\pm$$



Normalisation mode

- Same final states in signal and normalisation modes
- Signal mode partially reconstructed (missing neutrino $\bar{\nu}_\tau$)
- Normalisation mode fully reconstructed
- Helps to cancel out systematic uncertainties

Most relevant backgrounds are: Prompt decay: $B^0 \rightarrow D^* 3\pi^\pm$, double-charm decay: $B^0 \rightarrow D^* DX$ where $D \rightarrow 3\pi^\pm$, combinatorial backgrounds.

$R(D^*)$ measurement

$$R(D^*) = \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)}$$

$R(D^*)$ measurement

$$R(D^*) = \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)} = \underbrace{\frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \pi^+ \pi^- \pi^+)}}_{\mathcal{K}(D^*)} \times \underbrace{\frac{\mathcal{B}(B^0 \rightarrow D^{*-} \pi^+ \pi^- \pi^+)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)}}_{\text{External branching fractions}}$$

$R(D^*)$ measurement

$$R(D^*) = \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)} = \underbrace{\frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \pi^+ \pi^- \pi^+)}}_{\mathcal{K}(D^*)} \times \underbrace{\frac{\mathcal{B}(B^0 \rightarrow D^{*-} \pi^+ \pi^- \pi^+)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)}}_{\text{External branching fractions}}$$

We measure:

$$\mathcal{K}(D^*) \equiv \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} 3\pi^\pm)} = \frac{N_{\text{sig}}}{N_{\text{norm}}} \frac{\epsilon_{\text{norm}}}{\epsilon_{\text{sig}}} \frac{1}{\mathcal{B}(\tau^+ \rightarrow 3\pi^\pm \bar{\nu}_\tau) + \mathcal{B}(\tau^+ \rightarrow 3\pi^\pm (\pi^0) \bar{\nu}_\tau)}$$

$R(D^*)$ measurement

$$R(D^*) = \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)} = \underbrace{\frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \pi^+ \pi^- \pi^+)}}_{\mathcal{K}(D^*)} \times \underbrace{\frac{\mathcal{B}(B^0 \rightarrow D^{*-} \pi^+ \pi^- \pi^+)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)}}_{\text{External branching fractions}}$$

We measure:

$$\mathcal{K}(D^*) \equiv \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} 3\pi^\pm)} = \frac{N_{\text{sig}}}{N_{\text{norm}}} \frac{\epsilon_{\text{norm}}}{\epsilon_{\text{sig}}} \frac{1}{\mathcal{B}(\tau^+ \rightarrow 3\pi^\pm \bar{\nu}_\tau) + \mathcal{B}(\tau^+ \rightarrow 3\pi^\pm (\pi^0) \bar{\nu}_\tau)}$$

- N_{sig} from a 3D binned template fit:
 - $q^2 = (p_B - p_{D^*})^2$ momentum transferred to the leptonic system (8 bins),
 - τ^+ lifetime t_τ (8 bins),
 - Anti- D_s^+ BDT (6 bins).

$R(D^*)$ measurement

$$R(D^*) = \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)} = \underbrace{\frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \pi^+ \pi^- \pi^+)}}_{\mathcal{K}(D^*)} \times \underbrace{\frac{\mathcal{B}(B^0 \rightarrow D^{*-} \pi^+ \pi^- \pi^+)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)}}_{\text{External branching fractions}}$$

We measure:

$$\mathcal{K}(D^*) \equiv \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} 3\pi^\pm)} = \frac{N_{\text{sig}}}{N_{\text{norm}}} \frac{\epsilon_{\text{norm}}}{\epsilon_{\text{sig}}} \frac{1}{\mathcal{B}(\tau^+ \rightarrow 3\pi^\pm \bar{\nu}_\tau) + \mathcal{B}(\tau^+ \rightarrow 3\pi^\pm (\pi^0) \bar{\nu}_\tau)}$$

- N_{sig} from a 3D binned template fit:
 - $q^2 = (p_B - p_{D^*})^2$ momentum transferred to the leptonic system (8 bins),
 - τ^+ lifetime t_τ (8 bins),
 - Anti- D_s^+ BDT (6 bins).
- N_{norm} from an unbinned fit to $m(D^* 3\pi^\pm)$

R(D*) measurement

$$R(D^*) = \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)} = \underbrace{\frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \pi^+ \pi^- \pi^+)}}_{\mathcal{K}(D^*)} \times \underbrace{\frac{\mathcal{B}(B^0 \rightarrow D^{*-} \pi^+ \pi^- \pi^+)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)}}_{\text{External branching fractions}}$$

We measure:

$$\mathcal{K}(D^*) \equiv \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} 3\pi^\pm)} = \frac{N_{\text{sig}}}{N_{\text{norm}}} \frac{\epsilon_{\text{norm}}}{\epsilon_{\text{sig}}} \frac{1}{\mathcal{B}(\tau^+ \rightarrow 3\pi^\pm \bar{\nu}_\tau) + \mathcal{B}(\tau^+ \rightarrow 3\pi^\pm (\pi^0) \bar{\nu}_\tau)}$$



- N_{sig} from a 3D binned template fit:
 - $q^2 = (p_B - p_{D^*})^2$ momentum transferred to the leptonic system (8 bins),
 - τ^+ lifetime t_τ (8 bins),
 - Anti- D_s^+ BDT (6 bins).
- N_{norm} from an unbinned fit to $m(D^* 3\pi^\pm)$
- Efficiencies ϵ_{sig} and ϵ_{norm} extracted from MC samples

Selection

- Initial cuts (preselection and common cuts)
- Apply four BDTs (next slides):
 - $3\pi^\pm$ vertex detachment BDT
 - Anti-combinatorial background BDT
 - Charged isolation BDT
 - Anti- $B^0 \rightarrow D^{*-} D_s^+$ X BDT
- Remaining cuts:
 - Signal and normalisation modes

$3\pi^\pm$ vertex detachment BDT



- Remove 'prompt' background



Δz /uncertainty distribution of the simulated signal (red), double charm background (black) and prompt background (grey), after the initial cuts. A cut at 2σ is shown.

3π[±] vertex detachment BDT



- Remove 'prompt' background
- Training samples:
 - signal: MC $B^0 \rightarrow D^{*-} \tau^+ \nu_\tau$
 - background: $b\bar{b} \rightarrow D^{*-} 3\pi^\pm$ MC where 3π[±] don't come from the τ⁺ nor any D-meson



- Signal efficiency: 70%; background rejection: 90% at BDT > 0.2

Anti-combinatorial background BDT



- Remove D^{*-} and $3\pi^\pm$ from different hadrons
- Training samples:
 - signal: $B^0 \rightarrow D^{*-} \tau^+ \nu_\tau$ MC
 - background: wrong-sign data



Signal efficiency: 85% & background rejection: 70% at BDT > 0

Charged isolation BDT

- Remove events with extra charged tracks associated with signal ones
- Training samples: $b\bar{b} \rightarrow D^{*\mp} 3\pi^{\pm}$ MC
 - signal: **without** extra tracks
 - background: **with** extra tracks





Signal efficiency: 80% and background rejection 77% at $BDT > 0$

Anti- $B^0 \rightarrow D^{*-} D_s^+ X$ BDT

→ This BDT is used in final fit

- Distinguish $\tau^+ \rightarrow 3\pi^-$ from signal vs. $D_s^+ \rightarrow 3\pi^\pm X$ from $B^0 \rightarrow D^{*-} D_s^+ X$
- Training samples:
 - signal: MC $B^0 \rightarrow D^{*-} \tau^+ \nu_\tau$
 - background: MC-truth-matched $B^0 \rightarrow D^{*-} D_s^+ X$




Signal efficiency: 99.7%; background rejection: 31% at BDT > -0.2

Control Samples

D_s decay model


A simultaneous maximum likelihood binned fit to: $\min m(\pi^+\pi^-)$, $\max m(\pi^+\pi^-)$, $m(\pi^+\pi^+)$ and $m(3\pi^\pm)$



Control Samples

$B \rightarrow DD_s^+(X)$ control mode

- Fit to $m(D^* 3\pi^\pm)$



- We extract fractions that are used as constraints in the signal yield fit

Results

Signal fit - PDF

The signal yield is determined from a 3-dimensional maximum likelihood binned fit to q^2 (8 bins), decay time of the τ^+ -candidate (8 bins), t_τ , and the anti- D_s^+ BDT (6 bins).

- The total probability density function is:

$$\begin{aligned}
 \mathcal{P}_{\text{total}}(q^2, t_\tau, \text{BDT}) = & 1/N_{\text{total}} \times \{ N_{\text{sig}} [f_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \bar{\nu}_\tau} \mathcal{P}_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \bar{\nu}_\tau} + (1 - f_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \bar{\nu}_\tau}) \mathcal{P}_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \pi^0 \bar{\nu}_\tau} \\
 & + f_{D^{*+} \tau \nu} \mathcal{P}_{B \rightarrow D^{*+} \tau^+ \nu_\tau}] + N_{D^0}^{\text{same}} [\mathcal{P}_{B \rightarrow D^{*+} D^0 X \text{ SV}} + f_{D^0}^{v_1 - v_2} \mathcal{P}_{B \rightarrow D^{*+} D^0 X \text{ DV}}] \\
 & + N_{D_s^+} / k \times [\mathcal{P}_{B^0 \rightarrow D^{*+} D_s^+} + f_{D_s^+} \mathcal{P}_{B^0 \rightarrow D^{*+} D_s^+} + f_{D_s^0}^{*+} \mathcal{P}_{B^0 \rightarrow D^{*+} D_s^0} \\
 & + f_{D_{s1}^+} \mathcal{P}_{B^0 \rightarrow D^{*+} D_{s1}^+} + f_{D^{*+} D_s X} \mathcal{P}_{B \rightarrow D^{*+} D_s^+ X} + f_{B_s \rightarrow D^+ D_s^+ X} \mathcal{P}_{B_s^0 \rightarrow D^{*+} D_s^+ X}] \\
 & + N_{D_s^+} f_{D^+} \mathcal{P}_{B \rightarrow D^{*+} D^+ X} + N_{B \rightarrow D^{*+} - 3\pi^\pm X} \mathcal{P}_{B \rightarrow D^{*+} - 3\pi^\pm X} \\
 & + N_{B_1 - B_2} \mathcal{P}_{\text{combinatoric } B} + N_{\text{fake } D^0} \mathcal{P}_{\text{combinatoric } D^0} + N_{\text{fake } D^*} \mathcal{P}_{\text{combinatoric } D^{*-}} \}
 \end{aligned}$$

- 16 templates: 13 templates from MC , 3 templates from data
- 4 free parameters , 6 gaussian constrained parameters and 6 fixed parameters

Signal fit - PDF

The signal yield is determined from a 3-dimensional maximum likelihood binned fit to q^2 (8 bins), decay time of the τ^+ -candidate (8 bins), t_τ , and the anti- D_s^+ BDT (6 bins).

- The total probability density function is:

$$\begin{aligned}
 \mathcal{P}_{\text{total}}(q^2, t_\tau, \text{BDT}) = & 1/N_{\text{total}} \times \{ N_{\text{sig}} [f_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \bar{\nu}_\tau} \mathcal{P}_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \bar{\nu}_\tau} + (1 - f_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \bar{\nu}_\tau}) \mathcal{P}_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \pi^0 \bar{\nu}_\tau} \\
 & + f_{D^{*+} \tau \nu} \mathcal{P}_{B \rightarrow D^{*+} \tau + \nu_\tau}] + N_{D^0}^{\text{same}} [\mathcal{P}_{B \rightarrow D^{*-} D^0 X \text{ SV}} + f_{D^0}^{v_1 - v_2} \mathcal{P}_{B \rightarrow D^{*-} D^0 X \text{ DV}}] \\
 & + N_{D_s^+} / k \times [\mathcal{P}_{B^0 \rightarrow D^{*-} D_s^{*+}} + f_{D_s^+} \mathcal{P}_{B^0 \rightarrow D^{*-} D_s^+} + f_{D_{s0}^{*+}} \mathcal{P}_{B^0 \rightarrow D^{*-} D_{s0}^{*+}} \\
 & + f_{D_{s1}^{*+}} \mathcal{P}_{B^0 \rightarrow D^{*-} D_{s1}^{*+}} + f_{D^{*+} D_s X} \mathcal{P}_{B \rightarrow D^{*+} D_s^+ X} + f_{B_s \rightarrow D^{*+} D_s^+ X} \mathcal{P}_{B_s^0 \rightarrow D^{*-} D_s^+ X}] \\
 & + N_{D_s^+} f_{D^+} \mathcal{P}_{B \rightarrow D^{*-} D^+ X} + N_{B \rightarrow D^{*-} 3\pi^\pm X} \mathcal{P}_{B \rightarrow D^{*-} 3\pi^\pm X} \\
 & + N_{B_1 - B_2} \mathcal{P}_{\text{combinatoric } B} + N_{\text{fake } D^0} \mathcal{P}_{\text{combinatoric } D^0} + N_{\text{fake } D^*} \mathcal{P}_{\text{combinatoric } D^{*-}} \}
 \end{aligned}$$

- 16 templates: 13 templates from MC, 3 templates from data
- 4 free parameters, 6 gaussian constrained parameters and 6 fixed parameters

Signal fit - PDF

The signal yield is determined from a 3-dimensional maximum likelihood binned fit to q^2 (8 bins), decay time of the τ^+ -candidate (8 bins), t_τ , and the anti- D_s^+ BDT (6 bins).


- The total probability density function is:

$$\begin{aligned}
 \mathcal{P}_{\text{total}}(q^2, t_\tau, \text{BDT}) = & 1/N_{\text{total}} \times \{ N_{\text{sig}} [f_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \bar{\nu}_\tau} \mathcal{P}_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \bar{\nu}_\tau} + (1 - f_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \bar{\nu}_\tau}) \mathcal{P}_{\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \pi^0 \bar{\nu}_\tau} \\
 & + f_{D^{*+} \tau \nu} \mathcal{P}_{B \rightarrow D^{*+} \tau \nu}] + N_{D^0}^{\text{same}} [\mathcal{P}_{B \rightarrow D^{*-} D^0 X \text{ SV}} + f_{D^0}^{v_1 - v_2} \mathcal{P}_{B \rightarrow D^{*-} D^0 X \text{ DV}}] \\
 & + N_{D_s^+} / k \times [\mathcal{P}_{B^0 \rightarrow D^{*-} D_s^+} + f_{D_s^+} \mathcal{P}_{B^0 \rightarrow D^{*-} D_s^+} + f_{D_{s0}^{*+}} \mathcal{P}_{B^0 \rightarrow D^{*-} D_{s0}^{*+}} \\
 & + f_{D_{s1}^{*+}} \mathcal{P}_{B^0 \rightarrow D^{*-} D_{s1}^{*+}} + f_{D^{*+} D_s X} \mathcal{P}_{B \rightarrow D^{*+} D_s^+ X} + f_{B_s \rightarrow D^+ D_s^+ X} \mathcal{P}_{B_s^0 \rightarrow D^{*-} D_s^+ X}] \\
 & + N_{D_s^+} f_{D^+} \mathcal{P}_{B \rightarrow D^{*-} D^+ X} + N_{B \rightarrow D^{*-} 3\pi^\pm X} \mathcal{P}_{B \rightarrow D^{*-} 3\pi^\pm X} \\
 & + N_{B_1 - B_2} \mathcal{P}_{\text{combinatoric } B} + N_{\text{fake } D^0} \mathcal{P}_{\text{combinatoric } D^0} + N_{\text{fake } D^*} \mathcal{P}_{\text{combinatoric } D^{*-}} \}
 \end{aligned}$$

- 16 templates: 13 templates from MC , 3 templates from data
- 4 free parameters , 6 gaussian constrained parameters and 6 fixed parameters


Signal fit results

The signal yield is determined from a 3-dimensional maximum likelihood binned fit to q^2 (8 bins), decay time of the τ^+ -candidate (8 bins), t_τ , and the anti- D_s^+ BDT (6 bins).




Signal fit projection on q^2 (left), τ lifetime (middle) and the Anti- D_s BDT (right)

Run 1 and Run 2



LHCb cumulative integrated luminosity from 2010 till 2018

Year	\mathcal{L} (fb^{-1})	E (TeV)	Trigger efficiency
2011	1.11	3.5	
2012	2.08	4	
2015	0.33	6.5	increased by a
2016	1.67	6.5	factor of
2017	1.71	6.5	15%
2018	2.19	6.5	




Mass distribution of B_d for different datasets after applying normalisation cuts

- Run1 (2011+2012):** $R(D^*) = 0.283 \pm \underbrace{0.019}_{6.7\% \text{ stat.}} \pm \underbrace{0.029}_{10.3\% \text{ syst.}}$ (PRL.120.171802)
- Run2p1 (2015+2016):** Blinded $R(D^*)$ (Resmi P.K et al. LHCb internal review)
- Run2p2 (2017+2018):** Statistics increased by a factor > 2 w.r.t. 2016 (stat. uncertainty $\sim 3\%$)


LFU Anomalies: $R(D)$ vs $R(D^*)$ plot

$$R(D) \equiv \frac{\mathcal{B}(B^0 \rightarrow D^- \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^- \mu^+ \nu_\mu)} \quad \text{and} \quad R(D^*) \equiv \frac{\mathcal{B}(B^0 \rightarrow D^{*-} \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^{*-} \mu^+ \nu_\mu)}$$

3 experiments, 6 measurements, different analysis techniques



[Heavy Flavor Averaging Group, HFLAV]



All the measurements lie above the SM expectation



The current world-average measured $R(D)$ and $R(D^*)$ are 3.2σ away from the SM

Other R ratios


$$R(D) \equiv \frac{\mathcal{B}(B^0 \rightarrow D^- \tau^+ \nu_\tau)}{\mathcal{B}(B^0 \rightarrow D^- \mu^+ \nu_\mu)}$$

$$R(J/\psi) = \frac{\mathcal{B}(B_c^+ \rightarrow J/\psi \tau^+ \nu_\tau)}{\mathcal{B}(B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu)}$$

$$R(\Lambda_c^+) = \frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \tau \nu_\tau)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \mu \nu_\mu)}$$




[arxiv:1711.05623]



[arxiv:2201.03497]

[Heavy Flavor Averaging Group, HFLAV]

New Physics behind LFU?



Contributions for B to D^* semileptonic decay: Left: SM, Middle: Charged Higgs, Right: Leptoquark.

There are three typical candidates to account for the and $R(D)$ anomalies:

- Leptoquarks PRL 116, 081801 and PRD 94, 115021
- Two-Higgs-doublet models PRL 116, 081801
- Heavy vector bosons, e.g. W' JHEP 07 (2015) 142 1506.01705

Conclusions

Summary and Prospects

Summary

- $R(D^*)$ is an important analysis that may shed light on the intriguing Lepton Flavor Anomaly
- Complicated analysis because of many backgrounds due to p-p collisions and partially reconstructed signal
- Analysis of $R(D^*)$ hadronic for Run 2 part 1 (2015-2016) in final step of internal review

Prospects

- Complete Run 2 part 2 (2017-2018) analysis
- Use the full dataset Run1 + Run2 (2011-2018) to determine $R(D^*)$
 - Expected statistical uncertainty of the order of 3%
(while: 6.7% for run1 only and for 5% run1+run2p1)
- $R(D)$ and simultaneous measurements $R(D)$ and $R(D^*)$
- Angular analysis study of $B_d \rightarrow D^* \tau \nu_\tau$ decay
- Study other B-mesons

Thank you!

Backups

$R(D^*)$ hadronic ($\tau \rightarrow 3\pi\nu_\tau$) Systematics (run1) [PRL 120, 171802 2018]

$$R(D^*) = 0.280 \pm 0.018(\text{stat}) \pm 0.026(\text{syst}) \pm 0.013(\text{ext})$$

Breakdown of relative uncertainties:

Source	$\frac{\delta R(D^{*-})}{R(D^{*-})}$ [%]
Simulated sample size	4.7
Empty bins in templates	1.3
Signal decay model	1.8
$D_s^{*-} \tau \nu$ and $D_s^{*+} \tau \nu$ feed-downs	2.7
$D_s^+ \rightarrow 3\pi X$ decay model	2.5
$B \rightarrow D^{*-} D_s^+ X$, $D^{*-} D^+ X$, $D^{*-} D^0 X$ backgrounds	3.9
Combinatorial background	0.7
$B \rightarrow D^{*-} 3\pi X$ background	2.8
Efficiency ratio	3.9
Normalization channel efficiency (modeling of $B^0 \rightarrow D^{*-} 3\pi$)	2.0
Total systematic uncertainty	9.1
Statistical uncertainty	6.5
Total uncertainty	11.9

Will be improved in the next iteration of the analysis

$R(D^*)$ hadronic ($\tau \rightarrow 3\pi\nu_\tau$) Systematics [PRL 120, 171802 2018]

$$R(D^*) = 0.280 \pm 0.018(\text{stat}) \pm 0.026(\text{syst}) \pm 0.013(\text{ext})$$

Breakdown of relative uncertainties:

Source	$\frac{\delta R(D^{*-})}{R(D^{*-})}$ [%]	Future
Simulated sample size	4.7	Produce more MC !
Empty bins in templates	1.3	
Signal decay model	1.8	
$D^{**} \tau \nu$ and $D_s^{**} \tau \nu$ feed-downs	2.7	Measure $R(D_1(2420)^0)$
$D_s^+ \rightarrow 3\pi X$ decay model	2.5	BESIII
$B \rightarrow D^{*-} D_s^+ X, D^{*-} D^+ X, D^{*-} D^0 X$ bkgs	3.9	Improves with stat
Combinatorial background	0.7	
$B \rightarrow D^{*-} 3\pi X$ background	2.8	Kill with $ z\tau - zD > 5\sigma$
Efficiency ratio	3.9	Improves with stat
Normalization channel efficiency (modeling of $B^0 \rightarrow D^{*-} 3\pi$)	2.0	
Total systematic uncertainty	9.1	
Statistical uncertainty	6.5	
Total uncertainty	11.9	

$R(D^*)$ hadronic ($\tau \rightarrow 3\pi\nu_\tau$) [PRD 97,072013 2018]

List of the individual systematic uncertainties for $R(D^*)$:

Contribution	Value in %
$\mathcal{B}(\tau^+ \rightarrow 3\pi\bar{\nu}_\tau) / \mathcal{B}(\tau^+ \rightarrow 3\pi(\pi^0)\bar{\nu}_\tau)$	0.7
Form factors (template shapes)	0.7
Form factors (efficiency)	1.0
τ polarization effects	0.4
Other τ decays	1.0
$B \rightarrow D^{*+} \tau^+ \nu_\tau$	2.3
$B_S^0 \rightarrow D_S^{*+} \tau^+ \nu_\tau$ feed-down	1.5
$D_S^+ \rightarrow 3\pi X$ decay model	2.5
D_S^+ , D^0 and D^+ template shape	2.9
$B \rightarrow D^{*-} D_S^+(X)$ and $B \rightarrow D^{*-} D^0(X)$ decay model	2.6
$D^{*-} 3\pi X$ from B decays	2.8
Combinatorial background (shape + normalization)	0.7
Bias due to empty bins in templates	1.3
Size of simulation samples	4.1
Trigger acceptance	1.2
Trigger efficiency	1.0
Online selection	2.0
Offline selection	2.0
Charged-isolation algorithm	1.0
Particle identification	1.3
Normalization channel	1.0
Signal efficiencies (size of simulation samples)	1.7
Normalization channel efficiency (size of simulation samples)	1.6
Normalization channel efficiency (modeling of $B^0 \rightarrow D^{*-} 3\pi$)	2.0
Total uncertainty	9.1

Generated MC samples

Decay descr.	EventType	Generated evts [M]	Filtered evts [M]
$B^0 \rightarrow D^{*-} \tau^+ (\rightarrow 3\pi^\pm \bar{\nu}_\tau) \nu_\tau$	11160001	93.0	0.4
$B^0 \rightarrow D^{*-} \tau^+ (\rightarrow 3\pi^\pm \pi^0 \bar{\nu}_\tau) \nu_\tau$	11563020	90.0	0.3
$B^0 \rightarrow D^{*-} 3\pi^\pm$	11266018	363.0	1.0
$B^0 \rightarrow D^{**} \tau^+ (\rightarrow 3\pi^\pm \bar{\nu}_\tau) \nu_\tau$	11566431	7.0	0.09
$B_s^0 \rightarrow D^{*-} D_s^+ X$	13996612	50.0	0.4
$B^+ \rightarrow D^{*0} D_s^+ X$	12997613	354.0	4.0
$B^0 \rightarrow D^{*-} D_s^+ X$	11896612	692.0	8.0
$B^0 \rightarrow D^{*-} D_s^+ X$	11996413	42.0	0.4
$b\bar{b} \rightarrow D^{*-} 3\pi^\pm X$	27163970	8202.0	16.0
$b\bar{b} \rightarrow D^{*-} D^{\{0,+ \}} X$	27163971	427.0	2.0
Total	-	10320.0	32.0

$R(D^*)$ hadronic kinematics

Two-fold ambiguities in determining τ momentum:

$$|\vec{p}_\tau| = \frac{(m_{3\pi}^2 + m_\tau^2)|\vec{p}_{3\pi}| \cos \theta_{\tau,3\pi} \pm E_{3\pi} \sqrt{(m_\tau^2 - m_{3\pi}^2)^2 - 4m_\tau^2 |\vec{p}_{3\pi}|^2 \sin^2 \theta_{\tau,3\pi}}}{2(E_{3\pi}^2 - |\vec{p}_{3\pi}|^2 \cos^2 \theta_{\tau,3\pi})},$$

where $\theta_{\tau,3\pi}$ is the angle between the 3π system three-momentum and the τ line of flight.

Approximation: take the maximum allowed angle

$$\theta_{\tau,3\pi} \approx \theta_{\tau,3\pi}^{\max} = \arcsin \left(\frac{m_\tau^2 - m_{3\pi}^2}{2m_\tau |\vec{p}_{3\pi}|} \right),$$

The B^0 momentum is obtained similarly:

$$|\vec{p}_{B^0}| = \frac{(m_Y^2 + m_{B^0}^2)|\vec{p}_Y| \cos \theta_{B^0,Y} \pm E_Y \sqrt{(m_{B^0}^2 - m_Y^2)^2 - 4m_{B^0}^2 |\vec{p}_Y|^2 \sin^2 \theta_{B^0,Y}}}{2(E_Y^2 - |\vec{p}_Y|^2 \cos^2 \theta_{B^0,Y})}$$

with

$$\theta_{B^0,Y}^{\max} = \arcsin \left(\frac{m_{B^0}^2 - m_Y^2}{2m_{B^0} |\vec{p}_Y|} \right),$$

where Y represents the $D^{*-} \tau^+$ system.

Properties of charged leptons

Particle	Mass (MeV/c ²)	Lifetime	Main decay modes
e^-	0.5109989461(31)	$>6.6 \times 10^{26}$ years	None
μ^-	105.6583745(24)	2.1969811(22) μ s	$e^- \bar{\nu}_e \nu_\mu$
τ^-	1776.86(12)	290.3(5) fs	$\pi^- \pi^0 \nu_\tau$ (25.5%) $e^- \bar{\nu}_e \nu_\tau$ (17.8%) $\mu^- \bar{\nu}_\mu \nu_\tau$ (17.39%) $\pi^- \nu_\tau$ (10.8%) $\pi^- \pi^+ \pi^- \nu_\tau$ (9.3%)

τ lepton Branching Ratios [PDG 2018]

Mode	\mathcal{BR} (%)
$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	25.49 ± 0.09
$\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau$	17.82 ± 0.04
$\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau$	17.39 ± 0.04
$\tau^- \rightarrow \pi^- \nu_\tau$	10.82 ± 0.05
$\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau$	9.31 ± 0.05
$\tau^- \rightarrow \pi^- \pi^+ \pi^- \pi^0 \nu_\tau$	4.62 ± 0.05

D^* branching ratios

Mode	BR
$D^*(2007)^0 \rightarrow D^0 \pi^0$	$(64.7 \pm 0.9)\%$
$D^*(2007)^0 \rightarrow D^0 \gamma$	$(35.3 \pm 0.9)\%$
$D^*(2010)^+ \rightarrow D^0 \pi^+$	$(67.7 \pm 0.5)\%$
$D^*(2010)^+ \rightarrow D^+ \pi^0$	$(30.7 \pm 0.5)\%$
$D^*(2010)^+ \rightarrow D^+ \gamma$	$(1.6 \pm 0.4)\%$

Particle	Mass (MeV/c ²)	Lifetime
D^+	1869.65 ± 0.05	(1.040 ± 0.007) ps
D^0	1864.83 ± 0.05	(0.4101 ± 0.0015) ps
D_s^+	1968.34 ± 0.07	(0.504 ± 0.004) ps
Λ_c^+	2286.46 ± 0.14	(0.200 ± 0.006) ps
$D^*(2007)^0$	2006.85 ± 0.05	-
$D^*(2010)^-$	2010.26 ± 26	-

Relevant branching ratios

Mode	\mathcal{BR}
$B^0 \rightarrow D^*(2010)^- D_s^+$	$(8.0 \pm 1.1) \times 10^{-3}$
$B^0 \rightarrow D^*(2010)^- D_s^{*+}$	$(1.77 \pm 0.14) \times 10^{-2}$
$B^0 \rightarrow D^*(2010)^- D^0 K^+$	$(2.47 \pm 0.10 \pm 0.18) \times 10^{-3}$
$B^0 \rightarrow D^*(2010)^- D^*(2007) K^+$	$(10.6 \pm 0.33 \pm 0.86) \times 10^{-3}$
$B^0 \rightarrow D^*(2010)^- \pi^+ \pi^+ \pi^- \pi^0$	$(1.67 \pm 0.27)\%$
$B^0 \rightarrow D^*(2010)^- 3\pi^+ \pi^+ 2\pi^-$	$(4.7 \pm 0.9)\%$
$B^0 \rightarrow D^*(2010)^- D_{s0}(2317)^+$	$(1.5 \pm 0.6)\%$
$B^0 \rightarrow D^*(2010)^- D_{sJ}(2457)^+$	$(9.3 \pm 2.2) \times 10^{-3}$
$B^0 \rightarrow D^*(2010)^- D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*0} K^+ + D^{*+} K^0$	$(5.0 \pm 1.4) \times 10^{-3}$

Generated events 2017 and 2018

Decay descr.	EventType	Generated evts [M]	Filtered evts [M]
$B^0 \rightarrow D^{*-} \tau^+ (\rightarrow 3\pi^\pm \bar{\nu}_\tau) \nu_\tau$	11160001	65.0	0.3
$B^0 \rightarrow D^{*-} \tau^+ (\rightarrow 3\pi^\pm \pi^0 \bar{\nu}_\tau) \nu_\tau$	11563020	60.0	0.2
$B^0 \rightarrow D^{*-} 3\pi^\pm$	11266018	269.0	1.0
$B^0 \rightarrow D^{**} \tau^+ (\rightarrow 3\pi^\pm \bar{\nu}_\tau) \nu_\tau$	11566431	6.0	0.1
$B_s^0 \rightarrow D^{*-} D_s^+ X$	13996612	35.0	0.3
$B^+ \rightarrow D^{*0} D_s^+ X$	12997613	242.0	3.0
$B^0 \rightarrow D^{*-} D_s^+ X$	11896612	472.0	6.0
$B^0 \rightarrow D^{**} D_s^+ X$	11996413	29.0	0.3
$b\bar{b} \rightarrow D^{*-} 3\pi^\pm X$	27163970	5688.0	13.0
$b\bar{b} \rightarrow D^{*-} D^{\{0,+ \}} X$	27163971	302.0	1.0
Total	-	7170.0	26.0

Decay descr.	EventType	Generated evts [M]	Filtered evts [M]
$B^0 \rightarrow D^{*-} \tau^+ (\rightarrow 3\pi^\pm \bar{\nu}_\tau) \nu_\tau$	11160001	93.0	0.4
$B^0 \rightarrow D^{*-} \tau^+ (\rightarrow 3\pi^\pm \pi^0 \bar{\nu}_\tau) \nu_\tau$	11563020	90.0	0.3
$B^0 \rightarrow D^{*-} 3\pi^\pm$	11266018	363.0	1.0
$B^0 \rightarrow D^{**} \tau^+ (\rightarrow 3\pi^\pm \bar{\nu}_\tau) \nu_\tau$	11566431	7.0	0.09
$B_s^0 \rightarrow D^{*-} D_s^+ X$	13996612	50.0	0.4
$B^+ \rightarrow D^{*0} D_s^+ X$	12997613	354.0	4.0
$B^0 \rightarrow D^{*-} D_s^+ X$	11896612	692.0	8.0
$B^0 \rightarrow D^{**} D_s^+ X$	11996413	42.0	0.4
$b\bar{b} \rightarrow D^{*-} 3\pi^\pm X$	27163970	8202.0	16.0
$b\bar{b} \rightarrow D^{*-} D^{\{0,+ \}} X$	27163971	427.0	2.0
Total	-	10320.0	32.0

Selection efficiencies

Cut	Absolute efficiencies			Cumulative efficiencies		
	$3\pi\mathcal{D}_\tau$	$3\pi\pi^0\mathcal{D}_\tau$		$3\pi\mathcal{D}_\tau$	$3\pi\pi^0\mathcal{D}_\tau$	
Initial selection						
L0	89.51	86.60	89.08	89.51	86.60	89.08
Hlt1	89.76	87.32	90.92	87.14	83.88	88.02
Hlt2	79.90	77.31	90.33	73.25	69.02	85.10
$PV(\bar{D}^0) = PV(\tau^+)$	69.76	65.73	79.94	69.76	65.73	79.94
totCandidates = 1	60.89	52.22	71.97	58.06	49.87	67.75
$ \text{vtx}_z(\tau^+) - \text{vtx}_z(PV) /\text{error} > 10$	71.66	66.59	78.60	57.01	48.29	62.64
nSPDHits < 450	72.24	67.78	83.97	56.37	47.56	61.99
Signal selection						
$m(D^{*-}) - m(K^- \pi^+) \in [143, 148] \text{ MeV}/c^2$	94.63	93.98	-	94.63	93.98	-
$m(K^- \pi^+) \in [1840, 1890] \text{ MeV}/c^2$	97.36	97.39	-	92.28	91.70	-
$m(3\pi) < 1825 \text{ MeV}/c^2$	98.24	98.77	-	90.73	90.68	-
$m(B^0) < 5100 \text{ MeV}/c^2$	99.29	99.03	-	90.46	90.27	-
$q^2 \in [0, 12] \text{ GeV}^2/c^4$	97.52	97.22	-	88.74	88.53	-
combinatorial BDTD > 0	80.37	76.71	-	74.72	71.89	-
$ \text{vtx}_z(\tau^+) - \text{vtx}_z(B^0) /\text{error} > 2$	99.81	99.78	-	74.72	71.89	-
isolation BDT > 0	87.85	83.86	-	67.42	62.41	-
anti D_s^+ BDT > -0.2	98.30	86.10	-	67.12	54.87	-
PID	76.23	78.86	-	-	-	-
Normalisation selection						
$ \text{vtx}_z(\bar{D}^0) - \text{vtx}_z(\tau^+) /\text{error} > 4$	-	-	94.30	-	-	94.30
$m(D^* 3\pi^\pm) \in [5150, 5400] \text{ MeV}$	-	-	97.87	-	-	93.32
$m(D^{*-}) - m(\bar{D}^0) \in [143, 148] \text{ MeV}$	-	-	94.97	-	-	89.04
combinatorial BDTD > 0	-	-	81.37	-	-	74.19
isolation BDT > 0	-	-	88.33	-	-	66.94
PID	-	-	73.96	-	-	-

Signal fit templates

The signal yield is determined from a 3-dimensional maximum likelihood binned fit to q^2 (8 bins), decay time of the τ^+ -candidate (8 bins), t_τ , and the anti- D_s^+ BDT (6 bins).

There are 16 templates, 13 of them come from MC and three from data. The latter ones are combinatorial B , D^0 and D^* events. The templates are grouped into the 12 following categories, due to similar shapes:

- $B^0 \rightarrow D^{*-} \tau^+ \nu_\tau$ – the signal; includes $\tau^+ \rightarrow 3\pi^- \bar{\nu}_\tau$ and $\tau^+ \rightarrow 3\pi^\pm \pi^0 \bar{\nu}_\tau$
- $B^0 \rightarrow D^{**} \tau^+ \nu_\tau$ – excited D^{*-} states
- $B^0 \rightarrow D^{*-} D_s^{+(*)}$ – includes $B^0 \rightarrow D^{*-} D_s^+$, $B^0 \rightarrow D^{*-} D_s^{+*}$, $B^0 \rightarrow D^{*-} D_{s0}^{*+}$ and $B^0 \rightarrow D^{*-} D_{s1}^+$
- $B \rightarrow D^{**} D_s^+ X$
- $B_s^0 \rightarrow D^{*-} D_s^+ X$
- $B \rightarrow D^{*-} D^+ X$
- $B \rightarrow D^{*-} 3\pi^\pm X$

Signal fit templates

The signal yield is determined from a 3-dimensional maximum likelihood binned fit to q^2 (8 bins), decay time of the τ^+ -candidate (8 bins), t_τ , and the anti- D_s^+ BDT (6 bins).

There are 16 templates, 13 of them come from MC and three from data. The latter ones are combinatorial B , D^0 and D^* events. The templates are grouped into the 12 following categories, due to similar shapes:

- $B \rightarrow D^{*-} D^0 X$ SV, 'Same Vertex' where all 3 pions come from D^0
- $B \rightarrow D^{*-} D^0 X$ DV, 'Different Vertices' where at least 1 of the 3 pions comes from the D^0 vertex and the other(s) from a different vertex, e.g. the slow pion from D^{*-} is reconstructed as coming from the D^0
- combinatorial B^0 whose template is made from the collision data with the $D^{*\pm}$ of the same sign as the $3\pi^\pm$ system (*i.e.* wrong sign data w.r.t. the signal)
- combinatorial D^0
- combinatorial D^{*-} but genuine D^0


NTuples production

- 2016 (real data + MC) dataset (reproduced as a cross-check)
- 2017-2018 (real data + MC) dataset (new production)

Year	2016	2017	2018
# MC event types	14	13	13
# MC events [M]	21.8	39.8	49.4
# data events [M]	7	7.2	9.1

In total, we generate $\mathcal{O}(100M)$ events.

Normalisation fit result



Fit results

Sweight fit result at the $D^*3\pi^\pm$ mass peak using 2018 dataset


year	N_{norm}	N_{norm} expected	N_{bkg}	N_{bkg} expected	$N_{\text{norm}}/N_{\text{bkg}}$
2016	$26\,434 \pm 190$	-	$1\,446 \pm 107$	-	18.28
2017	$31\,200 \pm 207$	27 067	$2\,002 \pm 117$	1 481	15.58
2018	$37\,137 \pm 225$	34 664	$2\,170 \pm 126$	1 896	17.11

B^0 and τ vertex error reweighting


- Main background Prompt decay where $3\pi^\pm$ system comes directly from the B^0 vertex.
- To suppress this background we require a separation between the B^0 and τ vertex

$$\Delta z = vt_{x_z}(\tau^+) - vt_{x_z}(B^0) \quad \sigma_{\Delta z} = \sqrt{vt_{x_{err\ z}}(\tau^+)^2 + vt_{x_{err\ z}}(B^0)^2}$$


- Difference between 'run2p1' and 'run2p2' MC: B^0 vertex error in the beam direction
 - Due to reconstruction algorithms (applied for data but not for MC)
 - Need to apply simultaneous gradient-boosted weight for the B^0 and τ vertex error



Signal mode



(a) No reweighting



(b) Reweighted MC

B^0 vertex_z error from sweigted data and $D^*3\pi^\pm$ MC 2018 sample