

Lattice QCD w/ a focus on B and D form factors, $R(D^*)$, and quark masses

Andrew Lytle (FNAL-MILC collaboration)
University of Illinois @ Urbana-Champaign

04.07.23
Beauty 2023
Clermont-Ferrand, France

Introduction

- The LHC era has produced/is producing an incredible wealth of data
 - ▶ New processes
 - ▶ Ever increasing levels of precision
- Stringent Standard Model tests request corresponding levels of theoretical precision
 - ▶ → Tests of the CKM matrix
 - ▶ Detailed comparisons with observed data (R -ratios, shape data, ...)

Outline


- Lattice QCD - broad overview
- Leptonic decays (briefly)
- Semileptonic decays
 - ▶ $B_{(s)}$ decays
 - ▶ $D_{(s)}$ decays
- Quark masses
- Summary & Conclusion

Lattice QCD - Broad overview

- From-first-principles approach to QCD/hadronic physics.
- Discretize the Euclidean time path integral \rightarrow finite lattice spacing, a .
- Calculate hadronic observables using Monte Carlo & supercomputers, take $a \rightarrow 0$.
- Works well for
 - ▶ Leptonic decays (two-point functions)
 - ▶ Semileptonic decays (three-point functions)and is *systematically improvable*.

Lattice QCD - Broad overview

Example: MILC lattice ensembles 1712.09262




- 6 lattice spacings, finest $a \approx 0.03$ fm
- Physical pion masses at all but finest spacing

Leptonic decays

Leptonic decays - I

- Relatively straightforward to compute, with meson 2-point functions.
- Allow extraction of CKM matrix elements via leptonic decay measurements.
 - ▶ $f_D \rightarrow |V_{cd}| = 0.2173(47)_{\text{exp}}(28)_{\text{em}}(7)_{\text{latt}}$
 - ▶ $f_{D_s} \rightarrow |V_{cs}|$ (sl decays now competitive)
 - ▶ $f_B \rightarrow |V_{ub}|$
- $f_{B_{(s)}}$ used for SM prediction of rare processes $B_{(s)} \rightarrow \mu^+ \mu^-$
- In all cases lattice QCD uncertainties well below experimental uncertainties

Leptonic decays - II



- Sub-percent precision obtained, results from different collaborations in good agreement.
- To improve substantially requires including QED and strong-isospin breaking
- Substantial progress including QED corrections to these processes (cf radiative decays)

$$B_{(s)} \rightarrow \mu^+ \mu^- \text{ and } B \rightarrow \ell \nu$$

$$B_{(s)} \rightarrow \mu^+ \mu^-$$

- Highly suppressed FCNC process can give important constraints on new physics
- $\mathcal{B}(B_s \rightarrow \mu\mu)_{\text{SM}} = 3.64(4) f_{B_s} (8)_{\text{CKM}} (7)_{\text{other}} \times 10^{-9}$
 $\mathcal{B}(B_s \rightarrow \mu\mu)_{\text{exp}} = 2.69(37) \times 10^{-9}$ ATLAS-CONF-2020-049
- $\mathcal{B}(B^0 \rightarrow \mu\mu)_{\text{SM}} = 1.00(1) f_B (2)_{\text{CKM}} (2)_{\text{other}} \times 10^{-10}$
 $\mathcal{B}(B^0 \rightarrow \mu\mu)_{\text{exp}} < 1.6 \times 10^{-10}$

B leptonic decays

- $B \rightarrow \tau \nu$ may reach 3-5% precision at Belle II
- $B \rightarrow \mu \nu$ may reach $\sim 7\%$ precision at Belle II 1808.10567


$B_{(s)}$ -meson semileptonic decays

Semileptonic decays - I

SL Decay processes critical inputs for heavy flavor studies.

Lattice predictions needed for:

- Extracting CKM matrix elements from expt'l measurements
- Pure SM predictions of R-ratios
- SM predictions $\frac{d\Gamma}{dq^2}$, etc.



Lattice calculations based on 2- & 3-point correlators give matrix elements $\rightarrow f_i(q^2)$

Semileptonic decays - II

Tree level decays:


- $B \rightarrow D^{(*)} \rightarrow |V_{cb}|$
- $B \rightarrow \pi \rightarrow |V_{ub}|$
- $D \rightarrow K \rightarrow |V_{cs}|$
- Baryon decays: $\rightarrow |V_{ub}|/|V_{cb}|$

Many different processes and results, can only highlight a few illustrative developments!


Loop level decays:

- Lattice form factors important for $B \rightarrow K\ell^+\ell^-$
- $B \rightarrow K^*\ell^+\ell^-$ requires recent theory developments in handling resonances/multi-hadron final states.

Semileptonic decays - III




$B \rightarrow D^*$ at zero recoil from LQCD







FNAL/MILC 1403.0635

- $n_f = 2 + 1$ MILC asqtad ensembles
- Clover b with Fermilab interpretation
- $h_{A_1}(1) = 0.906(4)(12)$





HPQCD 1711.11013

- $n_f = 2 + 1 + 1$ MILC HISQ ensembles
- NRQCD b quark
- $h_{A_1}(1) = 0.895(10)(24)$
- $h_{A_1}^s(1) = 0.883(12)(28)$





Figs. courtesy A. Vaquero



Figs. courtesy A. Vaquero

$B \rightarrow D^*$ comparisons



Figs. courtesy A. Vaquero

Lattice refs:


- JLQCD 2306.05657
- HPQCD 2304.03137
- FNAL/MILC 2105.14019

New set of measurements based on LHC Run 1 data.

- CLN: $|V_{cb}| = 41.4(6)_{\text{stat}}(9)_{\text{syst}}(12)_{\text{ext}} \times 10^{-3}$
- BGL: $|V_{cb}| = 42.3(8)_{\text{stat}}(9)_{\text{syst}}(12)_{\text{ext}} \times 10^{-3}$
- In this analysis, LQCD inputs improve statistical precision by 20% and 50% for CLN and BGL, respectively.

See also talk by Mirco Dorigo @LHCP2020
indico.cern.ch/event/856696/contributions/3742179/

New CKM constraints from LHCb. Based on baryon decays
1504.01568, and first msm't of $B_s \rightarrow K\mu\nu$ 2012.05143.




Extractions use lattice form factor calculations of **baryon decays**
($\Lambda_b \rightarrow \Lambda_c$ and $\Lambda_b \rightarrow p$ 1503.01421) and $B_s \rightarrow K$ and $B_s \rightarrow D_s$
form factors, respectively.

$R(B_c \rightarrow J/\psi)$

First lattice calculation of $B_c \rightarrow J/\psi$ 2007.06956
(All form factors across q^2 range)

- LHCb 1711.05623
 $R(J/\psi) = 0.71(17)_{\text{stat}}(18)_{\text{syst}}$
- w/in 2σ of theory range [0.25, 0.28]
- LQCD result: 0.2582(38)



FNAL-MILC all-HISQ semileptonic decays

Treatment of heavy quarks

Treatment of c and especially b quarks challenging in lattice simulation due to lattice artifacts which grow as $(am_h)^n$


- May use an effective theory framework to handle the b quark.
 - ▶ Fermilab method, RHQ, OK, NRQCD
 - ▶ Pros: Solves problem w/ am_h artifacts.
 - ▶ Cons: Requires matching, can still have ap artifacts.
- Also possible to use relativistic fermion provided a is sufficiently small $am_c \ll 1$, $am_b < 1$.
 - ▶ Use improved actions e.g. $\mathcal{O}(a^2) \rightarrow \mathcal{O}(\alpha_s a^2)$
 - ▶ Pros: Absolutely normalised current, straightforward continuum extrap.
 - ▶ Cons: Numerically expensive, extrapolate $m_h \rightarrow m_b$.

allhisq simulations


- Here we simulate *all* quarks with the HISQ action.
- Unified treatment for wide range of $B_{(s)}$ (and $D_{(s)}$) to pseudoscalar transitions
 - ▶ $B_{(s)} \rightarrow D_{(s)}$
 - ▶ $B_{(s)}/D_{(s)} \rightarrow K$
 - ▶ $B/D \rightarrow \pi$
- Ensembles with (HISQ) sea quarks down to physical at each lattice spacing.
- Enables correlated studies of ff *ratios*.

See 2022 Lattice proceeding for more details. 2301.09229

$B_s \rightarrow K$



$B_s \rightarrow D_s$





Preliminary results show

- Good q^2 coverage
- Good statistical precision
- Data at/beyond $M_{B(s)}$ \rightarrow interpolation in heavy mass m_h

$D_{(s)}$ -meson semileptonic decays

$|V_{cs}|$ from $D \rightarrow K$



New result from HPQCD 2104.09883. See also ETMC calculations 1706.03017, 1706.03657, 1803.04807.



$$|V_{cs}| = 0.9663(53)_{\text{latt}}(39)_{\text{exp}}(19)_{\eta_{\text{EW}}}(40)_{\text{EM}}$$



Semileptonic determination of $|V_{cs}|$ now has smaller errors than leptonic determinations.

$D \rightarrow \pi, K$



References:

- FNAL-MILC 2212.12648
- HPQCD 21 2104.09883
- ETMC 17 1706.03017




Semileptonic decays - Summary

- Thus far, successful collaboration with experiment in the LHC era
 - ▶ Theory predictions for new channels
 - ▶ Improved kinematic range
 - ▶ Improved precision
- Lattice errors roughly commensurate with experimental errors. In the next 5 years or so, these should continue to improve and lattice error may become sub-dominant.
- To go beyond this requires adding EM and strong isospin breaking effects.


Quark masses

Quark masses - I

[1606.02266]



[1312.4974]





HL-LHC estimates $\sim 3\%$ $hb\bar{b}$, evidence for charm CMS:2022cju

Est'd final ILC precision in $hb\bar{b}$ ($hc\bar{c}$) coupling: ~ 0.3 (0.7)%.

Quark masses on the lattice

- Quarks are not asymptotic (physical) states due to confinement – mass cannot be measured directly.
- Quark masses are scheme and scale dependent, $m_q^{\text{scheme}}(\mu)$.
- Generally will quote results $m_q^{\overline{\text{MS}}}(\mu_{\text{ref}})$.
- Lattice input quark masses are non-universal (depend on discretisation), but can be connected to quark masses defined in a continuum scheme.


Quark masses - II




- A variety of different methods, with different systematics, have been developed. See 2110.15090 for a recent review.
- FLAG averages 2111.09849 $N_f = 2 + 1 + 1$
 $m_b(m_b) = 4.203(11)$ GeV $m_c(3 \text{ GeV}) = 0.988(11)$

Quark masses - III


- Since CHARM 2015 - Highlights
 - ▶ Many new results: 13 since 2015, 4 since FLAG19.
 m_c : 6 \rightarrow 13, $n_f = 3$: 5 \rightarrow 9
 m_b : 4 \rightarrow 9, $n_f = 4$: 5 \rightarrow 13
 - ▶ From more collaborations \rightarrow differing discretizations/lattice artifacts under control.
 - ▶ New techniques with different sources of systematic uncertainty.
- QED corrections are small ($\lesssim 0.2\%$ for charm).
- Precision continues to improve.
- Complementary techniques - good agreement at sub-percent level.



Time moments of charmonium correlators
1408.4169



Heavy-light meson masses and HQET
1802.04248



Quark-level diagrams on the lattice
1805.06225



See also Schroedinger functional methods: 2101.02694 1802.05243

Summary

- We have looked at several places where lattice calculations impact precision SM tests
- Lattice QCD allows stringent comparisons of the Standard Model with experiment
 - ▶ Required for precision CKM studies
 - ▶ Rare processes and flavor anomalies
- Lattice theorists have kept pace with experimental advances at LHC, Belle II, and BES III
 - ▶ Calculations relevant for new msmt's
 - ▶ Improving precision

Thank you!

$D \rightarrow \pi, K$ rates from LQCD



References:

- FNAL-MILC 2212.12648
- HPQCD 21 2104.09883
- ETMC 17 1706.03017

$D_{(s)}$ decay — spectator quark dependence

