



Numerical investigation of screened scalar-tensor theories in space-based experiments

Hugo Lévy, Joël Bergé, Jean-Philippe Uzan

Colloque national Action Dark Energy 2023 - 7ème édition, Annecy




Scalar-Tensor theories

Scalar field in the gravitational sector


1. The dark sector [1]

- Dark Matter
- Dark Energy
- + inflation paradigm



2. A 'true' scalar field exists in nature

- Discovery of the Higgs boson in 2012
- ... ?



3. More fundamental theories

- String theory as an effective 4-dimensional theory [1]
- $f(R)$ -theories \equiv scalar-tensor [2]




[1] A. Joyce et al, arXiv:1407.0059

[2] J. Velásquez and L. Castañeda, arXiv:1808.05615

Scalar field in the gravitational sector


1. The dark sector [1]

- Dark Matter
- **Dark Energy**
- + inflation paradigm




2. A 'true' scalar field exists in nature

- Discovery of the Higgs boson in 2012
- ... ?



3. More fundamental theories

- String theory as an effective 4-dimensional theory [1]
- $f(R)$ -theories \equiv scalar-tensor [2]




[1] A. Joyce et al, arXiv:1407.0059

[2] J. Velásquez and L. Castañeda, arXiv:1808.05615


Laboratory Tests



Atom interferometry [3]




Eöt-Wash torsion pendulum [4]



Casimir effect [5]

Solar System Tests



© Y. Gominet/IMCCE/Nasa



MICROSCOPE [6]

Astrophysical Tests



Galaxy rotation curves [7]



Cluster lensing [8]



NASA/Goddard Space Flight Center


Scalar radiation in BNS [9]

Length scales


Laboratory Tests



Atom interferometry [3]




Eöt-Wash torsion pendulum [4]



Casimir effect [5]

Solar System Tests



© Y. Gominet/IMCCE/Nasa



MICROSCOPE [6]

Astrophysical Tests



Galaxy rotation curves [7]



Cluster lensing [8]



NASA/Goddard Space Flight Center

Scalar radiation in BNS [9]

Length scales

Scalar fields playing *hide-and-seek*

Review of the most popular screening mechanisms [2]		
Classification	Type of Equation	Rule of Thumb
Weak coupling <ul style="list-style-type: none"> Symmetron Damour-Polyakov 	$\square\phi = \frac{dV_{\text{eff}}}{d\phi}$	Occurs in regions of high Newtonian potential
Large mass <ul style="list-style-type: none"> Chameleon 		
Large inertia <ul style="list-style-type: none"> K-Mouflage Vainshtein 	$\square\phi + A_1\partial_\mu [(\partial\phi)^2\partial^\mu\phi] + A_2T = 0$	Occurs in regions where the gravitational acceleration is large
	$6\square\phi + B_1 [(\square\phi)^2 - (\partial_\mu\partial_\nu\phi)^2] = B_2T^\mu{}_\mu$	Occurs in regions where spatial curvature is large

Take-home messages:

- Different mechanisms to ‘screen’ scalar fields from local tests of gravity (i.e. recover GR at Solar System scales)
- At the equation level, screening \equiv non-linearity

Scalar fields playing *hide-and-seek*

Review of the most popular screening mechanisms [2]		
Classification	Type of Equation	Rule of Thumb
Weak coupling <ul style="list-style-type: none"> Symmetron Damour-Polyakov 	$\square\phi = \frac{dV_{\text{eff}}}{d\phi}$ Non-linear Klein-Gordon equation	Occurs in regions of high Newtonian potential
Large mass <ul style="list-style-type: none"> Chameleon 		
Large inertia <ul style="list-style-type: none"> K-Mouflage 	$\square\phi + A_1\partial_\mu [(\partial\phi)^2\partial^\mu\phi] + A_2T = 0$	Occurs in regions where the gravitational acceleration is large
<ul style="list-style-type: none"> Vainshtein 	$6\square\phi + B_1 [(\square\phi)^2 - (\partial_\mu\partial_\nu\phi)^2] = B_2T^\mu{}_\mu$	Occurs in regions where spatial curvature is large

Take-home messages:

- Different mechanisms to ‘screen’ scalar fields from local tests of gravity (i.e. recover GR at Solar System scales)
- At the equation level, screening \equiv non-linearity

Numerical considerations

[arXiv:2209.07226](https://arxiv.org/abs/2209.07226)

Chameleon field equation


Field equation (in the Newtonian limit)	5 th force
$\Delta\phi = \frac{\beta}{M_{Pl}}\rho - n\frac{\Lambda^{n+4}}{\phi^{n+1}}$ <p>Free parameters: β, n, Λ Mass distribution: $\rho = \rho(\mathbf{x}, t)$ Unknown: $\phi = \phi(\mathbf{x}, t)$</p>	$\vec{F}_\phi = -m\frac{\beta}{M_{Pl}}\nabla\phi$ <p>Point-mass follows geodesics of the Jordan frame metric \neq Einstein frame geodesics</p>

Chameleon field equation

Field equation (in the Newtonian limit)	5 th force
$\Delta\phi = \frac{\beta}{M_{Pl}}\rho - n\frac{\Lambda^{n+4}}{\phi^{n+1}}$ <p>Free parameters: β, n, Λ Mass distribution: $\rho = \rho(\mathbf{x}, t)$ Unknown: $\phi = \phi(\mathbf{x}, t)$</p>	$\vec{F}_\phi = -m\frac{\beta}{M_{Pl}}\nabla\phi$ <p>Point-mass follows geodesics of the Jordan frame metric \neq Einstein frame geodesics</p>

Geometry can be quite complex!

✓ Finite Element Method can deal with complex geometries



Chameleon field equation

Field equation (in the Newtonian limit)	5 th force
$\Delta\phi = \frac{\beta}{M_{Pl}} \rho - n \frac{\Lambda^{n+4}}{\phi^{n+1}}$ <p>Free parameters: β, n, Λ Mass distribution: $\rho = \rho(\mathbf{x}, t)$ Unknown: $\phi = \phi(\mathbf{x}, t)$</p>	$\vec{F}_\phi = -m \frac{\beta}{M_{Pl}} \nabla\phi$ <p>Point-mass follows geodesics of the Jordan frame metric \neq Einstein frame geodesics</p>

Boundary conditions?
$\begin{cases} \alpha\Delta\phi = \rho - \phi^{-(n+1)} \\ \phi(\mathbf{x}) \xrightarrow{\ \mathbf{x}\ \rightarrow +\infty} \phi_{\text{vac}} \end{cases}$ <p style="text-align: right;">“well-posed problem”</p>

Ω


$$\begin{cases} \alpha \Delta \phi = \rho - \phi^{-(n+1)} \\ \phi(\mathbf{x}) \xrightarrow{\|\mathbf{x}\| \rightarrow +\infty} \phi_{\text{vac}} \end{cases}$$

Distance from the origin


0

 $+\infty$


Ω



Ω



Not possible to mesh a domain of infinite spatial extent...



Not possible to mesh a domain of infinite spatial extent...



Let's *compactify** *space*!



***(i.e. apply a global coordinate transform that will map the whole plane to a bounded domain)**

$$\mathcal{T}: \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

For instance


$$\mathbf{x} = (x, y) \mapsto \frac{R_{\text{cut}}}{1 + \|\mathbf{x}\|} (x, y)$$







Penrose diagram









Γ

Apply $\phi = \phi_\infty$
on Γ

Γ

Apply $\Phi = \Phi_0$
on $\Gamma = \Gamma_0$



One idea among (many) others!

Caveat: Applying such coordinate transforms leads to unbounded coefficients in the resulting PDE (weight regularisation technique [arxiv:2209.07226](https://arxiv.org/abs/2209.07226))


Inspired by
Grosch and Orszag (1977)
Zenginoglu (2011)
Chernogorova et al. (2016)
Boulmezaoud (2005)

Programming (low level code)

Physics (high level code)

Study Scripts (`from femtoscope import ...`)

- Custom nonlinear solver with line-search
- Implementation of 3 techniques to handle asymptotic boundary conditions
- 1D, 2D and 3D Finite Element Method




femtoscope

- **Poisson Class**
(+ analytical & semi-analytical solutions available)
- **Chameleon Class**
(+ few analytical approximations)

Third-party Python librairies



Gmsh



meshio



sfepy



NumPy

Application to space geodesy* [arxiv:2310.03769](https://arxiv.org/abs/2310.03769)


All computations are performed with




**Space geodesy is a scientific discipline that involves precise measurements and analysis of the Earth's shape, gravitational field, and the dynamic behavior of its surface using satellite-based technologies.*

Motivations


- A satellite in orbit is subject to both **Newtonian attraction** and **fifth force**
- Strong **impact of the local landform** on the scalar field in the **screened regime**
- **Mountain** \equiv deviation from spherical symmetry + analogy with the '**lightning-rod effect**' in chameleon and symmetron models [10]
- Can a satellite flying over a mountain **distinguish** between **Newtonian gravity** and **chameleon gravity**?





Testing screened scalar-tensor models with mountains (?!)

GRACE-FO



GRACE-FO




GRACE-FO




**Laser Ranging Interferometry:
precision of few tenths of microns**

That's $\sim 10^{-10}$ km!!!


$t = t_0$



$$t = t_0$$




$$t = t_1$$



$$d(t_1) > d(t_0)$$


$$t = t_2$$




$$d(t_2) < d(t_1)$$

$$t = t_2$$


d (inter-satellite distance)





Time




$$t = t_2$$







Interpreting the measurement in the framework of Newtonian gravity gives us a gravity map






Passage of the satellites above the mountain






The anomaly is well within the range of GRACE-FO precision!

$$\mathcal{O}(1\text{cm}) \gg \mathcal{O}(10^{-5}\text{cm})$$



Sources of degeneracy




$$\rho(\mathbf{x}) = \rho_0 [1 + \delta_\rho(\mathbf{x})]$$

Questions

Is it possible to absorb the chameleon anomaly in

- a small uncertainty in the satellite's initial state vector δ_x ?
 - a slight variation in the {Earth + Mountain} density δ_ρ ?
- (in the framework of Newtonian gravity)

Sources of degeneracy



$$\rho(\mathbf{x}) = \rho_0[1 + \delta_\rho(\mathbf{x})]$$

Questions

Is it possible to absorb the chameleon anomaly in

- a small uncertainty in the satellite's initial state vector δ_x ?
- **a slight variation in the {Earth + Mountain} density δ_ρ ?**
(in the framework of Newtonian gravity)


Modified Gravity

(Newtonian + chameleon accelerations)

vs

Newtonian Gravity

with extra point-mass




Modified Gravity

(Newtonian + chameleon accelerations)

vs


Newtonian Gravity

with extra point-mass



● = 0,001% the mass of the initial mountain


Anomaly
 $\sim \mathcal{O}(10^{-4} \text{ cm})$




How to lift the degeneracy?

Idea:



Perform the experiment at
different altitudes




How to lift the degeneracy?



Tensions come in...




Tensions come in...



The greater the tension, the tighter the potential constraints on the modified gravity model at stake

Conclusion

- *femtoscope*: solve semi-linear elliptic PDE using the Finite Element Method on unbounded domains (general purpose code)



- Application to scalar-tensor theories of gravity:

Linear Poisson equation

$$\begin{cases} \Delta\Phi = 4\pi G\rho(\mathbf{x}) \\ \Phi(\mathbf{x}) \xrightarrow{\|\mathbf{x}\|\rightarrow+\infty} 0 \end{cases}$$


Nonlinear Klein-Gordon equation

$$\begin{cases} \Delta\phi = dV_{\text{eff}}/d\phi \\ \phi(\mathbf{x}) \xrightarrow{\|\mathbf{x}\|\rightarrow+\infty} \phi \end{cases}$$

$$\mathbf{a} \sim -\nabla(\Phi + \phi)$$

Gravitational Acceleration


- Application to space geodesy: focus on the GRACE-FO configuration
- Can we detect / put constraints on the chameleon model in this context?



- In comparison, lab experiments offer a more controlled environment



- Application to space geodesy: focus on the GRACE-FO configuration
- Can we detect / put constraints on the chameleon model in this context?



- In comparison, lab experiments offer a more controlled environment



Thanks for your attention!

References

- [1] A. Joyce et al, *Beyond the Cosmological Standard Model*, Physics Reports, Vol. 568, 22 March 2015, p. 1-98
- [2] J. Velásquez and L. Castañeda, *Equivalence between Scalar-Tensor theories and $f(R)$ -gravity: From the action to Cosmological Perturbations*, Journal of Physics Communications, May 2020
- [3] B. Elder et al, *Chameleon dark energy and atom interferometry*, Phys. Rev. D, August 2016
- [4] A. Upadhye, *Dark energy fifth forces in torsion pendulum experiments*, Phys. Rev. D, November 2012
- [5] A. Almasi et al, *Force sensor for chameleon and Casimir force experiments with parallel-plate configuration*, Phys. Rev. D, May 2015
- [6] P. Touboul et al (MICROSCOPE Collaboration),
Phys. Rev. Lett. 129, 121102 – Published 14 September 2022
- [7] V. Vikram et al, *Astrophysical tests of modified gravity: Stellar and gaseous rotation curves in dwarf galaxies*, Phys. Rev. D, May 2018
- [8] H. Wilcox et al, *Simulation tests of galaxy cluster constraints on chameleon gravity*, MNRAS, October 2016
- [9] A. Upadhye, J. H. Steffen, *Monopole radiation in modified gravity*, arXiv:1306.6113 [astro-ph.CO], June 2013
- [10] K. Jones-Smith and F. Ferrer, *Detecting Chameleon Dark Energy via an Electrostatic Analogy*, Phys. Rev. Lett. 108, 221101 – Published 29 May 2012

Backup Slides

Tensions come in...



FEM in a nutshell (example on Poisson's equation)

strong form

$$\left[\text{Find } u : \Omega \rightarrow \mathbb{R} \right. \quad \left. \begin{cases} -\Delta u = f & \text{in } \Omega \\ u = 0 & \text{on } \Gamma := \partial\Omega \end{cases} \right. \quad (1)$$

FEM in a nutshell (example on Poisson's equation)

strong form

$$\left[\begin{array}{l} \text{Find } u : \Omega \rightarrow \mathbb{R} \end{array} \right. \left\{ \begin{array}{l} -\Delta u = f \quad \text{in } \Omega \\ u = 0 \quad \text{on } \Gamma := \partial\Omega \end{array} \right. \quad (1)$$

FEM Recipe

1. Multiply Eq. (1) by a test function v
2. Integrate over Ω
3. Perform an integration by parts

FEM in a nutshell (example on Poisson's equation)

strong form

$$\left[\text{Find } u : \Omega \rightarrow \mathbb{R} \quad \begin{cases} -\Delta u = f & \text{in } \Omega \\ u = 0 & \text{on } \Gamma := \partial\Omega \end{cases} \right.$$

$$\text{Find } u \in V := H_0^1(\Omega) \quad \forall v \in V, \quad \underbrace{\int_{\Omega} \nabla u \cdot \nabla v \, dx}_{a(u,v)} = \underbrace{\int_{\Omega} f v \, dx}_{l(v)}$$

FEM in a nutshell (example on Poisson's equation)


strong form

$$\left[\text{Find } u : \Omega \rightarrow \mathbb{R} \quad \begin{cases} -\Delta u = f & \text{in } \Omega \\ u = 0 & \text{on } \Gamma := \partial\Omega \end{cases} \right.$$

$$\text{Find } u \in V := H_0^1(\Omega) \quad \forall v \in V, \quad \underbrace{\int_{\Omega} \nabla u \cdot \nabla v \, dx}_{a(u,v)} = \underbrace{\int_{\Omega} f v \, dx}_{l(v)}$$

4. Look for u in a finite-dimensional subspace $V^h \subset V$
(e.g. space of piecewise polynomial functions)

The landscape of basis functions



FEM in a nutshell (example on Poisson's equation)

strong form

$$\text{Find } u : \Omega \rightarrow \mathbb{R} \quad \begin{cases} -\Delta u = f & \text{in } \Omega \\ u = 0 & \text{on } \Gamma := \partial\Omega \end{cases}$$

continuous

weak formulation

$$\text{Find } u \in V := H_0^1(\Omega) \quad \forall v \in V, \underbrace{\int_{\Omega} \nabla u \cdot \nabla v \, dx}_{a(u,v)} = \underbrace{\int_{\Omega} f v \, dx}_{l(v)}$$

$$\text{Find } u_h \in V_h \subset V \quad \forall i \in [1, N] \quad a(u_h, v_i) = l(v_i)$$


$$\text{Solve } AU = L \quad \begin{cases} A_{ij} = a(v_j, v_i) \\ L_i = l(v_i) \end{cases}$$

discrete


How to lift the degeneracy?

Idea n°2

Measuring gravitational redshift
using clocks [under investigation]



Chameleon constraints from MICROSCOPE



M. Pernot-Borràs et al.
(2019)

Chameleon constraints from MICROSCOPE

MICROSCOPE can do:



- weak equivalence principle [state of the art]
- generic long-range Yukawa 5th-force [state of the art]
- light dilaton [competitive]
- Lorentz invariance [state of the art]


MICROSCOPE cannot do:



- generic short-range Yukawa 5th-force [not competitive]
- chameleon 5th-force [not competitive]

MICROSCOPE was not designed for testing short-ranged modified gravity theories. Recent work challenges the claim on the ability of space experiments to detect chameleon-sourced violations of the WEP sourced by the Earth [2, 3].


Convergence Analysis (FEM)



Implemented techniques

- Compactification : $\Omega \rightarrow \tilde{\Omega}$ (global coordinate transform) + BC applied at the boundary of the compactified domain.
- “Connected” : domain splitting $\bar{\Omega} = \bar{\Omega}_{int} \cup \bar{\Omega}_{ext}$ and Kelvin inversion $\Omega_{ext} \rightarrow \tilde{\Omega}_{ext}$ + identification of the boundary DOFs $\partial\Omega_{int} \equiv \partial\tilde{\Omega}_{ext}$.
- “ping-pong” : domain splitting $\bar{\Omega} = \bar{\Omega}_{int} \cup \bar{\Omega}_{ext}$ and Kelvin inversion $\Omega_{ext} \rightarrow \tilde{\Omega}_{ext}$ + iterative method with DtN / NtD transmission conditions at the boundary

Convergence Analysis (FEM)



Building blocks of Scalar-Tensor theories

Metric Tensor

$$g_{\mu\nu}$$

Einstein-Hilbert action in General Relativity

$$S_{EH} = \frac{M_{Pl}^2}{2} \int d^4x \sqrt{-g} R + \int d^4x \sqrt{-g} L_m(g_{\mu\nu}, \psi_m^{(i)})$$

Building blocks of Scalar-Tensor theories

Metric Tensor

$g_{\mu\nu}$

+

Scalar Field

ϕ


Modified action with a scalar field (example)

$$S = \int d^4x \sqrt{-g} \left[\frac{M_{Pl}^2}{2} R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right] + \int d^4x \sqrt{-\tilde{g}} L_m \left(\Omega^2(\phi) g_{\mu\nu}, \psi_m^{(i)} \right)$$

Modified Gravity geodesics


$$u^\mu \nabla_\mu u^\rho = - \frac{\partial \ln \Omega}{\partial \phi} \perp^{\mu\rho} \partial_\mu \phi$$

Starting point



General Relativity geodesics

$$\frac{d^2 x^\sigma}{ds^2} + \Gamma_{\mu\nu}^\sigma \frac{dx^\mu}{ds} \frac{dx^\nu}{ds} = 0$$




$$\mathbf{F}_\phi = -m \frac{\partial \ln \Omega}{\partial \phi} \nabla \phi$$

Modified Gravity geodesics

$$u^\mu \nabla_\mu u^\rho = - \frac{\partial \ln \Omega}{\partial \phi} \perp^{\mu\rho} \partial_\mu \phi$$

Starting point



General Relativity geodesics

$$\frac{d^2 x^\sigma}{ds^2} + \Gamma_{\mu\nu}^\sigma \frac{dx^\mu}{ds} \frac{dx^\nu}{ds} = 0$$

Derivation can be found in M. Pernot-Borràs 2020, PhD thesis