

# *Hydrodynamisation of charm quarks in heavy ion collisions*

Stefan Floerchinger (Jena University)

work together with **Federica Capellino**, Andrea Dubla, Eduardo Grossi,  
Andreas Kirchner and Silvia Masciocchi

Strangeness in Quark Matter, Strasbourg  
June 06, 2024.




## Fluid dynamics



- long distances, long times or strong enough interactions
- quantum fields form a fluid!
- needs **macroscopic** fluid properties
  - thermodynamic equation of state  $p(T, \mu)$
  - shear and bulk viscosity  $\eta, \zeta$
  - heat conductivity
  - relaxation times
  - heavy quark diffusion coefficient  $\kappa_n$
- fixed by **microscopic** properties encoded in Lagrangian  $\mathcal{L}_{\text{QCD}}$

# High energy nuclear collisions



## Relativistic fluid dynamics

**Energy-momentum tensor** and conserved current

$$T^{\mu\nu} = \epsilon u^\mu u^\nu + (p + \pi_{\text{bulk}})\Delta^{\mu\nu} + \pi^{\mu\nu}$$

$$N^\mu = n u^\mu + \nu^\mu$$

- tensor decomposition using fluid velocity  $u^\mu$ ,  $\Delta^{\mu\nu} = g^{\mu\nu} + u^\mu u^\nu$
- thermodynamic equation of state  $p = p(T, \mu)$

Covariant **conservation laws**  $\nabla_\mu T^{\mu\nu} = 0$  and  $\nabla_\mu N^\mu = 0$  imply

- equation for energy density  $\epsilon$
- equation for fluid velocity  $u^\mu$
- equation for particle number density  $n$


Need **further evolution equations** [e.g Israel & Stewart]

- equation for shear stress  $\pi^{\mu\nu}$
- equation for bulk viscous pressure  $\pi_{\text{bulk}}$


$$\tau_{\text{bulk}} u^\mu \partial_\mu \pi_{\text{bulk}} + \dots + \pi_{\text{bulk}} = -\zeta \nabla_\mu u^\mu$$

- equation for diffusion current  $\nu^\mu$
- non-hydrodynamic degrees of freedom are needed for relativistic causality!

# Thermodynamics of QCD



[Borsányi *et al.* (2016), similar Bazavov *et al.* (2014)]



[Floerchinger, Grossi, Lion (2019)]


- equation of state at vanishing chemical potential is well known now
- at large temperature lattice QCD
- at small temperature hadron resonance gas approximation
- extensions to non-zero chemical potentials e. g. by Taylor expansion

## *Flow in heavy ion collisions*

### **FluidM:** Fluid dynamics of heavy ion collisions with Mode expansion

[Floerchinger & Wiedemann, PLB 728, 407 (2014), PRC 88, 044906 (2013), 89, 034914 (2014)]

[Floerchinger, Grossi & Lion, PRC 100, 014905 (2019)]




- background-fluctuation splitting + mode expansion
- analogous to cosmological perturbation theory
- substantially improved numerical performance (pseudospectral method)
- resonance decays included  
[Mazeliauskas, Floerchinger, Grossi & Teaney, EPJC 79, 284 (2019)]
- allows fast and precise comparison between theory and experiment

# Parameter estimation from theory-experiment comparison


[Vermunt, Seemann, Dubla, Floerchinger, Grossi, Kirchner, Masciocchi, Selyuzhenkov, PRC 108, 064908 (2023)]

- fluid models have parameters
- can be determined with Bayesian analysis from data
- here based on transverse momentum spectra of pions, kaons, protons
- data from Pb-Pb (2.76 TeV), Pb-Pb (5.02 TeV), Xe-Xe (5.44 TeV)



# Particle production at the Large Hadron Collider

[Vermunt, Seemann, Dubla, Floerchinger, Grossi, Kirchner, Masciocchi, Selyuzhenkov, PRC 108, 064908 (2023)]



- overall good description
- some deviations for pions at small  $p_T$



## *Fluid dynamics for heavy quarks from Fokker-Planck equation*

- phase-space distribution function  $f(t, \mathbf{x}, p)$
- currents are moments with respect to momenta

$$N^\mu(t, \mathbf{x}) = \int \frac{d^3 p}{(2\pi)^3 p^0} p^\mu f(t, \mathbf{x}, p)$$

- Boltzmann equation for time evolution

$$p^\mu \frac{\partial}{\partial x^\mu} f(t, \mathbf{x}, p) = C[f]$$

- heavy quarks get small “momentum kicks” from light partons
- Fokker-Planck approximation to collision kernel

$$C[f] = k^0 \frac{\partial}{\partial p^j} \left[ A^j f + \frac{\partial}{\partial p^k} \left[ B^{jk} f \right] \right]$$

- fluid dynamics from taking moments of the Fokker-Planck equation
- approximations justified for slow dynamics

## Equations of motion for charm current

- net heavy quark number current  $N_-^\mu = N_Q^\mu - \bar{N}_Q^\mu$  conserved in QCD but not in electroweak theory
- total integrated net quark number vanishes
- average quark number current  $N_+^\mu = (N_Q^\mu + \bar{N}_Q^\mu)/2$  approximately conserved for small temperatures  $T \ll m_Q$
- we work with

$$N^\mu = N_+^\mu = nu^\mu + \nu^\mu$$

- conservation law


$$\nabla_\mu N^\mu = u^\mu \partial_\mu n + n \nabla_\mu u^\mu + \nabla_\mu \nu^\mu = 0$$

- additional equation of motion

$$\tau_n \Delta^\rho{}_\sigma u^\lambda \nabla_\lambda \nu^\sigma + \nu^\rho + \kappa_n \Delta^{\rho\sigma} \partial_\sigma \left( \frac{\mu}{T} \right) = 0$$

- chemical potential  $\mu$  conjugate to heavy quark number
- heavy quark diffusion coefficient  $\kappa_n = D_s n$
- relaxation time  $\tau_n$

[Floerchinger & Grossi, JHEP 08 (2018) 186]



- dissipative fluid equations *can* be of hyperbolic type
- characteristic velocities depend on fluid fields
- need  $|\lambda^{(j)}| < c$  for relativistic causality
- works when relaxation times are large enough


## *Thermodynamic equation of state for charm*

- fluid dynamics needs a thermodynamic equation of state
- dependence of pressure on charm chemical potential not very well known
- we use a hadron resonance model approximation with sum over all measured charmed states

$$n(T, \mu) = \frac{T}{2\pi^2} \sum_{i \in \text{HRG}_c} q_i M_i^2 \exp\left(\frac{q_i \mu}{T}\right) K_2\left(\frac{M_i}{T}\right)$$

- yields larger values than gas of free charm quarks
- lattice results would be nice to have

## Constraints on charm quark diffusion on the lattice





- latest lattice results for heavy quark diffusion coefficient for  $N_f = 2 + 1$  flavor QCD indicate small  $D_s$  [HotQCD, PRL 130, 231902 (2023)]
- supports fast hydrodynamization of heavy quarks
- phenomenological analysis based on different transport models [ALICE, JHEP01(2022)174] and Bayesian analysis based on Langevin dynamics [Xu, Bernhard, Bass, Nahrgang, Cao, PRC 97, 014907 (2018)] support larger values of  $D_s$

## Applicability of fluid description

[Capellino, Beraudo, Dubla, Floerchinger, Masciocchi, Pawlowski, Selyuzhenkov, PRD 106, 034021 (2022)]

- Fokker-Planck equation yields relation for relaxation time  $\tau_n$  in terms of diffusion coefficient  $D_s$
- fluid dynamics applicable when the relaxation time is small compared to the dynamics
- for initial Bjorken expansion

$$\tau_n < 1/(\nabla_\mu u^\mu) = \tau$$



## *Initial conditions for charm current*

[Capellino, Dubla, Floerchinger, Grossi, Kirchner, Masciocchi, PRD 108, 116011 (2023)]

- initial density distribution from hard scattering

$$n(\tau_0, r) = \frac{1}{\tau_0} n_{\text{coll}}(r) \frac{1}{\sigma_{pp}^{\text{in}}} \frac{d\sigma^{Q\bar{Q}}}{dy}$$

$$\sigma_{pp}^{\text{in}} = 67.6 \text{ mb}, \quad \frac{d\sigma^{Q\bar{Q}}}{dy} = 0.463 \text{ mb} \text{ [Cacciari, Frixione, Nason, JHEP03(2001)006]}$$


- diffusion current initially assumed to vanish

$$\nu^\mu(\tau_0, r) = 0$$

- leads to parameter-free model for initial charm density and current

## Evolution of charm density and diffusion current

[Capellino, Dubla, Floerchinger, Grossi, Kirchner, Masciocchi, PRD 108, 116011 (2023)]




- Charm density expands and dilutes like energy density
- diffusion leads to further dilution



## Yields of charmed hadrons


[Capellino, Dubla, Floerchinger, Grossi, Kirchner, Masciocchi, PRD 108, 116011 (2023)]



- resonance decays from FASTRESO sizeable
- yield of  $\Lambda_c^+$  underpredicted, possibly missing higher resonances in PDG list?
- prediction for  $\Xi_c^+$  and  $\Omega_c^0$

## Transverse momentum spectra of charmed hadrons


[Capellino, Dubla, Floerchinger, Grossi, Kirchner, Masciocchi, PRD 108, 116011 (2023)]



- good agreement for D-mesons up to  $p_T \approx 4 - 5$  GeV
- some deviations for  $J/\Psi$  (dissipative correction?)





# Predictions for yields of multicharmed hadrons

[Capellino, PhD thesis (2024)]



# Predictions for transverse momentum spectra of multicharmed hadrons

[Capellino, PhD thesis (2024)]



## Conclusions

- fluid dynamic description for heavy quark currents
- no *chemical* equilibration assumed
- local *kinetic* equilibrium picture is useful
- on-top description: fluid velocity and temperature governed by QCD fluid with equation of state for 2+1 light flavors
- spectra of mesons and baryons with charm quarks well described up to transverse momenta of  $p_T \approx 4$  GeV
- total abundances depend on feed-down from resonance decays
- extension to bottom quark current should be attempted
- dissipative corrections at freeze-out seem small but should be studied