



UNIVERSITETET
I OSLO




First $D^0 + \bar{D}^0$ measurement in heavy-ion collisions at SPS energies with NA61/SHINE

Anastasia Merzlaya (UiO)
(for the NA61/SHINE Collaboration)

SQM24
04.06.2023


Motivation of open charm measurements

- Heavy quarks are produced in the hard scattering processes that occur in the early stage of the collision between partons of the incoming nuclei;
- By studying charm hadrons one can get insight into properties of the medium created in the collision;
- Such measurements can be in a big interest at the SPS energies, close to the threshold of QGP creation.
- There are no measurements of open charm in A+A collisions at SPS.




Models of charm production

- Predictions of charm yield differ by up to two orders of magnitude for central heavy-ion collisions at the top SPS energy (beam momentum $150A$ GeV/c, $\sqrt{s_{NN}} = 16.8$ GeV);



- Obtaining precise data on $D^0 + \bar{D}^0$ is expected to narrow the spectrum of viable theoretical models and thus learn about the charm quark and hadron production mechanisms.
 - **HSD:** Hadron-String Dynamics
O. Linnyk et al. Int. J. Mod. Phys. E 17 (2008), 1367-1439
 - **pQCD:** the scaled PYTHIA calculations
P. Braun-Munzinger et al. Phys. Lett. B 490 (2000), 196-202
 - **HRG:** Hadron Resonance Gas Model
M. I. Gorenstein et al. J.Phys.G 27 (2001) L47-L52
 - **Statistical Quark Coalescence:**
M. I. Gorenstein et al. Phys.Lett.B 509 (2001) 277-282
 - **Dynamical Quark Coalescence:** ALCOR and MICOR models extended to charm formation.
P. Levai et al. J.Phys.G 27 (2001) 703-706
 - **SMES:** A statistical model of the early stage
M. Gazdzicki et al., Acta Phys. Polon. B 30 (1999), 2705



p+A measurements of charm at low $\sqrt{s_{NN}}$



- Open charm measurements in π +A and p+A data from SPS and Fermilab experiments show:
 - PYTHIA reasonably describes energy dependence;
 - PYTHIA underestimates the $D^0 + \bar{D}^0$ production cross-section by the **factor 3-3.5**.


NA38/NA50 & NA60

- Indirect estimation of open charm yield using dimuons from semi-leptonic charm quark pair decays by NA50 and NA60;
- Open charm contribution was separated via the fit procedure from an inclusive dimuon distribution, which also contains charmonium and Drell-Yan components;
- Centrality-dependent scaling factor for open charm production in PYTHIA is needed to reproduce the di-muon background in the intermediate mass range.




Anomalous J/ψ suppression

- For central heavy-ion collisions ($N_{\text{part}} \sim 200$) anomalous J/ψ suppression is observed in In+In and Pb+Pb collisions by NA60;
- It was initially attributed to onset of QGP formation in nuclear collisions, however CNM explanations have been proposed:
 - Shadowing;
 - Nuclear absorption.
- Open charm measurements would provide another view to the anomalous J/ψ suppression observed by NA60.






The NA61/SHINE experiment at CERN SPS



Small Acceptance Vertex Detector

- Four stations at $z = 5, 10, 15$ and 20 cm;
- 16 MIMOSA sensors with pixel size $18.4 \times 18.4 \mu\text{m}^2$;
- Primary vertex resolution $\sigma_{x,y} \approx 1 \mu\text{m}$, $\sigma_z = 15 \mu\text{m}$;
- Secondary vertex resolution $\sigma_{x,y} \approx 10 \mu\text{m}$, $\sigma_z = 170 \mu\text{m}$ for D^0 and \bar{D}^0 .



- The SAVD acceptance for $D^0 + \bar{D}^0$ is

$$-0.5 < y < 1.0$$


$$0.2 < p_T < 2.0 \text{ GeV}/c$$

(analysis is performed in this single bin)




Event and track selection

- Event selection:
 - Data taking of **Xe+La 150A GeV/c** in 2017;
 - 1.93M **0-20%** central events;
- Track selection:
 - 3 or 4 SAVD hits (→ spatial resolution)
 - ≥ 10 TPC hits (→ momentum resolution)
 - No PID was applied.


$D^0 + \bar{D}^0$ reconstruction



| cut | value |
|---|-----------------------------|
| (a) Impact parameter of daughter d | $>36 \mu\text{m}$ |
| (b) Distance between primary and secondary vertices \mathbf{V}_z/γ | $>0.15 \text{ mm}$ |
| (c) Impact parameter of parent D | $<20 \mu\text{m}$ |
| Distance of closest approach DCA | $<42 \mu\text{m}$ |
| Parent momentum | $13 < p < 38 \text{ GeV}/c$ |





$D^0 + \bar{D}^0$ signal in 0-20% Xe+La at 150A GeV/c



- $K\pi$ -invariant mass distribution for 0-20% Xe+La at 150A GeV/c;
- This is the first direct observation of $D^0 + \bar{D}^0$ signal at the SPS energies with significance better than 5;


Simulations in GEANT4

- For obtaining the corrections the simulation in GEANT4 was performed:
 - The background was described using the EPOS model;
 - The signal phase space was parametrized using 3 models;
 - The yield of $D^0 + \bar{D}^0$ from the models not used;
 - Parametrized signal is used to enrich background event.



Visible yield of $D^0 + \bar{D}^0$ in 0-20% Xe+La at $150A$ GeV/ c

| correction with: | $N_{visible}(D^0 + \bar{D}^0)$ |
|------------------|--------------------------------|
| AMPT | 0.184 ± 0.032 (stat) |
| PHSD | 0.204 ± 0.036 (stat) |
| PYTHIA/Angantyr | 0.201 ± 0.035 (stat) |



| $N_{visible}(D^0 + \bar{D}^0)$ |
|--|
| 0.196 ± 0.035 (stat) ± 0.051 (syst) |

$$\begin{aligned} -0.5 < y < 1.0 \\ 0.2 < p_T < 2.0 \text{ GeV}/c \end{aligned}$$


- Systematic uncertainties include:
- Model-dependent phase space;
 - Track quality cut selection;
 - Spatial cuts selection;
 - Signal extraction procedure;
 - Background fitting procedure.

$\langle D^0 + \bar{D}^0 \rangle$ and dN/dy in 0-20% Xe+La at 150A GeV/c

| correction with: | $\frac{dN(D^0 + \bar{D}^0)}{dy}$ for $-0.5 < y < 1.0$ | Yield in 4π $\langle D^0 + \bar{D}^0 \rangle$ |
|------------------|---|--|
| AMPT | 0.129 ± 0.023 (stat) ± 0.035 (syst) | 0.218 ± 0.039 (stat) ± 0.060 (syst) |
| PHSD | 0.148 ± 0.026 (stat) ± 0.036 (syst) | 0.303 ± 0.054 (stat) ± 0.074 (syst) |
| PYTHIA/Angantyr | 0.147 ± 0.026 (stat) ± 0.037 (syst) | 0.300 ± 0.052 (stat) ± 0.075 (syst) |

- Extrapolation factors for AMPT significantly differ from PHSD and PYTHIA/Angantyr due to different phase space distribution of $D^0 + \bar{D}^0$:
 - AMPT: 84.1% of all $D^0 + \bar{D}^0$ are in the selected $y - p_T$ bin
 - PHSD: 67.4%
 - PYTHIA/Angantyr: 66.9%

Estimation of $\langle D^0 + \bar{D}^0 \rangle$ for Xe+La from p+A data




- One can estimate $D^0 + \bar{D}^0$ yield for Xe+La at $\sqrt{s_{NN}} = 16.8$ GeV from the extrapolation of π +A and p+A data:

$$\langle D^0 + \bar{D}^0 \rangle = 2 \times \frac{\sigma_{D^0 + \bar{D}^0}}{\sigma_{inelastic \text{ for } p+p = 31 \text{ mb}}} \times N_{coll} = 0.46$$

(the value was cross-checked with recent PYTHIA calculations, which shows similar result)

Comparison of $\langle D^0 + \bar{D}^0 \rangle$ with models



- Red band indicates theoretical uncertainty of the obtained result.
- Precision of the data is sufficient to discriminate between extreme model predictions;
- The dynamical microscopic models (AMPT, Pythia, PHSD) significantly underestimate $\langle D^0 + \bar{D}^0 \rangle$ while ALCOR and SMES overestimate it.
- The obtained results are below p+A extrapolation at the level of $\sim 2\sigma$:
 - Imprecision of the extrapolated p+A cross-section;
 - Fragmentation in A+A vs p+A?
 - N_{coll} vs N_{par} scaling?

Results and discussion

SMES: scaled from 0-20% Pb+Pb using $N_{\text{part}} = 173.7/272.5$


ALCOR (Dynamical Quark Coalescence): scaled from *J.Phys.G* 27 (2001) 703-706 using $N_{\text{coll}} = 331.1/598.8$

Summary & Outlook

- The first direct open charm observation in heavy-ion collisions at the SPS energies was done for Xe+La 0-20% central collisions at 150A GeV/c.
- Precision of the obtained result is sufficient to disentangle between theoretical models.
- After LS2, an upgraded version of Vertex Detector based on ALPIDE sensors designed for ALICE ITS was installed;
- Together with the upgrade of the TPC readout electronics and DAQ it allows data taking with 1 kHz rate.
- In 2022-2023 NA61/SHINE collected ~180M Pb+Pb events at 150A GeV/c.
The data should allow:
 - the p_T and y differential measurements of D^0 and \bar{D}^0 ;
 - measurements of other charm hadrons;→ Better insight into charm production mechanisms at energies close to production threshold.

Thank you for your attention!


The NA61/SHINE experiment



- Beam detectors
 - Position
 - Charge
 - Time
- of beam particles

The NA61/SHINE experiment at CERN SPS

The NA61/SHINE experiment



Beam detectors

- Position
- Charge
- Time

of beam particles

TPCs


- Momentum
- Charge
- PID (dE/dx)

of produced particles

ToF

- PID (mass)

The NA61/SHINE experiment



Beam detectors

- Position
- Charge
- Time

of beam particles

TPCs

- Momentum
- Charge
- PID (dE/dx)

of produced particles

ToF



- PID (mass)

PSD calorimeter


- energy

of projectile
spectators


Vertex Detector



- Main purpose of the **Vertex Detector** is the improvement of track resolution near the interaction point to allow reconstruction of secondary vertices;
- **SAVD** is positioned between the target and the VTPC-1;
- Four planes of coordinate-sensitive detectors are located at 5, 10, 15 and 20 cm distance from the target.





Main Vertex Detector components



- **MIMOSA-26AHR**
 - ❑ 1152x576 pixels of $18.4 \times 18.4 \mu\text{m}^2$
 - ❑ 3.5 μm resolution, 0.05% X0
 - ❑ Readout time: 115.2 μs , 50 μm thinPICSEL Group, IPHC Strasbourg
- **ALICE ITS ladder**
 - ❑ Ultra light carbon fibre
 - ❑ < 0.3% X0 including water coolingSt. Petersburg, CERN
- **CBM Micro Vertex Detector Prototype**
 - ❑ Sensor integration
 - ❑ Flex print cables, Front-end boards
 - ❑ Read-out based on TRB3 FPGA BoardGoethe Universitet Frankfurt am Main


dE/dx




Daughters of $D^0 + \overline{D}^0$ are in the range of 5-30 GeV/c, where π and K dE/dx curves overlap. So, no dE/dx used

$D^0 + \bar{D}^0$ in $y - p_T$ bin


AMPT




PHSD



PYTHIA/
Angantyr



Comparison $N_{4\pi}(c\bar{c})$ with models



- Values for the models are scaled from Pb+Pb 0-20%;
- Values for data scaled from $\langle D^0 + \bar{D}^0 \rangle$ using ratio provided by event generators:

| Ratio of $c\bar{c}$ decaying into $(D^0 + \bar{D}^0)$ | |
|---|-----|
| PHSD | 62% |
| PYTHIA/Angantyr | 35% |

- Comparison of the data and models show significant discrepancy between them: while some models (SMES) is overestimating the charm yield, other (HSD) underestimating it.
- The closest model predictions to obtained result are HRG and Quark Coalescence.

Results and discussion

Scaling with $N_{\text{coll}} = 331.1/598.8$
 Scaling with $N_{\text{part}} = 173.7/272.5$

[---] Stat uncertainty
 [---] Syst uncertainty