




Thermal dielectron measurement in Au+Au collisions with STAR BES-II data

Zhen Wang (王 楨) for the STAR collaboration
Shandong University

The 21st International Conference on Strangeness in Quark Matter
3-7 June 2024, Strasbourg, France



QCD phase diagram



STAR, arXiv: 2402.01998


Deconfined QCD matter produced at extreme high temperatures and/or baryon densities

Study phase structure in the $T - \mu_B$ plane

A “Little Bang” in heavy ion collisions

Relativistic Heavy-Ion Collisions

made by Chun Shen



Deconfined QCD matter produced at extreme high temperatures and/or baryon densities

Study phase structure in the $T - \mu_B$ plane

Extract the information from the final detected particles



C.Shen <https://u.osu.edu/vishnu/2014/08/06/sketch-of-relativistic-heavy-ion-collisions>

How to measure temperature


Hadrons
yields, p_T spectra

Hadrons:

- ✓ Large yields
- ✓ Infer QGP properties when the hadrons decouple
- ✓ Extract temperatures of chemical and kinetic freeze-out, T_{ch} and T_{kin}



How to measure temperature



Photons

p_T spectra



Dileptons

M_{ll} spectra


Electromagnetic Probes:

- ✓ Emitted from early stage to final stage
- ✓ Minimal interaction with medium

How to measure temperature



STAR: Phys.Lett.B 770 (2017), 451-458



Photons

p_T spectra

Dileptons

M_{ll} spectra

Electromagnetic Probes:

- ✓ Emitted from early stage to final stage
- ✓ Minimal interaction with medium

Photons:

- ✓ Extract T_{eff} from p_T spectra
- ✓ $T_{\text{eff}} \rightarrow T_{\text{QGP}}$: medium effect


Dileptons:

- ✓ Temperature measurement without distortion by medium flow effects
- ✓ Only observable to directly access in-medium spectral function


Thermal dileptons

QGP: $M^{3/2} * e^{-M/T}$

In-medium ρ : Relativistic Breit-Wigner * $e^{-M/T}$



Rapp, R. Nat. Phys. 15, 990–991 (2019).



inclusive dileptons

Interested signals


- QGP radiation
- In-medium ρ

Physical backgrounds (hadronic cocktails)

- $\pi^0, \eta, \eta' \rightarrow \gamma e^+ e^-$
- $\omega \rightarrow \pi^0 e^+ e^-$
- $\phi \rightarrow \eta e^+ e^-$
- $\omega, \phi, J/\psi \rightarrow e^+ e^-$
- $c\bar{c} \rightarrow e^+ e^- X$
- Drell-Yan


Invariant mass spectra of thermal dileptons can reveal temperature of the hot medium at both **QGP phase** and **hadronic phase**

STAR experiment and eID




Time Projection Chamber + Time of Flight detectors

- ✓ Charge particle tracking and momentum measurement
- ✓ Electron identification by dE/dx and velocity
- ✓ High purity electron samples



STAR BES-I Dielectron measurements

STAR: Phys.Rev.C 107 (2023) 6, L061901



$\sqrt{s_{NN}} = 200 \text{ GeV}$:


- High statistics
- Open charm dominate the intermediate mass region

$\sqrt{s_{NN}} = 19.6 - 62.4 \text{ GeV}$:


- Statistically limited

BES-II:

- ✓ Statistics ~ 10 times larger than that in the BES-I datasets



Dielectron spectra




←→ **LMR** ←→ **IMR** ←→ **LMR** ←→ **IMR**

Clear enhancement compared to hadronic cocktail in both low mass region (LMR) and intermediate mass region (IMR)

Excess yield


Excess = Data - Cocktail



Excess dilepton spectra in LMR at $\sqrt{s_{NN}} = 27$ and 54.4 GeV Au+Au collisions and at $\sqrt{s_{NN}} = 17.3$ GeV In+In collisions are similar

Temperature extraction from LMR

Excess = Data - Cocktail



Low mass region:

$$(a \cdot BW + b \cdot M^{3/2}) \cdot e^{-M/T}$$

- ✓ Similar T_{LMR} for STAR and NA60 measurements
- ✓ T_{LMR} around the pseudo critical temperature T_{pc} (156 MeV)

LMR

$$T_{\text{LMR}}^{54.4 \text{ GeV}} = 172 \pm 12(\text{stat.}) \pm 18(\text{sys.}) \text{ MeV}$$


$$T_{\text{LMR}}^{27 \text{ GeV}} = 167 \pm 21(\text{stat.}) \pm 18(\text{sys.}) \text{ MeV}$$

$$T_{\text{LMR}}^{17.3 \text{ GeV}} = 165 \pm 4 \text{ MeV}$$

T_{pc} : HotQCD, Phys.Lett.B 795 (2019) 15-21
 NA60: EPJC (2009)59 607-623

Temperature extraction from IMR

Excess = Data - Cocktail



Low mass region:

$$(a \cdot BW + b \cdot M^{3/2}) \cdot e^{-M/T}$$

- ✓ Similar T_{LMR} for STAR and NA60 measurements
- ✓ T_{LMR} around the pseudo critical temperature T_{pc} (156 MeV)

Intermediate mass region:

$$M^{3/2} \cdot e^{-M/T}$$


- ✓ QGP thermal radiation is predicted to be the dominant source in the IMR
- ✓ T_{IMR} is higher than the pseudo critical temperature T_{pc} (156 MeV) supporting that the emission is predominantly from deconfined partonic phase

$$T_{IMR}^{54.4 \text{ GeV}} = 303 \pm 59(\text{stat.}) \pm 28(\text{sys.}) \text{ MeV}$$

$$T_{IMR}^{27 \text{ GeV}} = 280 \pm 64(\text{stat.}) \pm 10(\text{sys.}) \text{ MeV}$$

T_{PC} : HotQCD, Phys.Lett.B 795 (2019) 15-21; NA60: EPJC (2009)59 607-623

Temperature v.s. μ_B



NA60: EPJC (2009) 59 607–623
 HADES: Nature Physics 15, 1040-1045 (2019)
 T_{ch} SH: P. Braun-Munzinger et al. Nature 561, 321-330 (2018)
 T_{ch} GCE/SCE: STAR PRC 96, 044904 (2017)
 T_{pc}: HotQCD, Phys.Lett.B 795 (2019) 15-21

Thermal dielectrons in LMR:


- ✓ T_{LMR} is close to the T_{pc} and T_{ch}
- ✓ Emitted from the hadronic phase, dominantly around the phase transition

Thermal dielectrons in IMR:

- ✓ T_{IMR} is higher than T_{LMR}, T_{pc} and T_{ch}
- ✓ Emitted from the partonic phase

T_{ch} : Chemical freeze-out temperature
 T_{pc} : Pseudo critical temperature

Temperature v.s. μ_B



NA60: EPJC (2009) 59 607–623
 HADES: Nature Physics 15, 1040-1045 (2019)
 Tch SH: P. Braun-Munzinger et al. Nature 561, 321-330 (2018)
 Tch GCE/SCE: STAR PRC 96, 044904 (2017)
 T_{pc} : HotQCD, Phys.Lett.B 795 (2019) 15-21

Thermal dielectrons in LMR:

- ✓ T_{LMR} is close to the T_{pc} and T_{ch}
- ✓ Emitted from the hadronic phase, dominantly around the phase transition

Thermal dielectrons in IMR:

- ✓ T_{IMR} is higher than T_{LMR} , T_{pc} and T_{ch}
- ✓ Emitted from the partonic phase


T_{ch} : Chemical freeze-out temperature

T_{pc} : Pseudo critical temperature

BES-II:

- ✓ Probing the thermal dilepton in the high baryon density region

Dielectron spectra for $\sqrt{s_{NN}} \leq 19.6$ GeV




BES-II collision mode at $\sqrt{s_{NN}} = 7.7 - 19.6$ GeV

- ✓ First dielectron measurements under $\sqrt{s_{NN}} = 19.6$ GeV at RHIC
- ✓ New 9.2 GeV dielectron spectra were obtained
- ✓ Excess observed in low mass region

BES-I V.S. BES-II

Excess = Data - Cocktail




Excess yield invariant mass spectra at $\sqrt{s_{NN}} = 19.6$ GeV

- ✓ BES-I and BES-II results are consistent
- ✓ Much better statistical and systematic uncertainties at BES-II than BES-I
 - ✓ Total error reduced by a factor of ~ 4
 - ✓ Better precision for extended analysis

BES-II @ 7.7 – 19.6 GeV

Excess = Data - Cocktail

Thermal dielectron spectra with STAR BES-II




Mass spectra of excess dielectron yields at different $\sqrt{s_{NN}} = 7.7 - 19.6$

- ✓ Different environment for medium interactions
 - ✓ Baryon chemical potential
 - ✓ Temperature

BES-II @ 7.7 – 19.6 GeV

Normalized excess yield



Mass spectrum of excess dielectron at different $\sqrt{s_{NN}}$

- ✓ Different environment for medium interactions
 - ✓ Baryon chemical potential
 - ✓ Temperature
- ✓ Integrated excess yield
 - ✓ Normalized by π^0 yield
 - ✓ New result @ 9.2 GeV
 - ✓ Hint of decreasing trend below 19.6 GeV

R. Rapp, Phys. Rev. C 63, 054907 (2001)

H. van Hees and R. Rapp, Phys. Rev. Lett. 97, 102301 (2006)


2024/6/4

Zhen Wang @ SQM 2024

18


Dielectron measurements with STAR BES-II and FXT program

STAR, arXiv: 2402.01998



NA60: EPJC (2009) 59 607–623
 HADES: Nature Physics 15, 1040-1045 (2019)
 T_{ch} SH: P. Braun-Munzinger et al. Nature 561, 321-330 (2018)
 T_{ch} GCE/SCE: STAR PRC 96, 044904 (2017)

- ✓ BES-II and FXT data will cover the large gap between the STAR and HADES data
- ✓ The normalized integrated excess yields in mass window $0.4 < M_{ee} < 0.75 \text{ GeV}/c^2$ were obtained
- ✓ Working on T extraction with BES-II data



Summary

Temperature @ 27 & 54.4 GeV:

- ✓ T_{LMR} : Close to T_{pc} and T_{ch} , provide a new insight to study the phase transition temperature
- ✓ T_{IMR} : Higher than T_{pc} and T_{ch} , first QGP temperature at RHIC

Thermal dielectron yields with BES-II:

- ✓ High precision measurement compared to BES-I measurements
- ✓ New thermal dielectron measurements at $\sqrt{s_{NN}} = 9.2$ GeV presented
- ✓ Excess yield spectra for different T and μ_B

Summary

Temperature @ 27 & 54.4 GeV:

- ✓ T_{LMR} : Close to T_{pc} and T_{ch} , provide a new insight to study the phase transition temperature
- ✓ T_{IMR} : Higher than T_{pc} and T_{ch} , first QGP temperature at RHIC

Thermal dielectron yields with BES-II:

- ✓ High precision measurement compared to BES-I measurements
- ✓ New thermal dielectron measurements at $\sqrt{s_{NN}} = 9.2$ GeV presented
- ✓ Excess yield spectra for different T and μ_B

Thanks for your attention!